arXiv:math/9802011v2 [math.AG] 5 Feb 1998

NEW PERIOD MAPPINGS FOR PLANE CURVE SINGULARITIES

RAINER H. KAENDERS

Abstract. For plane curve singularities we construct a mixed Hodge structure (MHS) over $\mathbb{Z}$ on the fundamental group of the Milnor fiber. The concept *nearby fundamental group* is introduced and we develop a theory of iterated integrals along elements of this group. We give an example for which the MHS on the nearby fundamental group detects a modulus which is invisible for the MHS on the vanishing cohomology.

Contents

Introduction 1
1. The Nearby Fundamental Group 6
2. Line Integrals on the Nearby Fundamental Group 20
3. Iterated Integrals on the Nearby Fundamental Group 34
4. Well-Definedness and Formal Connections 46
5. Plane Curve Singularities 66
References 74

Introduction

The building block for many invariants of plane curve singularities like the monodromy, the intersection form, the Seifert form, Coxeter-Dynkin diagrams, the mixed Hodge structure on the vanishing cohomology and the spectral pairs is the first integral homology group of the Milnor fiber or – which is the same – the abelianized fundamental group of the Milnor fiber.

The restriction to the abelianized version of this fundamental group causes some serious limitations. E. g. Lê Dũng Tráng [Lê72] proved that for irreducible plane curve singularities the monodromy (on homology) has finite order. However A'Campo [A'C73] was the first to discover that the monodromy on the fundamental group of the Milnor fiber (where the one boundary component is collapsed to one point, the basepoint) has infinite order, whenever the singularity has at least two Puiseux pairs. This shows that the monodromy on the fundamental group is richer than on homology.

For the study of analytical invariants of plane curve singularities, the first (co)homology group is of particular interest because of Hodge theory.

1991 *Mathematics Subject Classification.* Primary 14B05, 14H20; Secondary 32S35, 14F35 .

This work was partly supported by grant ERBCHBICT930403 (HCM) from the Eurpean Community and The Netherlands Organisation for Scientific Research (NWO).

Deligne [Del71] introduced in 1970/71 the notion of mixed Hodge structure (MHS) and showed that the cohomology of any complex algebraic variety carries such a MHS in a natural way. Schmid [Sch73] and Steenbrink [Ste76] determined a limit-MHS of a variation of Hodge structures coming from families of compact algebraic manifolds. This provided the background for the definition of the MHS on the vanishing cohomology of hypersurface singularities by Steenbrink [Ste77]. In the case of plane curve singularities this is a MHS on the first cohomology group of the Milnor fiber.

However, due to the fact that for irreducible plane curve singularities the monodromy on homology is finite, families of mutually right-left – but not necessarily right-equivalent plane curve singularities all have isomorphic MHSs on the vanishing cohomology. This holds, for example, for the family

$$f_\lambda(x,y) = \left(y^2 - x^3\right)^2 - \left(4\lambda x^5 y + \lambda^2 x^7\right), \quad \lambda \neq 0, \tag{0.1}$$

which is the singularity with Puiseux expansion

$$y = x^{\frac{3}{2}} + \sqrt{\lambda}\, x^{\frac{7}{4}}.$$

In 1987 Hain [Hai87a], [Hai87c] used Chen's iterated integrals to generalize Deligne's construction of a MHS on cohomology to a construction of a MHS on the homotopy Lie algebra for any complex algebraic variety with base point. In particular, this object contains a MHS which is called *the MHS on the fundamental group*. Already before, in 1978, Morgan [Mor78] had used Sullivan's minimal models to put a MHS on the homotopy Lie algebra of *smooth* complex algebraic varieties.

Hain also showed, in [Hai87b], that the local system of homotopy groups associated with a topologically trivial family of smooth pointed varieties underlies a good variation of $\mathbb{Q}$-MHSs, which implies by the earlier work of Steenbrink and Zucker [SZ85] that there is a limit $\mathbb{Q}$-MHS.

This provides the background of the problem[1] we study in this paper: Is there a generalization of the MHS on the vanishing (co)homology to something like 'a MHS on the fundamental group of the general Milnor fiber' in the case of plane curve singularities?

In this article we show that the construction of such a generalization is possible and we will call it *the mixed Hodge structure on the nearby fundamental group of a plane curve singularity*. We develop a theory that allows us to take iterated integrals along certain paths in the central fiber. These paths mimic paths in the regular fiber. The given approach seems to be even new for the integration of elements in the vanishing cohomology. Due to this integration theory, the constructed MHS is defined over the integers. For instance in the family f_λ defined on page 2 this $\mathbb{Z}$-MHS detects the modulus. The above formulated problem is the guideline of the paper.

We consider first the situation of a degenerating family $h : (Z, D^+) \to (\Delta, 0)$ of compact Riemann surfaces with r boundary components over the disk $\Delta \subset \mathbb{C}$, where the singular fiber, the fiber over 0, is a divisor with normal crossings $D = D^+ \cup D_0 \cup \cdots \cup D_{r-1}$. Its components are compact closed Riemann surfaces, except for the r components $D_0, \ldots, D_{r-1}$, which are disks. Such a degenerating family can be constructed from a plane curve singularity f by a process that is called

---

[1] This problem was proposed to us by J. H. M. Steenbrink.

'semistable reduction' (see 5.1). Each of the disks $D_0, \ldots, D_{r-1}$ corresponds to a branch of f.

In this situation, we want to mimic a path in a regular fiber by a path in the central fiber. Let us give a finite version of the ideas that we develop in an infinitesimal way in Chapter 1. Intuitively it is clear that the difficulties for this mimicry arise at the double points of the divisor with normal crossings D, which are all given locally by an equation: $x \cdot y = t$. If $t \neq 0$, this equation gives within the central fiber (locally) a 1-1 relation between points off the double point in one component and points off the double point in the other. This is obviously not true for $t = 0$. For $t \neq 0$, we consider then paths γ that approach the double point in one component and leave it in the other component in such a way that locally for all $\varepsilon > 0$ holds:

$$x \circ \gamma(\tau_0 - \varepsilon) \cdot y \circ \gamma(\tau_0 + \varepsilon) = t,$$

where $\gamma(\tau_0)$ is the double point. By the identification that is sketched in Figure 1, these paths can be thought of as being in the fiber over t.



FIGURE 1. Identification of a path in the regular fiber of the map $(x, y) \mapsto x \cdot y = t$ with a path in the singular fiber.

In this picture one can also observe that if $|t|$ is big, the path in the central fiber has to make a big detour compared to the corresponding path in the regular fiber. Therefore, the speed with which the path in the central fiber approaches and leaves the double point is a measure for the distance between the central fiber and the regular fiber in which the 'regular path' lives. We give an infinitesimal version of this idea in 1.2. Given a non zero tangent vector $\vec{v}$ in the tangent plane $T_0\Delta$ of the base space Δ and a non zero tangent vector $\vec{w}$ in the tangent plane $T_{p_0}D_0$ at the disk D_0, we introduce the notion of *a path over $\vec{v}$ based at $\vec{w}$*. Moreover, we define a *homotopy over $\vec{v}$ based at $\vec{w}$* or shorter, a *nearby homotopy*, in such a way that the set of all *nearby homotopy classes* becomes a group that we denote by:

$$\pi_1(Z_{\vec{v}}, \vec{w}).$$

The $\{\pi_1(Z_{\vec{v}},\vec{w})\}_{\vec{v}\in(T_0\Delta)^*,\vec{w}\in(T_{p_0}D_0)^*}$ form a local system on $(T_0\Delta)^*\times(T_{p_0}D_0)^*$, which is isomorphic to a corresponding local system that comes from the fundamental groups on the regular fibers (see 1.5 on page 13).

Denote by $\vec{J}=J_{\vec{v},\vec{w}}$ the augmentation ideal of the group ring $\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})$. In the Chapters 2 and 3, we will define for all $s\geq 0$ a $\mathbb{Z}$-MHS on

$$\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})\big/J^{s+1}_{\vec{v},\vec{w}}. \tag{0.2}$$

This will be done by putting a $\mathbb{Z}$-MHS on $(\vec{J}\big/\vec{J}^{s+1})^*$, the dual of $\vec{J}\big/\vec{J}^{s+1}\subset\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})\big/J^{s+1}_{\vec{v},\vec{w}}$.

The main ingredient for the definition of these $\mathbb{Z}$-MHSs is a differential graded algebra (dga) $A^\bullet$, which is made up from differential forms on the central fiber but whose cohomology is isomorphic to the cohomology of the regular fiber. It was suggested to us by Hain[2] to use a dga like $A^\bullet$ and he proposed the problem of finding a way to integrate elements of this dga along some kind of paths in the central fiber. We discovered that the right notion of *path* for this purpose turned out to be the notion of a path over $\vec{v}$ (based at $\vec{w}$). We will show that closed elements in A^1 can be integrated along paths in Z_0 over $\vec{v}$ in a natural way. This yields an integral structure on $H^1(A^\bullet)$, which is by this integration dual to $\vec{J}\big/\vec{J}^{\,2}$. On $A^\bullet$ we have filtrations $W_\bullet$ and $F^\bullet$ such that the induced filtrations together with the integral structure define a $\mathbb{Z}$-MHS on $H^1(A^\bullet)$ (see Theorem 2.11 on page 28).

Since $A^\bullet$ has an *algebra* structure, we can also define *iterated* line integrals along paths over $\vec{v}$. We shall do this in Chapter 3. This will give a way (see 3.9) to describe $\mathrm{Hom}_{\mathbb{Z}}(\vec{J}\big/\vec{J}^{s+1},\mathbb{C})$ in terms of the dga $A^\bullet$, similar to Chen's classical π_1-De-Rham theorem (cf. (4.4) in [Hai87c]).

These iterated integrals will lead to the definition of a $\mathbb{Z}$-MHS on $(\vec{J}\big/\vec{J}^{s+1})^*$, and hence to a $\mathbb{Z}$-MHS on (0.2) (see 3.12). And furthermore, we will see that the variation of $\mathbb{Z}$-MHS for all $s\geq 1$

$$\left\{J_{\vec{v},\vec{w}}\big/J^{s+1}_{\vec{v},\vec{w}}\right\}_{\vec{v}\in(T_0\Delta)^*,\ \vec{w}\in(T_{p_0}D_0)^*}$$

is a nilpotent orbit of MHS (see 2.4 and 3.5).

Let $\pi_1(Z_0,p_0)$ be the fundamental group of the central fiber and let $\mathbb{Z}\pi_1(Z_0,p_0)$ be its group ring with augmentation ideal J_0. We define a MHS on the fundamental group of the central fiber. This MHS is more or less a special case of the general construction of a MHS on the fundamental group of a complex algebraic varietiy as introduced by Hain [Hai87a]. We show that the obvious group homomorphism

$$c:\pi_1(Z_{\vec{v}},\vec{w})\to\pi_1(Z_0,p_0)$$

induces an inclusion of MHSs:

$$c^*:\left(J_0\big/J_0^{s+1}\right)^*\to\left(\vec{J}\big/\vec{J}^{s+1}\right)^*.$$

In Chapter 5 we show how all these considerations can be applied to study plane curve singularities

$$f=f_0\cdots f_{r-1}:(\mathbb{C}^2,0)\to(\mathbb{C},0).$$

Let t be a coordinate on $(\mathbb{C},0)$. The tangent vector $\frac{\partial}{\partial t}$ in $T_0\mathbb{C}$ yields a finite number of tangent vectors in the base space of the 'semistable reduction'. If f_0 is a distinguished branch of f and D_0 the disk corresponding to f_0, then $\frac{\partial}{\partial t}$ defines a finite number of tangent vectors in $T_{p_0}D_0$ in a natural way, which we call *the monstrance of* (f, f_0). We use this to define *the MHS on the nearby fundamental group of a plane curve singularity* f. This MHS depends on the tangent vector $\frac{\partial}{\partial t}$, which is preserved under right-equivalence. It might be interesting to note in this context that if a right-left-equivalence between two plane curve singularities preserves $\frac{\partial}{\partial t}$ that then the singularities are right-quivalent. This is a consequence of a result of J. Briançon and H. Skoda (see 5.8).

---

[2] The dga $A^\bullet$ is in principle a modified version of the complex $P^\bullet_{\mathbb{C}}(X^*)[\theta]$ of [Hai87b] for this special situation with non compact fibers.

Finally we consider the *MHS on the nearby fundamental group* of the example (0.1) on page 2 and show that the extension of H_1 by the part of weight -3 of $\vec{J}/\vec{J}^4$ detects the modulus that is hidden for the vanishing cohomology. Here we focus on the effects due to the infinite order of the monodromy.

In general we expect also interesting information from other parts of this MHS on the nearby fundamental group. Even the extension of H_1 by $H_1 \otimes H_1$ given by $\vec{J}/\vec{J}^3$ might contain interesting information. Also the decomposition of the monodromy into a unipotent and a semisimple part has still to be studied. The ideas described here for plane curve singularities have a counterpart for degenerating families of compact Riemann surfaces with a section (representing the basepoint in each fiber). Mutatis mutandis this is a special case of our construction.

In the first two sections we chose to gather a few technical proofs in appendices in order not to cloud the simplicity of the construction. We recommend to skip these appendices for the first reading.

Acknowledgements: This paper evolved from my thesis which was accepted at the Catholic University of Nijmegen. I am grateful to my advisor, Joseph Steenbrink for his continuous encouragement. I owe very much to many valuable conversations with Richard Hain and David Reed during a stay of three months at Duke University in spring 1996 and during a short visit of one week in spring 1997. As mentioned in the introduction, Richard Hain made some very useful suggestions for the problem I was looking at. Finally I would like to thank many people with whom I had interesting discussions and who gave encouragement during the preparation of my thesis.

1. THE NEARBY FUNDAMENTAL GROUP

In this section we describe our concept of *nearby fundamental group.*

1.1. The Setting.

Before we study plane curve singularities let us consider the folllowing setting. We are going to use this setting throughout the whole first three sections of the article.

Let

$$h : (Z,\ D^+) \longrightarrow (\Delta,\ 0),$$

be a map of space germs, where

- Z is a complex manifold of dimension 2,
- $\Delta = \{z \in \mathbb{C} \big| |z| < 1\}$,
- $Z_0 := h^{-1}(0) = \bigcup_{i\geq 0} D_i$ is a connected reduced divisor with normal crossings (DNC) with components D_i for $i \geq 0$. The components intersect each other mutually in either one or no point. If it exists, the intersection point between D_k and D_l for $k < l$ is denoted by p_{kl}.
- $D^+ := \bigcup_{i>r-1} D_i$ is the union of closed compact Riemann surfaces D_i.
- $D_0, \ldots , D_{r-1}$ are disjoint open disks ($r \geq 1$). Each disk D_i intersects D^+ in one point p_i. Note that each point $p_i \in \{p_0, \ldots , p_{r-1}\}$ is in particular an intersection point of two components, i. e. $p_i = p_{il}\}$ for some $l > r - 1$.
- Let $Z^* := Z \setminus h^{-1}(0)$ and $\Delta^* := \Delta \setminus \{0\}$ as well as $h^* := h_{\big| Z^*}$. Then $h^* : Z^* \longrightarrow \Delta^*$ is a locally trivial C^∞-fibration, where the closure of each fiber (in a bigger representative of (Z, D^+)) is a compact Riemann surface with r boundary components.

Moreover let us assume that we are given two nonzero tangent vectors

$$\vec{v} \in T_0\Delta \qquad \text{and} \qquad \vec{w} \in T_{p_0}D_0.$$

We make the following convention:

"[k<l]" means "all pairs k, l for which $D_k \cap D_l \neq \emptyset$ and $k < l$".
"[a<k<l]" means "all pairs k, l for which $D_k \cap D_l \neq \emptyset$ and $a < k < l$".

For $i \geq 0$ define

$$P_i := \bigcup_{j\neq i} D_i \cap D_j \qquad \text{and} \qquad D_i^* := D_i \setminus P_i.$$

In section 5 we will extract such data like above with additional actions of finite cyclic groups from a plane curve singularity modulo right-equivalence.

Remark 1.1. The construction which we are going to present can similarly be considered in the case $r = 0$. That is, a degenerating family of compact Riemann surfaces, where the role of tangent vector $\vec{w} \in T_{p_0}D_0$ is played by a point $w \in D^+ \setminus \bigcup_{[k<l]}\{p_{kl}\}$. The requirement that two components intersect in at most one point as well as that the components D_i have no self-intersection can easily be dropped. In our exposition it keeps the notation simple and it will be satisfied in the case of curve singularities.

1.2. Fundamental Group of the Nearby Fiber over a Tangent Vector.

Let us explain in this subsection, what we mean by *the fundamental group of the nearby fiber over a tangent vector.*

1.2.1. *The Hessian at a Double Point.* Consider the Hessian of h in a double point $p_{kl} = D_k \cap D_l$. It is a symmetric bilinear form:

$$H(h) : T_{p_{kl}}Z \times T_{p_{kl}}Z \longrightarrow T_0\Delta.$$

Since D_k and D_l intersect transversally we have $T_{p_{kl}}Z = T_{p_{kl}}D_k \oplus T_{p_{kl}}D_l$. Note that $H(h)(T_{p_{ij}}D_i \times T_{p_{ij}}D_i) = 0$ for $\{i,j\} = \{k,l\}$, that is to say that $T_{p_{kl}}D_k$ and $T_{p_{kl}}D_l$ are isotropic subspaces of $H(h)$ in $T_{p_{kl}}Z$. Therefore, the following restriction of the Hessian determines it completely:

$$\eta_{kl} : T_{p_{kl}}D_k \times T_{p_{kl}}D_l \longrightarrow T_0\Delta.$$

Putting all such maps for all double points together, we obtain a map:

$$\langle \cdot, \cdot \rangle : \coprod_{[k<l]} T_{p_{kl}}D_k \times T_{p_{kl}}D_l \longrightarrow T_0\Delta.$$

When we are given local coordinates (x, y) around a double point p_{kl} in Z and a coordinate t on Δ such that in these coordinates

$$h(x, y) = xy = t,$$

then we find

$$\left\langle a\frac{\partial}{\partial x},\, b\frac{\partial}{\partial y} \right\rangle = a \cdot b \,\frac{\partial}{\partial t}.$$

1.2.2. *Paths in Z_0 over $\vec{v}$.* Recall that $\vec{v}$ is a tangent vector in $T_0\Delta$. We define a *path in Z_0 over $\vec{v}$* or a *path in $Z_{\vec{v}}$* to be a continuous map

$$\gamma : [a, b] \longrightarrow Z_0,$$

which, when considered as map to Z, is piecewise smooth and which satisfies the following conditions:

- (i) γ is right- and left differentiable at all double points p_{kl}, $[k < l]$.
- (ii) For each double point p_{kl} with $[0 < k < l]$ there are two possibilities:
  - (a) either γ stays in the same component while passing p_{kl}, or
  - (b) it changes from one component to another.

Assume that $\tau_0 \in [a, b]$ is such that $\gamma(\tau_0) = p_{kl}$. If γ stays in the same component we require (and also for p_0)

$$-\dot{\gamma}^{\leq\tau_0}(\tau_0) = \dot{\gamma}_{\geq\tau_0}(\tau_0) \neq 0$$

and if γ changes from one component to another we impose

$$\left\langle -\dot{\gamma}^{\leq\tau_0}(\tau_0),\ \dot{\gamma}_{\geq\tau_0}(\tau_0) \right\rangle = \vec{v},$$

where $\gamma^{\leq\tau_0} : [a;\tau_0] \to Z_0$ and $\gamma_{\geq\tau_0} : [\tau_0; b] \to Z_0$ denote the respective restrictions of γ. If τ_0 is a resp. b, then we define $\dot{\gamma}^{\leq a}(a) := \dot{\gamma}^{\leq b}(b)$ resp. $\dot{\gamma}_{\geq b}(b) := \dot{\gamma}_{\geq a}(a)$.

1.2.3. *Closed Paths in Z_0 over $\vec{v}$ based at $\vec{w}$.* Let $\vec{w}$ be a tangent vector in $T_{p_0}D_0$. We define *a path over $\vec{v}$ based at $\vec{w}$* to be a path $\gamma : [a, b] \to D^+ \subset Z_0$ over $\vec{v}$ with $\gamma(a) = \gamma(b) = p_0$ such that holds:

$$\langle \vec{w},\, \dot{\gamma}_{\geq a}(a) \rangle = \vec{v} \quad \text{and} \quad \langle -\dot{\gamma}^{\leq b}(b),\, \vec{w} \rangle = \vec{v}.$$

It follows that: $\dot{\gamma}_{\geq a}(a) = -\dot{\gamma}^{\leq b}(b)$.

1.2.4. *Homotopy of Paths in Z_0 over $\vec{v}$ Relative to their Endpoints.* Let p and q be two points in Z_0 and let γ_0 and γ_1 be two paths over $\vec{v}$ both starting in p and ending in q. We say that there is a *homotopy over $\vec{v}$ relative to their endpoints p and q* between γ_0 and γ_1, if there are continuous maps a, $b : I \to \mathbb{R}$ from the unit interval I to $\mathbb{R}$ with $a(s) < b(s)$ for all $s \in I$ and

$$H : \big\{(t,s)\big| s \in I,\, t \in [a(s), b(s)]\big\} \longrightarrow Z_0$$

such that $H(\cdot, 0) = \gamma_0$ and $H(\cdot, 1) = \gamma_1$ as well as that for any $s \in I$ the map $H(\cdot, s) : [a(s), b(s)] \to Z_0$ is a path over $\vec{v}$ with endpoints p and q. If the context is clear we will call two paths in Z_0 over $\vec{v}$ with the same endpoints *nearby homotopic relative to their endpoints*, if there is a homotopy over $\vec{v}$ relative to their endpoints between them. Note that the notion of nearby homotopy relative to the endpoints p and q as defined above yields an equivalence relation on the set of all paths in Z_0 over $\vec{v}$ with endpoints p and q.

1.2.5. *Homotopy of Paths in Z_0 over $\vec{v}$ based at $\vec{w}$.* We say that there is a *homotopy over $\vec{v}$ based at $\vec{w}$* between two paths γ_0 and γ_1 in Z_0 over $\vec{v}$ based at $\vec{w}$, if there are continuous maps a, $b : I \to \mathbb{R}$ with $a(s) < b(s)$ for all $s \in I$ and

$$H : \big\{(t,s)\big| s \in I,\, t \in [a(s), b(s)]\big\} \longrightarrow Z_0$$

such that $H(\cdot, 0) = \gamma_0$ and $H(\cdot, 1) = \gamma_1$ as well as that for any $s \in I$ the map $H(\cdot, s) : [a(s), b(s)] \to Z_0$ is a path over $\vec{v}$ based at $\vec{w}$. If there is no chance of confusion we will call two paths in Z_0 over $\vec{v}$ based at $\vec{w}$ *nearby homotopic* when there is a homotopy over $\vec{v}$ based at $\vec{w}$ between them. Note that the notion of *nearby homotopy* as defined above gives an equivalence relation on the set of all paths in Z_0 over $\vec{v}$ based at $\vec{w}$.

Moreover, remark that a homotopy over $\vec{v}$ based at $\vec{w}$ between two paths over $\vec{v}$ based at $\vec{w}$ is in particular a homotopy over $\vec{v}$ relative to the basepoint p_0 between these paths.

Proposition 1.2.

(i) *Every path $\gamma : [a, b] \to Z_0$ over $\vec{v}$ (based at $\vec{w}$) is nearby homotopic relative to its endpoints (resp. based at $\vec{w}$) to a path $\gamma_I : I \to Z_0$ over $\vec{v}$ (based at $\vec{w}$).*

(ii) *Assume, two paths over $\vec{v}$, say $\gamma_0, \gamma_1 : I \to Z_0$ with $\gamma_0(0) = \gamma_1(0)$ and $\gamma_0(1) = \gamma_1(1)$ (resp. both based at $\vec{w}$), are nearby homotopic relative to their endpoints (resp. based at $\vec{w}$). Then there is a nearby homotopy*

$$H : I \times I \longrightarrow Z_0$$

such that $H(\cdot, 0) = \gamma_0$ and $H(\cdot, 1) = \gamma_1$ as well as $H(\cdot, s)$ is a path over $\vec{v}$ with the endpoints $\gamma_0(0) = \gamma_1(0)$ and $\gamma_0(1) = \gamma_1(1)$ (resp. based at $\vec{w}$) for all $s \in I$.

For the proof of this proposition one has to reparametrize the paths and the homotopies in such a way that the speed stays unchanged just in small neighbourhoods around the double points. For a detailled proof we refer to [Kae97], Prop. 5.3.



Figure 2. The disks on the left (right) show a coordinate plot in a component D_k (D_l) with coordinate x (y). In these coordinates h is given by $h(x,y) = x \cdot y$. The centers of the disks represent the intersection point $p_{kl} = D_k \cap D_l$.

Remark: Observe that for example each γ for which there is a $\lambda \in [0, \frac{1}{2}[$ such that

$$\gamma(\frac{1}{2} - \tau) = \gamma(\frac{1}{2} + \tau) \qquad \forall\, \tau \in [0; \lambda]$$

is *homotopic over $\vec{v}$ based at $\vec{w}$* with the path that does not go all the way to $\gamma(\frac{1}{2})$ but returns already from $\gamma(\frac{1}{2} - \lambda) = \gamma(\frac{1}{2} + \lambda)$, that is the path

$$\begin{array}{rcc} \tilde{\gamma} : [0;\, 1 - 2\lambda] & \longrightarrow & Z_0 \\ t & \longmapsto & \left\{ \begin{smallmatrix} \gamma(t) & \text{for } t \leq \frac{1}{2} - \lambda \\ \gamma(t + 2\lambda) & \text{for } t \geq \frac{1}{2} - \lambda \end{smallmatrix} \right. \end{array}.$$

Let $a(s) \equiv 0$ and $b(s) = 1 - 2s\lambda$. Then the homotopy is given by

$$\begin{array}{rcc} H\, \left\{(t, s) \middle| s \in I,\, t \in [a(s), b(s)]\right\} & \longrightarrow & Z_0 \\ t & \longmapsto & \left\{ \begin{smallmatrix} \gamma(t) & \text{for } t \leq \frac{1}{2} - s\lambda \\ \gamma(t + 2s\lambda) & \text{for } t \geq \frac{1}{2} - s\lambda \end{smallmatrix} \right. \end{array}.$$

We denote the set of all 'homotopy over $\vec{v}$ based at $\vec{w}$'-classes of paths in Z_0 over $\vec{v}$ based at $\vec{w}$ by:

$$\pi_1(Z_{\vec{v}},\, \vec{w}).$$

Again, if the context is clear we will refer to elements of this set as to *nearby homotopy classes.* $\pi_1(Z_{\vec{v}},\, \vec{w})$ itself will be called *the nearby fundamental group (over $\vec{v}$ based at $\vec{w}$).* The term 'group' is justified by the following proposition.

Proposition 1.3. *$\pi_1(Z_{\vec{v}},\, \vec{w})$ with the composition of 'paths in Z_0 over $\vec{v}$ based at $\vec{w}$' is a group.*

Proof: Note first that the composition of two paths in Z_0 over $\vec{v}$ based at $\vec{w}$ is again a path in Z_0 over $\vec{v}$ based at $\vec{w}$. Furthermore the composition of such paths gives a well-defined operation on the set of homotopy classes, which is associative.

All paths $\gamma : I \to Z_0$ over $\vec{v}$ based at $\vec{w}$ with $\gamma(\frac{1}{2} - \tau) = \gamma(\frac{1}{2} + \tau)$ for all $\tau \in [0; \frac{1}{2}]$ are mutually nearby homotopic. Call their nearby homotopy class e and note that it is the neutral element with respect to the above defined operation on homotopy classes. Given an element $[\gamma] \in \pi_1(Z_{\vec{v}},\, \vec{w})$, the inverse is represented by the path $t \mapsto \gamma(1 - t)$ for $t \in I$. $\square$

1.3. **The Local System of Nearby Fundamental Groups.** The association of $\pi_1(Z_{\vec{v}},\, \vec{w})$ to a pair of non zero tangent vectors $\vec{v} \in (T_0\Delta)^* := T_0\Delta \setminus \{0\}$ and $\vec{w} \in (T_{p_0}D_0)^* := T_{p_0}D_0 \setminus \{0\}$ gives rise to a local system[3] of groups on $(T_0\Delta)^* \times (T_{p_0}D_0)^*$ as we want to point out in the sequel. We call this local system *the local system of nearby fundamental groups* and denote it by:

$$\{\pi_1(Z_{\vec{v}},\, \vec{w})\}_{(\vec{v}, \vec{w}) \in (T_0\Delta)^* \times (T_{p_0}D_0)^*}\,.$$

For a homotopy class modulo endpoints represented by a path

$$\vec{\eta} = \vec{\alpha} \times \vec{\beta} : I \to (T_0\Delta)^* \times (T_{p_0}D_0)^*,$$

i. e. a morphism of the fundamental groupoid of $(T_0\Delta)^* \times (T_{p_0}D_0)^*$, we define

$$\vec{\eta}_* : \pi_1(Z_{\vec{\alpha}(0)},\, \vec{\beta}(0)) \longrightarrow \pi_1(Z_{\vec{\alpha}(1)},\, \vec{\beta}(1))$$

---

[3] By a *local system of groups (vector spaces etc.) on a topological space* Y we mean a contravariant functor from the fundamental groupoid $\Pi(Y)$ of Y to the category of groups (vector spaces etc.). Note that we multiply paths in their natural order, that is $\alpha * \beta$ means 'traverse α, then β'.

in the following way. Let γ, a path over $\vec{\alpha}(0)$ based at $\vec{\beta}(0)$, represent an element in $\pi_1(Z_{\vec{\alpha}(0)}, \vec{\beta}(0))$. Moreover, let $H : I \times I \to Z_0$ be a homotopy with the property that $H(\cdot, 0) = \gamma$ and $H(\cdot, s)$ is a path over $\vec{\alpha}(s)$ based at $\vec{\beta}(s)$ for all $s \in I$. Note that such a homotopy always exists. Then define:

$$\vec{\eta}_* \left([\gamma]\right) := [H(\cdot, 1)].$$

In the following proposition we state that this $[H(\cdot, 1)]$ is well-defined and does not depend on the special choice of H.

Proposition 1.4. *Let $\vec{\eta}_i = \vec{\alpha}_i \times \vec{\beta}_i : I \to (T_0\Delta)^* \times (T_{p_0}D_0)^*$ for $i = 1, 2$ be two paths which are homotopic modulo endpoints. Let moreover $H_i : I \times I \to Z_0$ for $i = 1, 2$ be two homotopies such that $H_i(\cdot, s)$ is a path over $\vec{\alpha}_i(s)$ based at $\vec{\beta}_i(s)$ for any $s \in I$. If there is a homotopy over $\vec{\alpha}_1(0) = \vec{\alpha}_2(0)$ based at $\vec{\beta}_1(0) = \vec{\beta}_2(0)$ between $H_1(\cdot, 0)$ and $H_2(\cdot, 0)$, then there is a homotopy over $\vec{\alpha}_1(1) = \vec{\alpha}_2(1)$ based at $\vec{\beta}_1(1) = \vec{\beta}_2(1)$ between $H_1(\cdot, 1)$ and $H_2(\cdot, 1)$.*

The proof of this proposition can be found in the Appendix 1.4.

1.3.1. *The Local System of Fundamental Groups.* The locally trivial fibration $h : Z^* \to \Delta^*$ together with a section $\sigma : \Delta \to Z$ defines a local system on Δ^* by associating $\pi_1(Z_t, \sigma(t))$ to a point $t \in \Delta^*$. For a homotopy class modulo endpoints represented by a path in Δ^* one considers trivializations of the fibration over this path, which have σ as a constant section. Such a homotopy induces a map from the fundamental group over the starting point to the fundamental group over the endpoint.

However, here we would like to consider a local system on $\Delta^* \times D_0^*$. The idea is that such a section σ corresponds to a point p in Z_0 by taking $p = \sigma(0)$. We want to consider those sections σ, where $\sigma(0) \in D_0$. In order to be able to define morphisms we perform the following construction.

Let t resp. p be coordinates on Δ resp. D_0 such that Δ resp. D_0 are open disks with radius 1 in these coordinates. Consider the map

$$\begin{aligned} \pi : \Delta \times D_0^* &\longrightarrow \Delta \\ (t, p) &\longmapsto |p| \cdot t. \end{aligned}$$

Observe that all fibers of this map are punctured disks although π is not a locally trivial fibration. Nonetheless, if we define for any $\delta > 0$ the annulus $D_0^\delta := \left\{p \in D_0 \middle| |p| > \delta\right\}$, then the restriction of π to $\Delta \times D_0^\delta$ is a locally trivial fibration.

Let now $P : \Delta \times D_0^* \to Z$ be a continuous map such that the following diagram commutes:

$$\begin{tikzcd} \Delta \times D_0^* \arrow[rr, "P"] \arrow[dr, "\pi"'] & & Z \arrow[dl, "h"] \\ & \Delta & \end{tikzcd}$$

and such that $P\Big|_{\{0\} \times D_0^*} : D_0^* \to D_0^*$ is the identity.

For $\delta > 0$ we define $\Delta_\delta := \left\{t \in \Delta \middle| |t| < \delta\right\}$ and $\Delta_\delta^* := \Delta_\delta \setminus \{0\}$. Then for any $p \in D_0^\delta$ we have a section of $\pi\Big|_{\Delta \times D_0^\delta}$

$$\begin{aligned} \check{\sigma}_p : \Delta_\delta &\longrightarrow \Delta \times D_0^\delta \\ \tau &\longmapsto \left(\tfrac{\tau}{|p|}, p\right). \end{aligned}$$



FIGURE 3. Picture of the map π (restricted to the real coordinates in $\Delta \times D_0^*$ and Δ).

Composed with P we obtain a section of $h\big|_{h^{-1}(\Delta_\delta)}$, which we denote by $\sigma_p : \Delta_\delta \to Z$. Note $\sigma_p(0) = p$.

In the following, we will for any $0 < \delta < 1$ define a local system of groups on $\Delta_\delta^* \times D_0^\delta$. This system will be called *the local system of fundamental groups* and we denote it by:

$$\{\pi_1(Z_t,\, \sigma_p(t))\}_{(t,p)\in\Delta_\delta^*\times D_0^\delta}\,.$$

To a point (t,p) in $\Delta_\delta^* \times D_0^\delta$ we associate the group $\pi_1\left(Z_t, \sigma_p(t)\right)$. With a homotopy class modulo endpoints represented by a path

$$\eta = \alpha \times \beta : I \longrightarrow \Delta_\delta^* \times D_0^\delta$$

we associate a group isomorphism

$$\eta_* : \pi_1\left(Z_{\alpha(0)}, \sigma_{\beta(0)}(\alpha(0))\right) \longrightarrow \pi_1\left(Z_{\alpha(1)}, \sigma_{\beta(1)}(\alpha(1))\right).$$

in the following way.

Let $\gamma : I \to Z_{\alpha(0)}$ represent an element of $\pi_1\left(Z_{\alpha(0)}, \sigma_{\beta(0)}(\alpha(0))\right)$ and let $H : I \times I \to Z$ be a continuous map such that $H(\cdot, 0) = \gamma$ and $H(\cdot, s)$ is a path in $Z_{\alpha(s)}$ based at $\sigma_{\beta(s)}\left(\alpha(s)\right)$. Then define:

$$\eta_*\left([\gamma]\right) := [H(\cdot, 1)] \in \pi_1\left(Z_{\alpha(1)}, \sigma_{\beta(1)}(\alpha(1))\right).$$

Define

$$\overset{\circ}{\Delta}_\delta := \left\{\left(\rho, e^{2\pi i\theta}\right) \in \mathbb{R}^{\geq 0} \times S^1 \middle| \rho \cdot e^{2\pi i\theta} \in \Delta_\delta\right\}.$$

In the Appendix 1.4.2 we will define a local system on $\overset{\circ}{\Delta}_\delta \times D_0^\delta$ (and prove that it is well-defined) such that its restriction to $\Delta_\delta^* \times D_0^\delta$ by means of the obvious embedding $\Delta_\delta^* \times D_0^\delta \hookrightarrow \overset{\circ}{\Delta}_\delta \times D_0^\delta$ is the local system above.

Note that the above defined local system depends on the choice of the map P. However, it will be a consequence of Theorem 1.5 below that two different such maps P yield isomorphic local systems.

1.3.2. *Comparison of the two Local Systems.* Here we compare the local system of nearby fundamental groups with the local system of fundamental groups on the fibration. We state the following theorem. Note that $(T_0\Delta)^*$, $(T_{p_0}D_0)^*$, Δ_δ^* and D_0^δ are oriented by their complex structures.

Theorem 1.5. *Any pair of orientation preserving homotopy equivalences*

$$(T_0\Delta)^* \longrightarrow \Delta_\delta^* \quad \text{and} \quad (T_{p_0}D_0)^* \longrightarrow D_0^\delta$$

induces an isomorphism of the local system of nearby fundamental groups on the product $(T_0\Delta)^* \times (T_{p_0}D_0)^*$ *with the local system of fundamental groups on* $\Delta_\delta^* \times D_0^\delta$.

The proof of this theorem will be given in Appendix 1.4. The link between the two local systems will be provided by a local system which we call *the local system of the real blow-up.* Here we will make use of a construction of A'Campo [A'C75].

1.4. **Appendix.** This Appendix consists of two parts. In the first part we give the proof of Proposition 1.4. In the second we compare the two local systems of 1.3 by constructing a third one and show that this is isomorphic to both of them.

1.4.1. *Proof of 1.4:* We know that there is a homotopy, say $\mathcal{H} : I \times I \to Z_0$, between $H_1(\cdot, 1)$ and $H_2(\cdot, 1)$ by using first H_1 as homotopy between $H_1(\cdot, 1)$ and $H_1(\cdot, 0)$, then using the homotopy between $H_1(\cdot, 0)$ and $H_2(\cdot, 0)$ and finally using H_2 as homotopy between $H_2(\cdot, 0)$ and $H_2(\cdot, 1)$. The question is: Can such a homotopy be realized as homotopy over $\vec{\alpha}_1(1) = \vec{\alpha}_2(1)$ based at $\vec{\beta}_1(1) = \vec{\beta}_2(1)$? Define $\vec{\lambda} := (\vec{\alpha}_1)^{-1} \star \vec{\alpha}_i(0) \star \vec{\alpha}_2$ and $\vec{\mu} := (\vec{\beta}_1)^{-1} \star \vec{\beta}_i(0) \star \vec{\beta}_2$. Then we can formulate the question more precisely: Given closed paths $\vec{\lambda} : I \to (T_0\Delta)^*$ and $\vec{\mu} : I \to (T_{p_0}D_0)^*$, which are homotopic modulo endpoints to the respective constant paths and given a homotopy $\mathcal{H} : I \times I \to Z_0$ such that $\mathcal{H}(\cdot, s)$ is a path over $\vec{\lambda}(s)$ based at $\vec{\mu}(s)$, can we then find a homotopy $\tilde{\mathcal{H}} : I \times I \to Z_0$ between $\mathcal{H}(\cdot, 0)$ and $\mathcal{H}(\cdot, 1)$ such that $\mathcal{H}(\cdot, s)$ is a path over $\vec{\lambda}(0) = \vec{\lambda}(1)$ based at $\vec{\mu}(0) = \vec{\mu}(1)$ for all $s \in I$?

We give a positive answer to this question by doing the following calculation locally around each double point of Z_0.

Let t be a coordinate on Δ such that $\frac{\partial}{\partial t} = \vec{\lambda}(0) = \vec{\lambda}(1)$. Write $\vec{\lambda}(\tau) = l(\tau)\, \frac{\partial}{\partial t}$. Choose around each double point p_{kl} local coordinates (x, y) in Z such that the function h in the coordinates (x, y) and t looks like: $h(x, y) = x \cdot y$. For the double point p_0 we may assume that y is the coordinate on D_0 and that $\frac{\partial}{\partial y} = \vec{\mu}(0) = \vec{\mu}(1)$. Write $\vec{\mu}(\tau) = m(\tau)\, \frac{\partial}{\partial y}$.

Let $\vec{L} : I \times I \to (T_0\Delta)^*$ resp. $\vec{M} : I \times I \to (T_{p_0}D_0)^*$ be the homotopies modulo endpoints between $\vec{\lambda}$ resp. $\vec{\mu}$ and the constant path $\vec{\lambda}(0) = \vec{\lambda}(1)$ resp. $\vec{\mu}(0) = \vec{\mu}(1)$. Write $\vec{L}(\tau, s) = L(\tau, s)\, \frac{\partial}{\partial t}$ and $\vec{M}(\tau, s) = M(\tau, s)\, \frac{\partial}{\partial y}$.

By $\sigma : \mathbb{R}^{\geq 0} \to [0; 1]$ we denote a bump function which is $\equiv 1$ near 0 and $\equiv 0$ on $\mathbb{R}^{\geq r_0}$ for a sufficiently small r_0 and strictly decreasing inbetween.

For $(\tau, s) \in I \times I$ where $\mathcal{H}(\tau, s)$ lies in an x-coordinate, we describe $x \circ \mathcal{H}$ by polar coordinates: $x \circ \mathcal{H}(\tau, s) = r(\tau, s)\, e^{2\pi i \varphi(\tau, s)}$.

Then modify the homotopy $\mathcal{H}$ to a homotopy $\tilde{\tilde{\mathcal{H}}}$ by setting:

$$x \circ \tilde{\tilde{\mathcal{H}}}(\tau, s) := \frac{1}{l(\sigma(r(\tau, s)) \cdot s)}\, x \circ \mathcal{H}(\tau, s).$$



Figure 4. Illustration of a homotopy between a path $\vec{\lambda}$ in $T_0\Delta$ and the constant path $\vec{\lambda}(0) = \vec{\lambda}(1)$.



Figure 5. Illustration of a homotopy $\mathcal{H}$ (over the path $\vec{\lambda}$) at a double point (other than p_0) in the x-coordinate.



Figure 6. Illustration of the homotopy $\tilde{\tilde{\mathcal{H}}}$ at a double point (other than p_0) in the x-coordinate.



Figure 7. Illustration of the homotopy between $\mathcal{H}(\cdot, 1)$ and $\tilde{\tilde{\mathcal{H}}}(\cdot, 1)$ at a double point (other than p_0) in the x-coordinate.

Around the point p_0 we define $\tilde{\tilde{\mathcal{H}}}$ by:

$$x \circ \tilde{\tilde{\mathcal{H}}}(\tau, s) := \frac{m(\sigma(r(\tau, s)) \cdot s)}{l(\sigma(r(\tau, s)) \cdot s)}\, x \circ \mathcal{H}(\tau, s).$$

Note that $\tilde{\tilde{\mathcal{H}}}$ is a homotopy over $\vec{\lambda}(0) = \vec{\lambda}(1)$ based at $\vec{\mu}(0) = \vec{\mu}(1)$.

It remains to show that there is a homotopy over $\vec{\lambda}(0) = \vec{\lambda}(1)$ based at $\vec{\mu}(0) = \vec{\mu}(1)$ between $\tilde{\tilde{\mathcal{H}}}(\cdot, 1)$ and $\mathcal{H}(\cdot, 1)$. This homotopy is realized by the the following deformation of $\mathcal{H}(\cdot, 1)$ in the neighbourhood of each double point:

$$(\tau, s) \mapsto \frac{1}{L(\sigma(r(\tau, 1)), s)}\, x \circ \mathcal{H}(\tau, 1)$$

respectively around p_0:

$$(\tau, s) \mapsto \frac{M(\sigma(r(\tau, 1)), s)}{L(\sigma(r(\tau, 1)), s)}\, x \circ \mathcal{H}(\tau, 1). \qquad \square$$

1.4.2. *The Real Blow-up of the Fibration.* In this part of the Appendix we will compare the local system of nearby fundamental groups with the local system of fundamental groups. The link between these two will be given by a local system, which we call *the local system of the real blow-up* and which we are going to construct in the sequel.

We want to define an extension of Z^* to a topological space $\overset{\circ}{Z}$ and a map

$$\overset{\circ}{h}: \overset{\circ}{Z} \longrightarrow \overset{\circ}{\Delta},$$

in such a way that the diagram

$$\begin{tikzcd}
\overset{\circ}{Z} \arrow[d, "\overset{\circ}{h}"'] & Z^* \arrow[l, hook'] \arrow[d, "h^*"] \\
\overset{\circ}{\Delta} & \Delta^* \arrow[l, hook']
\end{tikzcd}$$

commutes and that $\overset{\circ}{h}: \overset{\circ}{Z} \to \overset{\circ}{\Delta}$ is a locally trivial topological fibration.

Choose a coordinate t on Δ and choose local coordinates $(x, y) : W_{kl} = U^k_{kl} \times U^l_{kl} \to \mathbb{C}^2$ around any double point $p_{kl} \in Z$ for $[k < l]$ such that the function h in these coordinates looks like

$$h(x, y) = x \cdot y \qquad \text{for all } |x| < 2,\ |y| < 2.$$

Consider the map

$$\begin{array}{rcl} \chi : \left(\mathbb{R}^{\geq 0} \times S^1\right) \times \left(\mathbb{R}^{\geq 0} \times S^1\right) & \longrightarrow & \left(\mathbb{R}^{\geq 0} \times S^1\right) \\ \left((R_1, e^{2\pi i \phi_1})\ ,\ (R_2, e^{2\pi i \phi_2})\right) & \longmapsto & \left(R_1 R_2, e^{2\pi i (\phi_1 + \phi_2)}\right). \end{array}$$

Then define for any pair k, l with $[k < l]$:

$$\overset{\circ}{U}{}^{k}_{kl} := \left\{ \left(R, e^{2\pi i \phi}\right) \in \mathbb{R}^{\geq 0} \times S^1 \middle| R e^{2\pi i \phi} \in x(U^k_{kl}) \right\}$$

and

$$\overset{\circ}{U}{}^{l}_{kl} := \left\{ \left(R, e^{2\pi i \phi}\right) \in \mathbb{R}^{\geq 0} \times S^1 \middle| R e^{2\pi i \phi} \in y(U^l_{kl}) \right\}.$$

There are the obvious maps $\overset{\circ}{U}{}^{k}_{kl} \to U^k_{kl}$ and $\overset{\circ}{U}{}^{l}_{kl} \to U^l_{kl}$.

We construct $\overset{\circ}{Z}$ as follows. Recall that there are obvious maps $\overset{\circ}{\Delta} \to \Delta$ and $\Delta^* \hookrightarrow \overset{\circ}{\Delta}$. Define first:

$$\breve{Z} := Z \times_{\Delta} \overset{\circ}{\Delta}.$$

In $\breve{Z}$ we have the subspaces

$$\begin{aligned} \breve{W}_{kl} &:= W_{kl} \times_{\Delta} \overset{\circ}{\Delta} = \left(U^k_{kl} \times U^l_{kl}\right) \times_{\Delta} \overset{\circ}{\Delta} \\ \breve{W}^*_{kl} &:= \left({U^*}^k_{kl} \times {U^*}^l_{kl}\right) \times_{\Delta^*} \Delta^*. \end{aligned}$$

Now we use $\chi_{\big|\overset{\circ}{U}^k_{kl} \times \overset{\circ}{U}^l_{kl}}$ and the identity map to define

$$\overset{\circ}{W}_{kl} := \left(\overset{\circ}{U}^k_{kl} \times \overset{\circ}{U}^l_{kl}\right) \times_{\overset{\circ}{\Delta}} \overset{\circ}{\Delta} .$$

Finally we construct $\overset{\circ}{Z}$ by replacing $\breve{W}_{kl}$ in $\breve{Z}$ by $\overset{\circ}{W}_{kl}$, glued along the common subspace W^*_{kl}. The projection to the second factor in the above mentioned fibered products yields the well-defined map we are looking after:

$$\overset{\circ}{h} : \overset{\circ}{Z} \longrightarrow \overset{\circ}{\Delta} .$$

Note that the maps $\overset{\circ}{U}^k_{kl} \to U^k_{kl}$ and $\overset{\circ}{U}^l_{kl} \to U^l_{kl}$ induce a map $\overset{\circ}{Z} \to Z$ over $\overset{\circ}{\Delta} \to \Delta$.
Remark: It seems that this construction was considered first by A'Campo in [A'C75]. There is a coordinate-free way to construct $\overset{\circ}{Z}$ by using the theory of log-structures (cf. e. g. [KN95], [Ill94], [Kat89], [Ste95]). We decided not to introduce the language of log-structures here since we use the space $\overset{\circ}{Z}$ just as a link between the two earlier defined local systems and because we need the coordinates anyway. The following theorem is proved in greater generality by Usui [Usu96].

Theorem 1.6. $\overset{\circ}{h} : \overset{\circ}{Z} \to \overset{\circ}{\Delta}$ *is a locally trivial topological fibration.*

This theorem and its proof can be found in in A'Campo [A'C75] p. 241. For a direct proof in this special situation we refer to [Kae97], Thm. 5.6, p. 81.

Having this theorem at hand we can define a third local system of groups – this time on $\overset{\circ}{\Delta}_\delta \times D^\delta_0$. The purpose of this local system here is that it serves as link between the local system of nearby fundamental groups and the local system of fundamental groups on the fibration. This third local system will be refered to as *the local system of the real blow-up*. We denote it by:

$$\left\{\pi_1\left(\overset{\circ}{Z}_{(\rho, e^{2\pi i\theta})}, \overset{\circ}{\sigma}_p\left(\rho, e^{2\pi i\theta}\right)\right)\right\}_{\left((\rho, e^{2\pi i\theta}), p\right) \in \overset{\circ}{\Delta}_\delta \times D^\delta_0} .$$

It is an extension of the local system of fundamental groups on the fibration on $\Delta^*_\delta \times D^\delta_0$ to $\overset{\circ}{\Delta}_\delta \times D^\delta_0$ and it is defined as follows:

Observe that for $p \in D^\delta_0$ the section $\sigma_p : \Delta_\delta \to Z$ can be pulled back in a unique way to a section $\overset{\circ}{\sigma}_p : \overset{\circ}{\Delta}_\delta \to \overset{\circ}{Z}$. To any $\left((\rho, e^{2\pi i\theta}), p\right) \in \overset{\circ}{\Delta}_\delta \times D^\delta_0$ we associate the group

$$\pi_1\left(\overset{\circ}{Z}_{(\rho, e^{2\pi i\theta})}, \overset{\circ}{\sigma}_p\left(\rho, e^{2\pi i\theta}\right)\right)$$

and to a homotopy class modulo endpoints represented by a path

$$\eta = \alpha \times \beta : I \longrightarrow \overset{\circ}{\Delta}_\delta \times D^\delta_0$$

we associate similarly as before a group isomorphism

$$\eta_* : \pi_1\left(\overset{\circ}{Z}_{\alpha(0)}, \overset{\circ}{\sigma}_{\beta(0)}\left(\alpha(0)\right)\right) \longrightarrow \pi_1\left(\overset{\circ}{Z}_{\alpha(1)}, \overset{\circ}{\sigma}_{\beta(1)}\left(\alpha(1)\right)\right) .$$

Let $\gamma : I \longrightarrow \overset{\circ}{Z}$ be an element of $\pi_1\left(\overset{\circ}{Z}_{\alpha(0)}, \overset{\circ}{\sigma}_{\beta(0)}(\alpha(0))\right)$ and let $H : I \times I \longrightarrow \overset{\circ}{H}$ be a continuous map such that $H(\cdot, 0) = \gamma$ and for any $s \in I$ is $H(\cdot, s)$ a path in $\overset{\circ}{Z}_{\alpha(s)}$ based at $\overset{\circ}{\sigma}_{\beta(s)}(\alpha(s))$. Then define:

$$\eta_*([\gamma]) := [H(\cdot, 1)] \in \pi_1\left(\overset{\circ}{Z}_{\alpha(1)}, \overset{\circ}{\sigma}_{\beta(1)}(\alpha(1))\right).$$

The following proposition justifies this definition and moreover the definition of the local system of fundamental groups of the fibration.

Proposition 1.7. *Let $\eta_i = \alpha_i \times \beta_i : I \rightarrow \overset{\circ}{\Delta}_\delta \times D_0^\delta$ for $i = 1, 2$ be two paths, which are homotopic modulo endpoints. Let moreover $H_i : I \times I \rightarrow \overset{\circ}{Z}$ for $i = 1, 2$ be two homotopies such that $H_i(\cdot, s)$ is a path in $\overset{\circ}{Z}_{\alpha_i(s)}$ based at $\overset{\circ}{\sigma}_{\beta_i(s)}(\alpha_i(s))$ for any $s \in I$. If $H_1(\cdot, 0)$ and $H_2(\cdot, 0)$ represent the same homotopy class in*

$$\pi_1\left(\overset{\circ}{Z}_{\alpha_i(0)}, \overset{\circ}{\sigma}_{\beta_i(0)}(\alpha_i(0))\right)$$

with $i = 1$ or 2, then $H_1(\cdot, 1)$ and $H_2(\cdot, 1)$ represent the same homotopy class in

$$\pi_1\left(\overset{\circ}{Z}_{\alpha_i(1)}, \overset{\circ}{\sigma}_{\beta_i(1)}(\alpha_i(1))\right)$$

with $i = 1$ or 2.

Proof: Similar as in the proof of Proposition 1.4 the assertion may be reduced to: If $\lambda : I \rightarrow \overset{\circ}{\Delta}_\delta$ and $\mu : I \rightarrow D_0^\delta$ are closed paths, which are homotopic modulo endpoint to the respective constant paths and given a homotopy $\mathcal{H} : I \times I \rightarrow \overset{\circ}{Z}$ such that $\mathcal{H}(\cdot, s)$ is a path in $\overset{\circ}{Z}_{\lambda(s)}$ based at $\overset{\circ}{\sigma}_{\mu(s)}(\lambda(s))$ for any $s \in I$, then there is a $\tilde{\mathcal{H}} : I \times I \rightarrow \overset{\circ}{Z}$, a homotopy between $\mathcal{H}(\cdot, 0)$ and $\mathcal{H}(\cdot, 1)$ such that $\tilde{\mathcal{H}}(\cdot, s)$ is a path in $\overset{\circ}{Z}_{\lambda(0)} = \overset{\circ}{Z}_{\lambda(1)}$ based at $\overset{\circ}{\sigma}_{\mu(0)}(\lambda(0)) = \overset{\circ}{\sigma}_{\mu(1)}(\lambda(1))$ for any $s \in I$.

Observe first that there is a homotopy $\tilde{\tilde{\mathcal{H}}} : I \times I \rightarrow \overset{\circ}{Z}_{\lambda(0)} = \overset{\circ}{Z}_{\lambda(1)}$ between $\mathcal{H}(\cdot, 0)$ and $\mathcal{H}(\cdot, 1)$ lying completely in the one fiber over $\lambda(0) = \lambda(1)$ but not necessarily leaving the base point $\overset{\circ}{\sigma}_{\mu(0)}(\lambda(0)) = \overset{\circ}{\sigma}_{\mu(1)}(\lambda(1))$ fixed. This is true for the following reason. Let $L : I \times I \rightarrow \overset{\circ}{\Delta}_\delta$ be the homotopy between λ and the constant path $\lambda(0) = \lambda(1)$. Then the pull-back bundle $L^* \overset{\circ}{Z}$ is globally trivial, i. e. there is a trivialization

$$\begin{tikzcd}
L^* \overset{\circ}{Z} \arrow[rr, "\Phi"] \arrow[dr, "L^* h"] & & (I \times I) \times \overset{\circ}{Z}_{\lambda(0)} \arrow[dl, "pr_2"] \\
& I \times I &
\end{tikzcd}.$$

The homotopy $\mathcal{H}$ can be considered as a map to $L^* \overset{\circ}{Z}$ (since $\mathcal{H}(\cdot, s)$ is a path in $\overset{\circ}{Z}_{\lambda(s)}$ for $s \in I$) and therefore to $(I \times I) \times \overset{\circ}{Z}_{\lambda(0)}$. The projection to the second factor is a homotopy in $\overset{\circ}{Z}_{\lambda(0)}$ between $\mathcal{H}(\cdot, 0)$ and $\mathcal{H}(\cdot, 1)$. If the path, which is formed by the trace of the base point is zero-homotopic, then we may find a homotopy between $\mathcal{H}(\cdot, 0)$ and $\mathcal{H}(\cdot, 1)$ leaving the base point $\overset{\circ}{\sigma}_{\mu(0)}(\lambda(0)) = \overset{\circ}{\sigma}_{\mu(1)}(\lambda(1))$ fixed.

Note that a path in $(I \times I) \times \overset{\circ}{Z}_{\lambda(0)}$ is zero homotopic if and only if its projection to $\overset{\circ}{Z}_{\lambda(0)}$ is. The above mentioned trace of the base point is the projection to $\overset{\circ}{Z}_{\lambda(0)}$

of the path

$$\phi\left(\mathcal{H}(0,\cdot)\right) = \phi\left(\mathcal{H}(1,\cdot)\right).$$

But this path is zero-homotopic if and only if $\mathcal{H}(0,\cdot) = \mathcal{H}(1,\cdot)$ is. Note that $\mathcal{H}(0,s) = \mathcal{H}(1,s) = \overset{\circ}{\sigma}_{\mu(s)}(\lambda(s))$. Therefore, a homotopy to the constant path $\overset{\circ}{\sigma}_{\mu(0)}(\lambda(0)) = \overset{\circ}{\sigma}_{\mu(1)}(\lambda(1))$ is given by: $(s,\tau) \longmapsto \overset{\circ}{\sigma}_{M(s,\tau)}(L(s,\tau))$. ☐

Let us conclude this section by proving Theorem 1.5, which asserts that any pair of orientation preserving homotopy equivalences

$$(T_0\Delta)^* \longrightarrow \Delta_\delta^* \quad \text{and} \quad (T_{p_0}D_0)^* \longrightarrow D_0^\delta$$

induces an isomorphism of the local system of nearby fundamental groups on the product $(T_0\Delta)^* \times (T_{p_0}D_0)^*$ with the local system of fundamental groups on $\Delta_\delta^* \times D_0^\delta$.

Proof of 1.5: Observe that $\Delta_\delta^* \subset \overset{\circ}{\Delta}_\delta$ and call $S := \overset{\circ}{\Delta}_\delta \setminus \Delta_\delta^*$. By definition of both local systems, the inclusion $\Delta_\delta^* \times D_0^\delta \subset \overset{\circ}{\Delta}_\delta \times D_0^\delta$ induces an isomorphism of the local system of fundamental groups on the fibration with the local system of the real blow up. The latter local system is also equivalent to its restriction to $S \times D_0^\delta$ by the inclusion $S \times D_0^\delta \subset \overset{\circ}{\Delta}_\delta \times D_0^\delta$. We call this restriction to $S \times D_0^\delta$ *the local system on the soul of the real blow-up*.

Hence, the assertion of the theorem is equivalent with: Any pair of orientation preserving homotopy equivalences

$$a : (T_0\Delta)^* \longrightarrow S \quad \text{and} \quad b : (T_{p_0}D_0)^* \longrightarrow D_0^\delta$$

induces an isomorphism of the local system of nearby fundamental groups to the local system on the soul of the real blow-up.

We have to show that there is an isomorphism between the group associated to one pair of tangent vectors $(\vec{v}_0, \vec{w}_0) \in (T_0\Delta)^* \times (T_{p_0}D_0)^*$ and the group associated to $(a(\vec{v}_0), b(\vec{w}_0)) \in S \times D_0^\delta$, which is equivariant with respect to the action of $\pi_1\left((T_0\Delta)^* \times (T_{p_0}D_0)^*, (\vec{v}_0,\vec{w}_0)\right)$ and the action of $\pi_1\left(S \times D_0^\delta, (a(\vec{v}_0), b(\vec{w}_0))\right)$.

Let us first construct the isomorphism and show that it is equivariant. Associated to each double point p_{kl} for $[k<l]$ we have from the definition of $\overset{\circ}{Z}$ coordinates $\left(R_1, e^{2\pi i\phi_1}\right)$, $\left(R_2, e^{2\pi i\phi_2}\right)$ in $\overset{\circ}{Z}$ and coordinates $\left(\rho, e^{2\pi i\theta}\right)$ on $\overset{\circ}{\Delta}$ such that in these coordinates

$$\overset{\circ}{h}\left(\left(R_1, e^{2\pi i\phi_1}\right), \left(R_2, e^{2\pi i\phi_2}\right)\right) = \left(R_1R_2, e^{2\pi i(\phi_1+\phi_2)}\right).$$

Here we may assume that $a(\vec{v}_0) = (0,1) \in S$ and $b(\vec{w}_0) = (1,0) \in \overset{\circ}{\bar{D}}_0$.

We will define an isomorphism

$$\mathcal{J} : \pi_1\left(Z_{\vec{v}_0}, \vec{w}_0\right) \longrightarrow \pi_1\left(\overset{\circ}{Z}_{(0,1)}, \overset{\circ}{\sigma}_{(1,0)}(0,1)\right).$$

Let $[\gamma]$ be a nearby homotopy class in $\pi_1\left(Z_{\vec{v}_0}, \vec{w}_0\right)$ represented by the path γ over $\vec{v}_0$ based at $\vec{w}_0$. Note that any piecewise smooth path in $\mathbb{C}$ has a unique lifting with respect to the map

$$\begin{array}{rccc} \overset{\circ}{\mathbb{C}} := & \mathbb{R}^{\geq 0} \times S^1 & \to & \mathbb{C} \\ & \left(r, e^{2\pi i\varphi}\right) & \mapsto & r \cdot e^{2\pi i\varphi}. \end{array}$$

Therefore, there is a unique lifting of $\gamma \in Z_0$ to a path $\overset{\bullet}{\gamma}$ in $\overset{\circ}{Z}$. The condition that γ lies over $\vec{v}_0$ implies that $\overset{\bullet}{\gamma}$ is even in $\overset{\circ}{Z}_{(0,1)}$. and that it is connected. Define then $\overset{\circ}{\gamma}:= \sigma\star\overset{\bullet}{\gamma}\star\sigma^{-1}$, where σ is the straight path in $\overset{\circ}{D}_0$ defined by $t \mapsto (t, e^{2\pi i0}) = (1-t, 1)$, then we obtain a path in $\overset{\circ}{Z}_{(0,1)}$, which is based at $\overset{\circ}{\sigma}_{(1,0)}\ (0,1) = (1,0) \in \overset{\circ}{\bar{D}}_0$.

The map $\gamma\ \mapsto \overset{\circ}{\gamma}$ induces a well-defined map from the group $\pi_1\left(Z_{\vec{v}_0}, \vec{w}_0\right)$ to $\pi_1(\overset{\circ}{Z}_{(0,1)}, \overset{\circ}{\sigma}_{(1,0)}\ (0,1))$, which is certainly a group homomorphism. Define $\mathcal{J}$ by $\mathcal{J}(\gamma) := \overset{\circ}{\gamma}$. It remains to show that $\mathcal{J}$ is an *iso*morphism.

$\mathcal{J}$ **is surjective:** Call the sets $\left\{(0, e^{2\pi i\varphi}),\ (0, e^{-2\pi i\varphi}) \middle| \varphi \in [0;1[\right\} \subset \overset{\circ}{P}_{kl} \cap \overset{\circ}{Z}_{(0,1)}$ for $[k<l]$ *vanishing cycles*. We can give $\overset{\circ}{Z}_{(0,1)}$ a differentiable structure in such a way that it coincides with the already given differentiable structure off the vanishing cycles coming from $Z_0 \setminus \bigcup_{[k<l]}\{p_{kl}\}$. It is well-known that any element in the fundamental group of a smooth manifold (with boundary) can be represented by a smooth path (cf. e. g. [tD91], II, 1.10). Moreover observe that by a local construction any such smooth path is homotopic to a smooth path, which intersects the vanishing cycles transversally in a finite number of parameter values, the first and the last of which are the only intersections with the vanishing cycle corresponding to p_0. Moreover, we may assume that $\overset{\odot}{\gamma}$ can be written as $\overset{\odot}{\gamma} = \sigma \star \overset{\bullet}{\gamma} \star \sigma^{-1}$ for some closed path $\overset{\bullet}{\gamma}$, which lies, apart from the base point, entirely in D^+.

A path $\overset{\odot}{\gamma}$ in $\overset{\circ}{Z}_{(0,1)}$ with these properties, composed with the map $\overset{\circ}{Z}_{(0,1)} \rightarrow Z_0$ (collapsing the vanishing cycles) yields a path γ in $D^+ \subset Z_0$. By reparametrizing the path $\overset{\odot}{\gamma}$ appropriately we may assume that it lies over $\vec{v}_0$ and is based at $\vec{w}_0$. And hence we find: $\overset{\circ}{\gamma} = \overset{\odot}{\gamma}$.

$\mathcal{J}$ **is injective:** Here we have to convince ourselves from the following fact. If we are given two homotopic paths $\overset{\circ}{\gamma}_1$ and $\overset{\circ}{\gamma}_2$ in $\overset{\circ}{Z}_{(0,1)}$ based at $(1,0) \in \overset{\circ}{\bar{D}}_0$, which are images under $\mathcal{J}$ of two paths γ_1 and γ_2 over $\vec{v}_0$ based at $\vec{w}_0$, then these latter paths are homotopic over $\vec{v}_0$ based at $\vec{w}_0$.

Let $H : I \times I \rightarrow \overset{\circ}{Z}_{(0,1)}$ be a homotopy between the paths $\overset{\circ}{\gamma}_1$ and $\overset{\circ}{\gamma}_2$ in $\overset{\circ}{Z}_{(0,1)}$ based at $(1,0) \in \overset{\circ}{\bar{D}}_0$. Using again the differentiable structure on $\overset{\circ}{Z}_{(0,1)}$ like above we may assume that H is smooth (cf. e. g. [tD91], II, 1.10) and that for each $s \in I$ the path $H(\cdot, s)$ intersects the the vanishing cycles transversally in a finite number of parameter values, the first and the last of which are the only intersections with the vanishing cycle corresponding to p_0. Moreover, we may assume that $H(\cdot, s)$ can be written as $H(\cdot, s) = \sigma \star \eta_s \star \sigma^{-1}$ for some closed path η_s, which lies, apart from the base point, entirely in D^+.

Composed with $\overset{\circ}{Z}_{(0,1)} \rightarrow Z_0$ we obtain a homotopy between γ_1 and γ_2 over $\vec{v}_0$ based at $\vec{w}_0$. Finally, we study the monodromy actions. Let

$$\vec{\eta} = \vec{\alpha} \times \vec{\beta} : I \longrightarrow (T_0\Delta)^* \times (T_{p_0}D_0)^*$$

be a path representing a morphism in the fundamental groupoid of $(T_0\Delta)^* \times (T_{p_0}D_0)^*$. Then define $\eta := (a \circ \vec{\alpha}) \times \left(b \circ \vec{\beta}\right) : I \longrightarrow S \times D_0^\delta$. And let γ be a path over $\vec{v}_0$ based at $\vec{w}_0$ representing an element in $\pi_1\left(Z_{\vec{v}_0}, \vec{w}_0\right)$. It is to show: $\mathcal{J} \circ \vec{\eta}_* \left([\gamma]\right) = \eta_* \circ \mathcal{J}\left([\gamma]\right)$.

Let $H : I \times I \to Z_0$ be a homotopy with the property that $H(\cdot, 0) = \gamma$ and $H(\cdot, s)$ is a path over $\vec{\alpha}(s)$ based at $\vec{\beta}(s)$ for every $s \in I$. Like the construction of $\overset{\circ}{\gamma}$ from γ on page 19 we construct a homotopy $\overset{\circ}{H}\colon I \times I \to \overset{\circ}{Z}$ from H such that $\overset{\circ}{H}(\cdot, 0) = \overset{\circ}{\gamma}$ and such that $\overset{\circ}{H}(\cdot, s)$ is a path $\overset{\circ}{Z}_{a \circ \vec{\alpha}(s)}$ based at $b \circ \vec{\beta}(s)$. This can be achieved by the definition: $\overset{\circ}{H}(\tau, s) := (H(\cdot, s))^{\circ}(\tau)$. Now compute:

$$\mathcal{J} \circ \vec{\eta}_* ([\gamma]) = \mathcal{J}([H(\cdot, 1)]) = \left[\overset{\circ}{H}(\cdot, 1)\right] \text{ and } \eta_* \circ \mathcal{J}([\gamma]) = \eta_* \left([\overset{\circ}{\gamma}]\right) = \left[\overset{\circ}{H}(\cdot, 1)\right].$$

□

2. Line Integrals on the Nearby Fundamental Group

In Section 1 we defined the group

$$\pi_1(Z_{\vec{v}}, \vec{w}).$$

Consider the group ring $\mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w})$ and let $J_{\vec{v}\vec{w}}$ or simply $\vec{J}$ be the augmentation ideal. By definition, this is the set of elements in $\mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w})$, whose sum of coefficients vanishes. Our aim is to put a $\mathbb{Z}$-MHS on

$$\mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w}) \Big/ \vec{J}^{s+1}$$

for any $s \geq 1$. This will be done by putting a MHS on $(\vec{J}/\vec{J}^{s+1})$. Note that multiplication induces an isomorphism $\vec{J}^s/\vec{J}^{s+1} \xrightarrow{\cong} (\vec{J}/\vec{J}^{\,2})^{\otimes s}$ (recall that: $\pi_1(Z_{\vec{v}}, \vec{w}) \cong \pi_1(Z_t, p)$ for some $t \in \Delta^*$ and $p \in Z_t$). The short exact sequence

$$0 \to \vec{J}^s/\vec{J}^{s+1} \to \vec{J}/\vec{J}^{s+1} \to \vec{J}/\vec{J}^s \to 0 \tag{2.1}$$

will be a short exact sequence of MHSs.

The main ingredient for the definition of these MHSs is a differential graded algebra (dga) $A^\bullet$, which is made up from differential forms on the central fiber but whose cohomology is isomorphic to the cohomology of a regular fiber. It was suggested to us by Hain[4] to use a dga like $A^\bullet$ and he proposed the problem of finding a way to integrate elements of this dga somehow on the central fiber. We will see that the notion of *a path over $\vec{v}$ (based at $\vec{w}$)* will be the right notion of path for this purpose. We show that closed elements in A^1 can be integrated along paths in Z_0 over $\vec{v}$ in a natural way. This yields an integral structure on $H^1(A^\bullet)$, which is by this integration dual to $\vec{J}/\vec{J}^{\,2}$. On $A^\bullet$ we have filtrations $W_\bullet$ and $F^\bullet$ such that the induced filtrations together with the integral structure define a $\mathbb{Z}$-MHS on $H^1(A^\bullet)$. It will turn out that the integral lattice of this $\mathbb{Z}$-MHS depends on $\vec{v}$ (and not on $\vec{w}$). The dependence upon $\vec{v}$ is such that $\{(J_{\vec{v},\vec{w}}/J^2_{\vec{v},\vec{w}})^*\}_{\vec{v} \in (T_0\Delta)^*}$ is a nilpotent orbit of MHSs.

Since $A^\bullet$ has an algebra-structure, we can also define *iterated line integrals* along paths over $\vec{v}$. We shall do this in section 3. This will give a way to describe $(\vec{J}/\vec{J}^{s+1})^*$ in terms of the dga $A^\bullet$ similar to Chen's π_1-De Rham theorem (cf. [Che75b], [Che77b] or [Hai87c]). Finally it will lead to the definition of a MHS on $(\vec{J}/\vec{J}^{s+1})^*$ such that the short exact sequence (2.1) becomes a short exact sequence of MHSs.

---

[4] The dga $A^\bullet$ is in principle a modified version of the complex $P^\bullet_{\mathbb{C}}(X^*)[\theta]$ of [Hai87b] for this special situation with non compact fibers.

2.1. **A Differential Graded Algebra of Differential Forms on the Central Fiber.** Let $(\Lambda^\bullet, d) := (\bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})[\log t], d)$ be the dga defined in the following way. Let $\bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})$ be the free graded-commutative $\mathbb{C}$-algebra with unit, generated by the symbols $\frac{dx}{x}, \frac{dy}{y}$ in degree 1.

Let then $\Lambda^\bullet := \bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})[\log t]$ be the algebra $\bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})$, where the symbol[5] $\log t$ or u has been added in degree 0. The differential d on $\Lambda^\bullet$ is defined by the Leibniz-rule, $d\big(\frac{dx}{x}\big) = d\big(\frac{dy}{y}\big) = 0$ and

$$d\ (\log t) := \frac{dx}{x} + \frac{dy}{y}.$$

Note that $(\Lambda^\bullet, d)$ computes the cohomology of the fibers of the map $\Delta^* \times \Delta^* \to \Delta^*$ given by $(x, y) \mapsto xy$.

For each component D_i of D^+ let

$$E^\bullet(D_i \log P_i) := \Gamma\left(\Omega^\bullet(D_i \log P_i) \otimes_{\Omega^0(D_i)} \mathcal{E}^{0,\bullet}(D_i)\right)$$

be the logarithmic de Rham complex. Here $\mathcal{E}^{0,\bullet}(D_i)$ denotes the sheaf of C^∞-forms of type $(0, \bullet)$. Note that $(E^\bullet(D_i \log P_i), d)$ computes the cohomology of $D_i \setminus P_i$.

Finally let $\bigwedge^\bullet(\frac{dp}{p})$ be the differential graded-commutative $\mathbb{C}$-algebra with unit generated by the symbol $\frac{dp}{p}$ in degree 1, where the differential d is defined to be always the zero-map. $\left(\bigwedge^\bullet(\frac{dp}{p}), d\right)$ computes the cohomology of $\mathbb{C}^*$. Now we consider r copies $\left(\bigwedge_i^\bullet(\frac{dp}{p}), d\right)$ $i = 0, \dots, r-1$ of $\left(\bigwedge^\bullet(\frac{dp}{p}), d\right)$ and think of an element g_i in $\bigwedge_i^0(\frac{dp}{p}) = \mathbb{C}$ as a function on the point p_i, i. e. $g_i(p_i) := g_i$. We refer to the complex number ρ of an element $\rho\frac{dp}{p} \in \bigwedge_i^1(\frac{dp}{p})$ as *the residue in* p_i of this element. Write $\mathrm{Res}_{p_i}(\rho\frac{dp}{p}) = \rho$.

The dga $A^\bullet$, which we are going to define will be a sub-dga of

$$B^\bullet := \bigoplus_{0 \le i \le r-1} \bigwedge\nolimits_i^\bullet(\frac{dp}{p}) \oplus \bigoplus_{i > r-1} E^\bullet(D_i \log P_i) \oplus \bigoplus_{[k<l]} E^\bullet(\Delta^1) \otimes_{\mathbb{C}} \Lambda^\bullet,$$

where Δ^1 denotes the one-dimensional standard-simplex (the unit interval) and $E^\bullet(\Delta^1) = \mathbb{C}[\xi] \oplus \mathbb{C}[\xi]d\xi$ the dga of real polynomial differential forms on Δ^1.

The elements of $A^\bullet$ in $B^\bullet$ will be called *the compatible elements* in $B^\bullet$. Hence, we will define $A^\bullet$ now by defining, when an element of $B^\bullet$ is compatible. Observe that $B^n = 0$ for $n \ge 4$.

We shall use P_{kl}, K_{kl}, L_{kl}, H_{kl}, R_{kl}, S_{kl}, T_{kl} and U_{kl} to denote elements in $\mathbb{C}[\xi, u]$. If the context is clear we omit the indices k, l. Moreover it will be useful to expand these elements of $\mathbb{C}[\xi, u]$ as polynomials in u with coefficients in $\mathbb{C}[\xi]$. That means for instance

$$P_{kl} = P_{kl}(\xi, u) = P = P_0 + P_1 u + P_2 u^2 + \cdots + P_m u^m \text{ and } P' = \frac{\partial P}{\partial \xi}.$$

$\mathbf{A^0}$: An element $f = \sum_{i \ge 0} g_i + \sum_{[k<l]} P_{kl} \in B^0$, where $g_i \in \bigwedge_i^0(\frac{dp}{p})$ for $i \le r-1$ and $g_i \in E^0(D_i)$ for $i > r-1$ is called *compatible* iff for $[k < l]$ holds:

$$P_{kl}(0, u) = g_k(p_{kl}) \text{ and } P_{kl}(1, u) = g_l(p_{kl}).$$

---

[5] We will use u for convenience or if confusion with the function 'log' might occur.

$\mathbf{A^1}$: We call an element in B^1,

$$\varphi = \sum_{i\geq 0} \omega_i + \sum_{[k<l]} \left(K_{kl} \frac{dx}{x} + L_{kl} \frac{dy}{y} + H_{kl} d\xi \right),$$

where $\omega_i \in \bigwedge^1(\frac{dp}{p})$ for $i \leq r-1$ and $\omega_i \in E^1(D_i \log P_i)$ for $i > r-1$, a *compatible* element iff for $[k<l]$ holds:

$$\begin{array}{rcl} K_{kl}(0,u) = & \mathrm{Res}_{p_{kl}}\omega_k & , \quad L_{kl}(0,u) = 0, \\ K_{kl}(1,u) = & 0 & , \quad L_{kl}(1,u) = \mathrm{Res}_{p_{kl}}\omega_l. \end{array}$$

$\mathbf{A^2}$: Let us call an element

$$\phi = \sum_{i>r-1} \Omega_i + \sum_{[k<l]} \left(R_{kl}\, d\xi \wedge \frac{dx}{x} + S_{kl}\, d\xi \wedge \frac{dy}{y} + T_{kl}\, \frac{dx}{x} \wedge \frac{dy}{y} \right) \in B^2,$$

with $\Omega_i \in E^2(D_i \log P_i)$ for $i > r-1$, *compatible* iff for $[k<l]$ holds:

$$T_{kl}(0,u) = T_{kl}(1,u) = 0.$$

$\mathbf{A^3}$: We define $A^3 := B^3$. That is, all elements of B^3 are *compatible*.

Observe that holds $dA^i \subset A^{i+1}$ for $i \geq 0$ and that $A^\bullet$ is a dga.

Definition 2.1. On $A^\bullet$ we define an *augmentation map* a by:

$$\begin{array}{rccc} a: & A^0 & \to & \mathbb{C} \\ & \sum_{i\geq 0} g_i + \sum_{[k<l]} P_{kl} & \mapsto & g_0. \end{array}$$

An alternative definition of $A^\bullet$ can be given as follows. Consider the surjective map of complexes

$$\Phi : B^\bullet \longrightarrow \bigoplus_{[k<l]} \Lambda^\bullet \oplus \Lambda^\bullet,$$

which sends $\sum_{i\geq 0} g_i + \sum_{[k<l]} P_{kl} \in B^0$ to

$$\sum_{[k<l]} \{(P_{kl}(0,u) - g_k(p_{kl})) \oplus (P_{kl}(1,u) - g_l(p_{kl}))\}$$

and $\sum_{i\geq 0} \omega_i + \sum_{[k<l]} \left(K_{kl} \frac{dx}{x} + L_{kl} \frac{dy}{y} + H_{kl} d\xi \right) \in B^1$ to

$$\begin{aligned} \sum_{[k<l]} \Bigg\{ & \left[(K_{kl}(0,u) - \mathrm{Res}_{p_{kl}}\omega_k) \frac{dx}{x} + L_{kl}(0,u) \frac{dy}{y} \right] \\ \oplus \quad & \left[K_{kl}(1,u) \frac{dx}{x} + (L_{kl}(1,u) - \mathrm{Res}_{p_{kl}}\omega_l) \frac{dy}{y} \right] \Bigg\}. \end{aligned}$$

Moreover, Φ maps

$$\sum_{i>r-1} \Omega_i + \sum_{[k<l]} R_{kl}\, d\xi \wedge \frac{dx}{x} + S_{kl}\, d\xi \wedge \frac{dy}{y} + T_{kl}\, \frac{dx}{x} \wedge \frac{dy}{y} \in B^2$$

to $\sum_{[k<l]} T_{kl}(0,u) \frac{dx}{x} \wedge \frac{dy}{y} \oplus T_{kl}(1,u) \frac{dx}{x} \wedge \frac{dy}{y}$. It is easy to check that Φ is indeed a surjective map of complexes. We find:

$$A^\bullet = \ker \Phi.$$

Define $C^\bullet := \sum_{[k<l]} \Lambda^\bullet \oplus \Lambda^\bullet$ and note that we have the short exact sequence

$$0 \longrightarrow A^\bullet \longrightarrow B^\bullet \longrightarrow C^\bullet \longrightarrow 0,$$

which yields a long exact cohomology sequence (as $H^i(B^\bullet) = 0$ for $i \geq 2$)

$$(2.2)\quad 0 \to H^0(A^\bullet) \to H^0(B^\bullet) \to H^0(C^\bullet) \\ \to H^1(A^\bullet) \to H^1(B^\bullet) \to H^1(C^\bullet) \to H^2(A^\bullet) \to 0.$$

The complex $(A^\bullet, d)$ computes the cohomology of the regular fiber as the following theorem asserts.

Theorem 2.2. *$H^\bullet(A^\bullet) \cong H^\bullet(Z_t; \mathbb{C})$ for any $t \in \Delta^*$.*

The proof of this theorem can be found in the Appendix 2.5 on page 33.

2.2. **Integration in the Nearby Fiber along Paths over a Tangent Vector.**

Here we are going to define integrals of a closed element

$$\varphi = \sum_{i \geq 0} \omega_i + \sum_{[k<l]} \left(K_{kl} \frac{dx}{x} + L_{kl} \frac{dy}{y} + H_{kl} d\xi \right) \in A^1$$

along paths over $\vec{v}$. We define those integrals in four steps by defining them first in special cases.

(a): Let $\gamma : [a; b] \to Z_0$ be a path over $\vec{v}$, which meets the set of double points only once with parameter value $\tau_0 \in]a; b[$, where it changes from D_k to D_l. Let $(x, y) : W_{kl} = U_{kl}^k \times U_{kl}^l \to \mathbb{C}^2$ and $t : \Delta \to \mathbb{C}$ be coordinates such that $h(x, y) = x \cdot y$ and $\vec{v} = \frac{\partial}{\partial t}$. Define $\gamma_x(\tau)$ (resp. $\gamma_y(\tau)$) to be $x(\gamma(\tau))$ (resp. $y(\gamma(\tau))$) for all $\tau \in [a; b]$ with $\gamma(\tau) \in U_{kl}^k$ (resp. $\gamma(\tau) \in U_{kl}^l$). Then define:

$$\int_\gamma \varphi := \lim_{\varepsilon \to 0} \left\{ \int_{\gamma^{\leq \tau_0 - \varepsilon}} \omega_k + \int_0^1 H\left(\xi, \log\left(\gamma_x(\tau_0 - \varepsilon)\gamma_y(\tau_0 + \varepsilon)\right)\right) d\xi + \int_{\gamma_{\geq \tau_0 + \varepsilon}} \omega_l \right\}.$$

(b): Let $\gamma : [a; b] \to Z_0$ be a path over $\vec{v}$, which meets the set of double points only once with parameter value $\tau_0 \in]a; b[$, where it stays in one component D_k. Then define:

$$\int_\gamma \varphi := \lim_{\varepsilon \to 0} \left\{ \int_{\gamma^{\leq \tau_0 - \varepsilon}} \omega_k + \int_{\gamma_{\geq \tau_0 + \varepsilon}} \omega_k \right\}.$$

(c): Now let $\gamma : [a; b] \to Z_0$ be a path over $\vec{v}$ such that $\gamma(a)$ and $\gamma(b)$ are no double points. Then there is a finite number of parameter values $\tau_1, \dots, \tau_N$ with $a < \tau_1 < \cdots < \tau_N < b$, which are mapped onto double points. Choose a $\tau_i^* \in]\tau_i; \tau_{i+1}[$ for $i = 1, \dots, N-1$ and let $\tau_0^* := a$ and $\tau_N^* := b$. Then define for $i = 1, \dots, N$ the paths $\gamma_i := \gamma_{|[\tau_{i-1}^*, \tau_i^*]}$ and we define:

$$\int_\gamma \varphi := \sum_{i=1}^N \int_{\gamma_i} \varphi.$$

(d): Finally let $\gamma : [a; b] \to Z_0$ be a path in Z_0 over $\vec{v}$ based at $\vec{w}$. The coordinate $p : D_0 \to \mathbb{C}$ allows us to consider D_0 as part of the complex plane. There, in the complex plane, we have the differential form (and not the symbol) $\frac{dp}{p}$. Let σ be the straight path $\tau \mapsto (1-\tau)$ in $\mathbb{C}$, then we define

$$\int\limits_{\gamma} \varphi := \int\limits_{\sigma\star\gamma\star\sigma^{-1}} \varphi.$$

Theorem 2.3. *For a path $\gamma : [a;b] \to Z_0$ over $\vec{v}$ based at $\vec{w}$ and a closed $\varphi \in A^1$ holds:*

(i) *$\int_\gamma \varphi$ is well-defined and finite,*

(ii) *$\int_\gamma \varphi$ does not depend on the choice of coordinates,*

(iii) *$\int_\gamma \varphi = 0$, if φ is exact.*

Before proving 2.3 let us recall the following well-known lemma.

Lemma 2.4. *On a local coordinate (U, z) of D_i around a double point p_{kl} any closed $\omega \in E^1(D_i \log P_i)$ can be written as $\omega = Res_{p_{kl}}\omega \cdot \frac{dz}{z} + \psi$, where ψ is a smooth 1-form in $E^1(U)$.*

Proof of 2.3: Let $\varphi = \sum_{i\geq 0} \omega_i + \sum_{[k<l]} K_{kl}\frac{dx}{x} + L_{kl}\frac{dy}{y} + H_{kl}d\xi$. To prove (i), we have to show that integrals of type (a) and (b) are well-defined, finite and do not depend on the choice of coordinates (note that (d) is similar to (b)). Assume that we are given coordinates x, y and t like in (a) and write

$$\omega_k = \mathrm{Res}_{p_{kl}}\omega_k\frac{dx}{x} + \psi_k \quad \text{ and } \quad \omega_l = \mathrm{Res}_{p_{kl}}\omega_l\frac{dx}{x} + \psi_l.$$

ad (a) That φ is closed implies $K' = \frac{\partial H}{\partial u}$, $L' = \frac{\partial H}{\partial u}$ and $\frac{\partial L}{\partial u} = \frac{\partial K}{\partial u}$. Let $H = H_0 + H_1 u + \cdots + H_m u^m$. Then $iH_i = K'_{i-1} = L'_{i-1}$ for $i = 1, \ldots, m$. Since $K(0,u)$, $K(1,u)$, $L(0,u)$, $L(1,u)$ are constants in $\mathbb{C}$ we have $K_i(0) = K_i(1) = L_i(0) = L_i(1) = 0$ for $i \geq 1$. Moreover $\mathrm{Res}_{p_{kl}}\omega_k = -\mathrm{Res}_{p_{kl}}\omega_l$ and because of $H_1 = K'_0 = L'_0$ we obtain $\int_0^1 H_1(\xi)d\xi = \mathrm{Res}_{p_{kl}}\omega_l$. Therefore we derived:

$$\int\limits_0^1 H(\xi,u)d\xi = \int\limits_0^1 H_0(\xi)d\xi + \mathrm{Res}_{p_{kl}}\omega_l \cdot u.$$

Now we may compute as follows: recall $\gamma(\tau_0) = p_{kl}$ and choose ε_0 such that $\gamma([\tau_0 - \varepsilon_0; \tau_0 + \varepsilon_0]) \subset W_{kl}$ and $\gamma_x([\tau_0 - \varepsilon_0; \tau_0])$ as well as $\gamma_y([\tau_0; \tau_0 + \varepsilon_0])$ lie within an open set of $\mathbb{C}$, where the logarithm is univalent. Then we find, when we split up $\omega_k = \mathrm{Res}_{p_{kl}}\,\omega_k\frac{dx}{x} + \psi_k$ and $\omega_l = \mathrm{Res}_{p_{kl}}\,\omega_l\frac{dy}{y} + \psi_l$:

$$\int\limits_{\gamma} \varphi = \lim_{\varepsilon\to 0}\left\{ \int\limits_{\gamma^{\leq\tau_0-\varepsilon}} \omega_k + \int\limits_0^1 H\left(\xi, \log\left(\gamma_x(\tau_0-\varepsilon)\gamma_y(\tau_0+\varepsilon)\right)\right) d\xi + \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} \omega_l \right\}$$

$$= \int\limits_{\gamma^{\leq\tau_0-\varepsilon_0}} \omega_k + \int\limits_{\gamma^{\leq\tau_0}_{\geq\tau_0-\varepsilon_0}} \psi_k + \int\limits_{\gamma_{\geq\tau_0+\varepsilon_0}} \omega_l + \int\limits_{\gamma^{\leq\tau_0+\varepsilon_0}_{\geq\tau_0}} \psi_l + \int\limits_0^1 H_0(\xi)d\xi$$

$$+ \lim_{\varepsilon\to 0}\left\{ \int\limits_{\gamma^{\leq\tau_0-\varepsilon}_{\geq\tau_0-\varepsilon_0}} \mathrm{Res}_{p_{kl}}\,\omega_k\frac{dx}{x} + \mathrm{Res}_{p_kl}\,\omega_l \log\left(\gamma_x(\tau_0-\varepsilon)\gamma_y(\tau_0+\varepsilon)\right) + \int\limits_{\gamma^{\leq\tau+\varepsilon_0}_{\geq\tau_0+\varepsilon}} \mathrm{Res}_{p_{kl}}\,\omega_l\frac{dy}{y} \right\}.$$

And then remark that

$$\lim_{\varepsilon\to 0}\left\{\int\limits_{\gamma^{\le\tau_0-\varepsilon}_{\ge\tau_0+\varepsilon_0}}\frac{dx}{x}-\log\left(\gamma_x(\tau_0-\varepsilon)\right)+\int\limits_{\gamma^{\le\tau_0+\varepsilon_0}_{\ge\tau_0+\varepsilon}}-\frac{dy}{y}+\log\left(\gamma_y(\tau_0+\varepsilon)\right)\right\}$$
$$=-\log\left(\gamma_x(\tau_0-\varepsilon_0)\right)+\log\left(\gamma_y(\tau_0+\varepsilon_0)\right).$$

This shows that the integral in (a) is finite.

Once we know that $\int_\gamma \varphi$ converges, we may write $\int_\gamma \varphi$ as

$$\int\limits_0^1 H_0 d\xi+\lim_{\varepsilon\to 0}\left\{\int\limits_{\gamma^{\le\tau_0-\varepsilon}}\omega_k+\mathrm{Res}_{p_{kl}}\omega_l\log\left(\gamma_x(\tau_0-\varepsilon)\gamma_y(\tau_0+\varepsilon)\right)+\int\limits_{\gamma_{\ge\tau_0+\varepsilon}}\omega_l\right\}$$

$$=\int\limits_0^1 H_0 d\xi+\lim_{\varepsilon\to 0}\left\{\int\limits_{\gamma^{\le\tau_0-\varepsilon}}\omega_k+\mathrm{Res}_{p_{kl}}\omega_l\log\left(\varepsilon^2\right)+\int\limits_{\gamma_{\ge\tau_0+\varepsilon}}\omega_l\right.$$
$$\left.+\mathrm{Res}_{p_{kl}}\omega_l\log\left(\frac{\gamma_x(\tau_0-\varepsilon)}{\varepsilon}\cdot\frac{\gamma_y(\tau_0-\varepsilon)}{\varepsilon}\right)\right\}.$$

Observe that

$$\lim_{\varepsilon\to 0}\left\{\frac{\gamma_x(\tau_0-\varepsilon)}{\varepsilon}\cdot\frac{\gamma_y(\tau_0+\varepsilon)}{\varepsilon}\right\}=-\dot\gamma_x^{\le\tau_0}\cdot\dot\gamma_y{}_{\ge\tau_0}$$

and

$$-\dot\gamma_x^{\le\tau_0}\cdot\dot\gamma_y{}_{\ge\tau_0}\frac{\partial}{\partial t}=\langle-\dot\gamma_{\le\tau_0},\dot\gamma_{\ge\tau_0}\rangle=\frac{\partial}{\partial t}.$$

Hence, the second summand vanishes, i. e. :

$$\int\limits_\gamma\varphi=\int\limits_0^1 H_0 d\xi+\lim_{\varepsilon\to 0}\left\{\int\limits_{\gamma^{\le\tau_0-\varepsilon}}\omega_k+\mathrm{Res}_{p_{kl}}\omega_l\log\left(\varepsilon^2\right)+\int\limits_{\gamma_{\ge\tau_0+\varepsilon}}\omega_l\right\}.$$

This expression obviously does not depend upon the choice of the coordinates (x,y) and t. If $\varphi=d(\sum_{i\ge 0}g_i+\sum_{[k<l]}P_{kl})=\sum_{i\ge 0}dg_i+\sum_{[k<l]}\frac{\partial P_{kl}}{\partial u}(\frac{dx}{x}+\frac{dy}{y})+P'_{kl}\,d\xi$, then

$$\begin{aligned}\int_\gamma\varphi &= P_{kl}(1,u)-P_{kl}(0,u)\\ &+ \lim_{\varepsilon\to 0}\left\{g_k(\gamma(\tau_0-\varepsilon))-g_k(\gamma(0))+g_l(\gamma(1))-g_l(\gamma(\tau_0+\varepsilon))\right\}\\ &= g_l(\gamma(1))-g_k(\gamma(0)).\end{aligned}$$

ad (b) The definition does not depend on the choice of coordinates. We just have to show that the integral converges.

$$\int\limits_{\gamma} \varphi = \int\limits_{\gamma^{\le \tau_0 - \varepsilon_0}} \omega_k + \int\limits_{\gamma^{\le \tau_0}_{\ge \tau_0 - \varepsilon_0}} \psi_k + \int\limits_{\gamma_{\ge \tau_0 + \varepsilon_0}} \omega_l + \int\limits_{\gamma^{\le \tau_0 + \varepsilon_0}_{\ge \tau_0}} \psi_l$$

$$+ \operatorname{Res}_{p_{kl}} \omega_k \lim_{\varepsilon \longrightarrow 0} \left\{ \int\limits_{\gamma^{\le \tau_0 - \varepsilon}_{\ge \tau_0 - \varepsilon_0}} \frac{dx}{x} + \int\limits_{\gamma^{\le \tau_0 + \varepsilon_0}_{\ge \tau_0 + \varepsilon}} \frac{dx}{x} \right\}.$$

And further

$$\lim_{\varepsilon \longrightarrow 0} \left\{ \int\limits_{\gamma^{\le \tau_0 - \varepsilon}_{\ge \tau_0 - \varepsilon_0}} \frac{dx}{x} + \int\limits_{\gamma^{\le \tau_0 + \varepsilon_0}_{\ge \tau_0 + \varepsilon}} \frac{dx}{x} \right\}$$

$$= \log(\gamma_x(\tau_0 + \varepsilon_0)) - \log(\gamma_x(\tau_0 - \varepsilon_0)) + \lim_{\varepsilon \longrightarrow 0} \left(\log(\gamma_x(\tau_0 - \varepsilon)) - \log(\gamma_x(\tau_0 + \varepsilon))\right).$$

Now is

$$\begin{aligned} & \lim_{\varepsilon \longrightarrow 0} \left\{\log(\gamma_x(\tau_0 - \varepsilon)) - \log(\gamma_x(\tau_0 + \varepsilon))\right\} \\ = & \lim_{\varepsilon \longrightarrow 0} \left\{ \log\left(\frac{\gamma_x(\tau_0 - \varepsilon)}{\varepsilon}\right) - \log\left(\frac{\gamma_x(\tau_0 + \varepsilon)}{\varepsilon}\right) \right\} \\ = & \log\left(-\dot{\gamma}_x^{\le \tau_0}\right) - \log\left(\dot{\gamma}_{x\ \ge \tau_0}\right) \\ = & \ 0 \quad \text{by the definition of a path over } \vec{v}. \end{aligned}$$

If φ is exact, we have $\int_\gamma \varphi = g_k(\gamma(1)) - g_k(\gamma(0))$. □

Remark 2.5. We see from the proof of Theorem 2.3 that we have in the case of an integral of type (a) like on page 23:

$$\int\limits_{\gamma} \varphi = \int\limits_{0}^{1} H_0 d\xi + \lim_{\varepsilon \longrightarrow 0} \left\{ \int\limits_{\gamma^{\le \tau_0 - \varepsilon}} \omega_k + \operatorname{Res}_{p_{kl}} \omega_l \log\left(\varepsilon^2\right) + \int\limits_{\gamma_{\ge \tau_0 + \varepsilon}} \omega_l \right\},$$

$$\begin{aligned} = & \lim_{\varepsilon \longrightarrow 0} \left\{ \int\limits_{\gamma^{\le \tau_0 - \varepsilon}} \omega_k - \operatorname{Res}_{p_{kl}} \omega_k \log\left(\gamma_x(\tau_0 - \varepsilon)\right) \right\} \\ + & \lim_{\varepsilon \longrightarrow 0} \left\{ \int\limits_{\gamma_{\ge \tau_0 + \varepsilon}} \omega_l + \operatorname{Res}_{p_{kl}} \omega_l \log\left(\gamma_y(\tau_0 + \varepsilon)\right) \right\} \\ + & \int\limits_{0}^{1} H_{kl}(\xi, 0) d\xi. \end{aligned}$$

Remark 2.6. Observe that if one considers in (a) the closed

$$\Xi := K \frac{dx}{x} + L \frac{dy}{y} + H\, d\xi \in \left(\bigwedge\nolimits^{\bullet} \left(\frac{dx}{x}, \frac{dy}{y}\right)[\log t] \otimes E^{\bullet}(\Delta^1) \right)^1$$

as element in

$$\left(\bigwedge\nolimits^{\bullet}(\frac{dx}{x},\frac{dy}{y})[\log x, \log y] \otimes E^{\bullet}(\Delta^1)\right)^1,$$

which is defined in the obvious way (for the precise definition see 4.1.1), by setting

$$d\,\log x = \frac{dx}{x} \qquad \text{and} \qquad d\log y = \frac{dy}{y},$$

then Ξ is even exact. If $P \in \mathbb{C}[\xi, \log x, \log y]$ is such that $dP = \Xi$, then

$$\int_{[0;1]} H(\xi, \log t) d\xi = P(1, \log x, \log y) - P(0, \log x, \log y).$$

Next we state that $\int_\gamma \varphi$ does not depend on the choice of a path γ within a homotopy class in $\pi_1(Z_{\vec{v}}, \vec{w})$.

Theorem 2.7. *Let $\varphi \in A^1$ be closed and let γ_0, $\gamma_1 : I \to Z_0$ be two paths over $\vec{v}$ based at $\vec{w}$, which are nearby homotopic. Then*

$$\int_{\gamma_0} \varphi = \int_{\gamma_1} \varphi.$$

We will prove the corresponding theorem for iterated integrals in general as Theorem 4.16. The Theorem 2.7 above is then just a special case of 4.16.

Theorem 2.8. *The integration of closed elements in A^1 along paths over $\vec{v}$ based at $\vec{w}$ defines an isomorphism*

$$H^1(A^\bullet) \xrightarrow{\cong} \mathrm{Hom}_{\mathbb{Z}}\left(\vec{J}/\vec{J}^{\,2}; \mathbb{C}\right).$$

Proof: Since we know that both $\mathbb{C}$-vector spaces have the same dimension, it suffices to show that the integration map is injective. Note that

$$\mathrm{Hom}_{\mathbb{Z}}\left(\vec{J}/\vec{J}^{\,2}; \mathbb{C}\right) \hookrightarrow \mathrm{Hom}_{\mathbb{Z}}\left(\vec{J}; \mathbb{C}\right) \cong \mathrm{Hom}_{\mathbb{Z}}\left(\pi_1(Z_{\vec{v}}, \vec{w}); \mathbb{C}\right).$$

Assume $\varphi = \sum_{i\geq 0} \omega_i + \sum_{[k<l]} K_{kl}\frac{dx}{x} + L_{kl}\frac{dy}{y} + H_{kl} d\xi \in A^1$ is closed and that

$$\int_\gamma \varphi = 0 \qquad \text{for any } [\gamma] \in \pi_1(Z_{\vec{v}}, \vec{w}).$$

We claim that φ is exact. First, observe that all residues of φ are necessarily zero, as one sees by integration along a path over $\vec{v}$ based at $\vec{w}$, which approaches a double point, turns once around this double point and returns as it came. Moreover all the ω_i are exact since the expression $g_i(p) := \int_{\vec{w}}^p \varphi$ for $p \in D_i$ makes sense and $dg_i = \omega_i$. Now define

$$f := \sum_{i\geq 0} g_i + \sum_{[k<l]} \int_0^\xi H(\xi, u) d\xi \in B^0$$

and note that it satisfies the compatibility condition and that $df = \varphi$. □

The Theorem 2.8 allows us to define an integral lattice $H^1(A^\bullet)_{\mathbb{Z}}$ within $H^1(A^\bullet)$ in the following way.

Definition 2.9. Define $H^1(A^\bullet)_{\mathbb{Z}}$ to be the lattice of those cohomology classes which correspond to $\mathrm{Hom}_{\mathbb{Z}}(\vec{J}/\vec{J}^{\,2}; \mathbb{Z})$ via the integration map. $H^1(A^\bullet)_{\mathbb{Q}}$ is then defined as $H^1(A^\bullet)_{\mathbb{Q}} := \mathrm{Hom}_{\mathbb{Z}}(\vec{J}/\vec{J}^{\,2}; \mathbb{Q}) \subset H^1(A^\bullet)$.

Remark: Note that in contrast to $H^1(A^\bullet)$ the lattice $H^1(A^\bullet)_{\mathbb{Z}}$ depends on the tangent vectors $\vec{v}$ and $\vec{w}$.

2.3. **The Hodge- and Weight Filtration.** Next we define a decreasing filtration $F^\bullet$ and an increasing filtration $W_\bullet$ on $A^\bullet$ by defining such filtrations on $B^\bullet$. On $\Lambda^\bullet$ we set

$$F^p\Lambda^\bullet := \bigoplus_{n+m\geq p} \bigwedge\nolimits^n (\frac{dx}{x},\ \frac{dy}{y})(\log t)^m$$

$$W_l\Lambda^\bullet := \bigoplus_{n+2\cdot m\leq l} \bigwedge\nolimits^n (\frac{dx}{x},\ \frac{dy}{y})(\log t)^m.$$

In particular we have: $\log t \in W_2 \cap F^1$.

Then define on $E^\bullet(\Delta^1)\otimes\Lambda^\bullet$ the filtrations $F^\bullet$ and $W_\bullet$ by:

$$\begin{aligned} F^p\left(E^\bullet(\Delta^1)\otimes\Lambda^\bullet\right) &:= E^\bullet(\Delta^1)\otimes F^p\Lambda^\bullet \\ W_l\left(E^\alpha(\Delta^1)\otimes\Lambda^\beta\right) &:= E^\alpha(\Delta^1)\otimes W_{l+\alpha}\Lambda^\beta. \end{aligned}$$

On $E^\bullet(D_i\log P_i)$ for $i>0$ we have the classical *Hodge filtration*:

$$F^pE^\bullet(D_i\log P_i) := \bigoplus_{n\geq p}\Gamma\left(\Omega^n(D_i\log P_i)\otimes_{\Omega^0(D_i)}\mathcal{E}^{0,\bullet-n}(D_i)\right)$$

and the *weight filtration* $W_\bullet$. This filtration $W_\bullet$ is given by:

$$W_l\ E^p(D_i\log P_i) = E^l(D_i\log P_i)\otimes E^{p-l}(D_i) \quad \text{with} \quad 0\leq l\leq p.$$

Finally, define for $i=0,\ldots,r-1$ on $\bigwedge_i^\bullet(\frac{dp}{p}) = \bigwedge^\bullet(\frac{dp}{p})$:

$$W_l\bigwedge\nolimits^\bullet(\frac{dp}{p}) := \bigoplus_{n\leq l}\bigwedge\nolimits^n(\frac{dp}{p})$$

$$F^p\bigwedge\nolimits^\bullet(\frac{dp}{p}) := \bigoplus_{n\geq p}\bigwedge\nolimits^n(\frac{dp}{p}).$$

Putting all these filtrations together we obtain filtrations $W_\bullet$ and $F^\bullet$ on $A^\bullet$. In particular, $W_\bullet$ and $F^\bullet$ allow us to define filtrations on the cohomology groups in the following way.

Definition 2.10. The *weight filtration* $W_\bullet$ on $H^\bullet(A^\bullet)$ is defined by:

$$W_{l+m}H^m(A^\bullet) := \operatorname{im}\left\{H^m(W_lA^\bullet)\to H^m(A^\bullet)\right\}.$$

And define the *Hodge filtration* $F^\bullet$ on $H^\bullet(A^\bullet)$ as:

$$F^pH^m(A^\bullet) := \operatorname{im}\left\{H^m(F^pA^\bullet)\to H^m(A^\bullet)\right\}.$$

On the $\mathbb{Q}$-vector space $H^1(A^\bullet)_{\mathbb{Q}}$, the weight filtration $W_\bullet$ on $H^1(A^\bullet)$ induces the filtration:

$$W_lH^1(A^\bullet)_{\mathbb{Q}} := W_lH^1(A^\bullet)\cap H^1(A^\bullet)_{\mathbb{Q}}.$$

Observe that $Gr_l^W\left(H^1(A^\bullet)_{\mathbb{Q}}\right) = \operatorname{im}\left\{W_lH^1(A^\bullet)_{\mathbb{Q}}\to Gr_l^WH^1(A^\bullet)\right\}$.

We want to finish this subsection by proving the following theorem.

Theorem 2.11.

$$(J_{\vec{v}\vec{w}}/J^2_{\vec{v}\vec{w}})^* := \left(H^1(A^\bullet)_{\mathbb{Z}},\ (H^1(A^\bullet)_{\mathbb{Q}},W_\bullet),\ (H^1(A^\bullet),W_\bullet,F^\bullet)\right)$$

is a $\mathbb{Z}$-MHS of possible weights 0, 1, and 2.

Remark 2.12. If one computes the weight spectral sequence $(A^\bullet,\ W_\bullet)$ one finds infinitely many E_1-terms, which are nonzero, each of which is an infinite dimensional pure Hodge structure. Nevertheless, the spectral sequence degenerates at E_2. In the language of [Hai87a] the E_1-term is the image under a certain De Rham functor applied to the finite dimensional E_1-terms of the spectral sequences of $(E^\bullet(D_i \log P_i), W_\bullet)$ for $i > r-1$, of $(\bigwedge_i^\bullet(\frac{dp}{p}), W_\bullet)$ for $i \leq r-1$ and of $(\Lambda^\bullet, W_\bullet)$ (cf. [Hai87a], the proof of (5.5.1)).

However, if one just wants to prove that $H^1(A^\bullet)$ is a MHS, the weight spectral sequence of $(A^\bullet,\ W_\bullet)$ suggests a shortcut which avoids going all the way through the spectral sequences. This is how we will prove 2.11.

Furthermore one could consider $(A^\bullet,\ W_\bullet,\ F^\bullet)$ as part of a mixed Hodge complex, which takes the integral structure (coming from integration along paths over $\vec{v}$ based at $\vec{w}$) into account. We omit that construction, since we do not need it for our further investigations.

Proof: Since the category of MHSs is abelian, we can prove Theorem 2.11 if we show that $W_2H^1(A^\bullet) = H^1(A^\bullet)$ and that there are exact sequences

(i) $0 \to Gr_2^W H^1(A^\bullet) \xrightarrow{\alpha} \bigoplus_{[k<l]} H^0(p_{kl})(-1) \xrightarrow{\beta} \bigoplus_{i>r-1} H^2(D_i) \to 0,$

(ii) $0 \to Gr_1^W H^1(A^\bullet) \xrightarrow{\cong} \bigoplus_{i>r-1} H^1(D_i) \to 0,$

(iii) $\bigoplus_{i\geq 0} H^0(D_i) \xrightarrow{\gamma} \bigoplus_{[k<l]} H^0(p_{kl}) \xrightarrow{\delta} Gr_0^W H^1(A^\bullet) \to 0,$

where all the maps are defined over $\mathbb{Q}$ and are strict with respect to the respective Hodge filtrations. $Gr_l^W H^1(A^\bullet)$ is then a Hodge structure of weight l since all the other terms are Hodge structures of appropriate weights.

Note that if we are given a $[\varphi] = [\sum_{i\geq 0} \omega_i + \sum_{[k<l]} K_{kl}\frac{dx}{x} + L_{kl}\frac{dy}{y} + H_{kl}d\xi] \in H^1(A^\bullet)$, then we can always write with complex numbers ρ_{kl}, $c_{kl} \in \mathbb{C}$

$$[\varphi] = [\sum_{i\geq 0} \omega_i + \sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}\,\Theta + c_{kl}d\xi],$$

where $\Theta := (1-\xi)\frac{dx}{x} - \xi\frac{dy}{y} - \log t\, d\xi \in W_1\left[\left(E^\bullet(\Delta^1)\otimes\Lambda^\bullet\right)^1\right]$.

This shows that $W_2H^1(A^\bullet) = H^1(A^\bullet)$ and the following two equivalences:

$$[\varphi] \in W_1H^1(A^\bullet) \Leftrightarrow \forall [k<l]:\ \rho_{kl} = 0 \text{ and } [\varphi] \in W_0H^1(A^\bullet) \Leftrightarrow \forall i \geq 0:\ \omega_i = 0.$$

ad (i): We identify $H^0(p_{kl})(-1)$ with $\mathbb{C}(-1)$ such that $H^0(p_{kl})(-1)_{\mathbb{Z}}$ becomes $\frac{1}{2\pi i}\mathbb{Z} \subset \mathbb{C}$ and $Gr_F^1\mathbb{C} = \mathbb{C}$. Then define the map α by

$$\alpha\left([\sum_{i\geq 0}\omega_i + \sum_{[k<l]}\frac{1}{2\pi i}\rho_{kl}\,\Theta]\right) := \sum_{[k<l]}\frac{1}{2\pi i}\rho_{kl}.$$

If $[\sum_{i\geq 0}\omega_i + \sum_{[k<l]}\frac{1}{2\pi i}\rho_{kl}\,\Theta]$ represents an element in $H^1(A^\bullet)_{\mathbb{Z}}$ then it is mapped to $H^0(p_{kl})(-1)_{\mathbb{Z}}$ under α, since all the numbers ρ_{kl} have to be integers (they are the values of an integral along a path around a double point).

Now define the map β in the following way. Given an element $\sum_{[k<l]}\frac{1}{2\pi i}\rho_{kl} \in \bigoplus_{[k<l]} H^0(p_{kl})(-1)$, choose a $\sum_{i\geq 0}\omega_i + \sum_{[k<l]}\frac{1}{2\pi i}\rho_{kl}\,\Theta \in A^1$ such that for $i > r-1$

all the $d\omega_i$ have no singularities[6], i. e. $d\omega_i \in E^2(D_i)$. Then define:

$$\beta\left(\sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}\right) := \sum_{i>r-1} [d\omega_i].$$

Note that α and β are well-defined and that they yield a short exact sequence (i), where all the maps are strict with respect to the filtrations $F^\bullet$.

ad (ii): The isomorphism sends $[\sum_{i\geq 0}\omega_i] \in W_1H^1(A^\bullet)$ to $\sum_{i\geq r-1}[\omega_i] \in \bigoplus_{i>0} H^1(D_i)$. All the remaining verifications are obvious.

ad (iii): The maps γ and δ are defined in the following way:

$$\begin{array}{ccccccc} \bigoplus\limits_{i\geq 0} H^0(D_i) & \xrightarrow{\gamma} & \bigoplus\limits_{[k<l]} H^0(p_{kl}) & \xrightarrow{\delta} & Gr_0^W H^1(A^\bullet) & \to 0 \\ \sum_{i\geq 0} g_i & \mapsto & \sum_{[k<l]}(g_l - g_k) & & & \\ & & \sum_{[k<l]} c_{kl} & \mapsto & [\sum_{[k<l]} c_{kl}\; d\xi] & . \end{array}$$

Note that the kernel of γ has dimension 1. Also this sequence is exact, defined over $\mathbb{Q}$ and respects the filtrations $F^\bullet$. This accomplishes the proof. □

Remark 2.13. Using the short exact sequences (i), (ii) and (iii) from the proof of 2.11 we may derive that, if $g(D_i)$ denotes the genus of D_i for $i > r-1$:

$$\begin{aligned} \dim_{\mathbb{C}}\ Gr_2^W H^1(A^\bullet) &= (\sum_{[k<l]} 1) - (\sum_{i>r-1} 1), \\ \dim_{\mathbb{C}}\ Gr_1^W H^1(A^\bullet) &= \sum_{i>r-1} 2g(D_i), \\ \dim_{\mathbb{C}}\ Gr_0^W H^1(A^\bullet) &= (\sum_{[k<l]} 1) - (\sum_{i\geq 0} 1) + 1. \end{aligned}$$

2.4. **The Variation of Mixed Hodge Structure is a Nilpotent Orbit.**

Now it is not difficult anymore to investigate the monodromy of the local system of nearby fundamental groups and to show that the variation of Hodge structure

$$\left(J_{\vec{v},\vec{w}}/J^2_{\vec{v},\vec{w}}\right)^*_{\vec{v}\in(T_0\Delta)^*}$$

is a nilpotent orbit of Hodge structure.

Remark 2.14. The term *nilpotent orbit of mixed Hodge structure* is a generalization (for one variable) of the definition of a nilpotent orbit of MHS given in [Sch73] and [CK82] (Definition 3.1); specifically, [CK82]: (3.1) (iv) is not satisfied. A nilpotent orbit of MHS will be defined properly in Proposition & Definition 2.17.

First of all, we define a nilpotent chain morphism

$$N : A^\bullet \to A^\bullet$$

by defining it in the following way on $B^\bullet$. Let N be the zero map on $\bigoplus_{i\leq r-1} \bigwedge_i^\bullet(\frac{dp}{p}) \oplus \bigoplus_{i>r-1} E^\bullet(D_i \log P_i)$ and define it on $E^\bullet(\Delta^1)\otimes\Lambda^\bullet$ as

$$\begin{array}{cccc} N: & E^\bullet(\Delta^1)\otimes\Lambda^\bullet & \to & E^\bullet(\Delta^1)\otimes\Lambda^\bullet \\ & \Xi & \mapsto & \frac{d}{d(-u)}\Xi = -\frac{d}{du}\Xi\ . \end{array}$$

---

[6] This is always possible. It can be done similarly to the following: define a bump-function $b : \mathbb{C} \to [0;1]$ around $0 \in \mathbb{C}$ such that $b(z) \equiv 1$ in a small neighbourhood of $0 \in \mathbb{C}$. Then $b(z)\rho\frac{dz}{z}$ has residue ρ and $d(b(z)\rho\frac{dz}{z}) \in E^2(\mathbb{C})$.

Note that N respects the compatibility relations and hence defines a chain map $N : A^\bullet \to A^\bullet$. Concerning the filtrations, N induces maps

$$N : W_{l+1}A^\bullet \to W_{l-1}A^\bullet \quad \text{and} \quad N : F^pA^\bullet \to F^{p-1}A^\bullet.$$

In particular this chain map induces a map $N : H^1(A^\bullet) \to H^1(A^\bullet)$ (note that we are committing 'abuse of notation' here).

Proposition 2.15. *The filtration* $W_\bullet$ *on* $H^1(A^\bullet)$ *and* $N : H^1(A^\bullet) \to H^1(A^\bullet)$ *satisfy* $N^2 = 0$ *and*

(i) $W_{-1}H^1(A^\bullet) = 0$
(ii) $W_0\ H^1(A^\bullet) = \operatorname{im}\{N : H^1(A^\bullet) \to H^1(A^\bullet)\}$
(iii) $W_1\ H^1(A^\bullet) = \ker\{N : H^1(A^\bullet) \to H^1(A^\bullet)\}$
(iv) $W_2\ H^1(A^\bullet) = H^1(A^\bullet)$.

Proof: First, recall that we can always write a $[\varphi] \in H^1(A^\bullet)$ as

$$[\varphi] = [\sum_{i\geq 0} \omega_i + \sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}\,\Theta + c_{kl}d\xi],$$

where the ω_i are closed forms in $E^1(D_i \log P)$, the c_{kl} are complex numbers and $\Theta := (1-\xi)\frac{dx}{x} - \xi\frac{dy}{y} - \log t\, d\xi \in W_1\,(E^\bullet \otimes \Lambda^\bullet)^1$. This shows $N^2 = 0$ and $W_{-1}H^1(A^\bullet) = 0$ as well as $W_2\ H^1(A^\bullet) = H^1(A^\bullet)$.

ad(ii): Since the sum of the residues for any closed form ω_i is zero, we have

$$W_0\ H^1(A^\bullet) = \left\{ [\sum_{[k<l]} c_{kl}d\xi] \right\},$$

$$\operatorname{im} N = \left\{ [\sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}d\xi] \middle| \sum_{[i<l]} \frac{1}{2\pi i}\rho_{il} - \sum_{[k<i]} \frac{1}{2\pi i}\rho_{ki} = 0 \text{ for all } i \right\}.$$

Obviously: $\operatorname{im}\{N : H^1(A^\bullet) \to H^1(A^\bullet)\} \subseteq W_0\ H^1(A^\bullet)$. Now let $[\sum_{[k<l]} c_{kl}d\xi]$ be in $W_0\ H^1(A^\bullet)$ and let $m := \sum_{[k<l]} 1$. Define complex numbers

$$g_i := \frac{1}{m}\left(\sum_{[k<i]} c_{ki} - \sum_{[i<l]} c_{il}\right).$$

Note that $\sum_{i\geq 0} g_i = 0$. It is easy to check that

$$[\sum_{[k<l]} c_{kl}d\xi] = [\sum_{[k<l]} (c_{kl} - (g_l - g_k))d\xi] \in \operatorname{im} N.$$

ad(iii): Observe that on the one hand

$$W_1\ H^1(A^\bullet) = \left\{ [\sum_{i>0} \omega_i + \sum_{[k<l]} c_{kl}d\xi] \right\}$$

and on the other hand

$$\ker N = \left\{ [\sum_{i\geq 0} \omega_i + \sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}\,\Theta + c_{kl}d\xi] \,\middle|\, [\sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}\ d\xi] = 0 \right\}.$$

Hence, we have to show: $[\sum_{[k<l]} \frac{1}{2\pi i}\rho_{kl}\ d\xi] = 0 \in H^1(A^\bullet) \Rightarrow$ all $\rho_{kl} = 0$. For simplicity define $R_{kl} := \frac{1}{2\pi i}\rho_{kl}$. Let $g_i \in \mathbb{C}$ be complex numbers such that $d\left(\sum_{i\geq 0} g_i + \sum_{[k<l]}\{g_k + \xi(g_l - g_k)\}\right) = \sum_{[k<l]} R_{kl}\ d\xi$, i e. $(g_l - g_k) = R_{kl}\ \forall [k<l]$. We show that the real parts, $\Re(R_{kl})$, of all R_{kl} are zero and leave it to the reader to show that the imaginary parts are zero too.

Let i_0 be an index such that $\Re(g_{i_0}) = \max_i \Re(g_i)$. As a consequence, all the real parts of residues at ω_{i_0}, that is the numbers $\Re(\mathrm{Res}_{p_{kl}}\,\omega_{i_0})$ for $\{k,l\} \ni i_0$ and $[k<l]$, are non negative ($\mathrm{Res}_{p_{i_0 l}}\,\omega_{i_0} = R_{i_0 l} = g_{i_0} - g_l$ and $\mathrm{Res}_{p_{k i_0}}\,\omega_{i_0} = -R_{k i_0} = g_{i_0} - g_k$). Now the sum of the residues of the form ω_{i_0} has to be zero. This can only happen if for all the indices i with $D_i \cap D_{i_0} \neq \emptyset$ holds: $g_i = g_{i_0}$. Since the system od curves D_i is connected, this shows $\Re(g_i) = \Re(g_{i_0})$ for all i and hence $\Re(R_{kl}) = 0$ for all $[k<l]$. □

Remark 2.16. Proposition 2.15 shows that $W_\bullet$ is *the weight filtration of N relative to $W_\bullet$* as defined in [SZ85] (2.5). The proof of Proposition 2.15 is trivial, when one uses Proposition (2.14) in [SZ85].

Let $T \in \mathrm{Aut}\left(H^1(A^\bullet)\right)$ be the monodromy of the local system

$$\left\{\left(J_{\vec{v},\vec{w}}/J^2_{\vec{v},\vec{w}}\right)^*_{\mathbb{Z}}\right\}_{\vec{v}\in(T_0\Delta)^*}.$$

Recall that by means of Theorem 1.5 the automorphism T can be identified with the monodromy of the local system $\{H^1(Z_t;\mathbb{Z})\}_{t\in\Delta^*}$.

A basic example of *a variation of mixed Hodge structure* is given by the following definition. For a complex number $\lambda \in \mathbb{C}^*$ and a nilpotent vectorspace endomorphism A define the endomorphism λ^A to be $e^{\log\lambda A} = \mathbf{1} + \log\lambda A + \frac{1}{2}\log^2\lambda A^2 + \cdots$.

Proposition & Definition 2.17. Let H be a $\mathbb{Z}$-MHS and $N : H \to H$ be a nilpotent endomorphism such that $N(F^pH) \subset F^{p-1}H$ and $N(W_{l+1}H) \subset W_{l-1}H$. Then for any $\lambda \in \mathbb{C}^*$, the triple

$$H_\lambda := \left(\lambda^{-N}H_{\mathbb{Z}}, (H_{\mathbb{Q}}, W_\bullet), (H_{\mathbb{C}}, W_\bullet, F^\bullet)\right)$$

is a MHS and the family of MHSs $\{H_\lambda\}_{\lambda\in\mathbb{C}^*}$ is called *nilpotent orbit of mixed Hodge structure.* □

In the next theorem we describe, how the lattice $(J_{\vec{v},\vec{w}}/J^2_{\vec{v},\vec{w}})^*_{\mathbb{Z}} = H^1(A^\bullet)_{\mathbb{Z}} \subset H^1(A^\bullet)$ moves, when we move $\vec{v}$ and $\vec{w}$.

Theorem 2.18. *$e^{-2\pi i\, N} = T$ and for λ, $\mu \in \mathbb{C}$ holds:*

$$\left(J_{\lambda\vec{v},\mu\vec{w}}/J^2_{\lambda\vec{v},\mu\vec{w}}\right)^*_{\mathbb{Z}} = \lambda^{-N}\left(J_{\vec{v},\vec{w}}/J^2_{\vec{v},\vec{w}}\right)^*_{\mathbb{Z}} \subset H^1(A^\bullet).$$

Proof: Note that $N^2 = 0$ implies $\lambda^N = \mathbf{1} + \log\lambda N$. Let $\eta = \lambda \times \mu : [0;1] \to (\mathbb{C}\setminus\mathbb{R}^{\leq 0}) \times (\mathbb{C}\setminus\mathbb{R}^{\leq 0})$ be a path with $\eta(0) = (1,1)$ and let $H : [0;1]\times[0;1] \to Z_0$ be a homotopy such that $H(\cdot,s)$ is a path over $\lambda(s)\vec{v}$ based at $\mu(s)\vec{w}$. Define $\gamma_{\vec{v},\vec{w}} := H(\cdot,0)$.

Using the definitions (a), (b), (c) and (d) on page 23, it is easy to compute for a closed $\varphi \in A^1$:

$$\int_{H(\cdot,s)}\varphi = \int_{\gamma_{\vec{v},\vec{w}}}\varphi + \log\lambda(s)\int_{\gamma_{\vec{v},\vec{w}}}N\varphi = \int_{\gamma_{\vec{v},\vec{w}}}\lambda(s)^N\varphi.$$

Suppose $\int_{\gamma_{\vec{v},\vec{w}}}\varphi$ is an integer, then $\int_{H(\cdot,s)}\lambda(s)^{-N}\varphi$ is an integer too. □

The following statement is a direct consequence of Theorem 2.15 and Theorem 2.18.

Theorem 2.19. *The family of mixed Hodge structures*

$$\left\{ \left(J_{\vec{v},\vec{w}} / J^2_{\vec{v},\vec{w}} \right)^* \right\}_{\vec{v} \in (T_0\Delta)^*}$$

is a nilpotent orbit of mixed Hodge structure. □

Consider the graph Γ, whose vertices are the components D_i, where two vertices D_k and D_l are connected by an edge if and only if $D_k \cap D_l \neq \emptyset$. Then we have the following corollary of Theorem 2.15 and Theorem 2.18.

Corollary 2.20. Γ is a tree if and only if $T = id$.

Proof: By Theorem 2.15 and Theorem 2.18 $T = id$ is equivalent to $N = 0$ or $\ker N = W_1 H^1(A^\bullet) = W_2 H^1(A^\bullet) = H^1(A^\bullet)$. We observed in Remark 2.13 that holds:

$$\dim_{\mathbb{C}} \; Gr_2^W H^1(A^\bullet) = (\sum_{[k<l]} 1) - (\sum_{i \geq 0} 1) + 1.$$

Since Γ is connected, we have: Γ is a tree $\Leftrightarrow$ #edges = #vertices − 1. □

2.5. **Appendix.** We still have to give the proof of Theorem 2.2.
Proof of 2.2: The only closed elements in A^0 are the constant compatible functions. Therefore: $H^0(A^\bullet) \cong H^0(Z_t; \mathbb{C}) \cong \mathbb{C}$.

We make use of the fact that the map $\overset{\circ}{h}{:}\overset{\circ}{Z} \to \overset{\circ}{\Delta}$ in 1.4.2 is a locally trivial fibration and prove: $H^\bullet(A^\bullet) \cong H^\bullet(\overset{\circ}{Z}_{(0,1)}; \mathbb{C})$.

Recall that we can build up $\overset{\circ}{Z}_{(0,1)}$ in the following way. For each p_{kl} with $[k < l]$ we have the open disks $U_{kl}^k \subset D_k$ and $U_{kl}^l \subset D_l$ around the double point p_{kl}. Now define

$$B_{kl} := \left(\overset{\circ}{U}{}_{kl}^{k} \amalg \overset{\circ}{U}{}_{kl}^{l} \right) \Big/ \sim \qquad ,$$

where "$\sim$" denotes the equivalence relation:

$$\left(R_1, e^{2\pi i \phi_1}\right) \sim \left(R_2, e^{2\pi i \phi_2}\right) \;:\Leftrightarrow\; R_1 = R_2 = 0 \text{ and } \phi_1 = -\phi_2.$$

Then ${U_{kl}^k}^* \amalg {U_{kl}^l}^*$ can be considered as subspace of B_{kl} and as subspace of $\coprod_{i \geq 0} D_i^* = \coprod_{i \geq 0} D_i \setminus P_i$. We construct the fiber $\overset{\circ}{Z}_{(0,1)}$ by glueing $\coprod_{[k<l]} B_{kl}$ and $\coprod_{i \geq 0} D_i^*$ together along the ${U_{kl}^k}^* \amalg {U_{kl}^l}^*$'s. Note that $(\coprod_{i \geq 0} D_i^*, \coprod_{[k<l]} B_{kl})$ is an excisive couple (cf. [Spa66] p. 188) for the space $\overset{\circ}{Z}_{(0,1)}$, which is to say that we may apply Mayer-Vietoris. Hence, there is an exact sequence

$$(2.3) \quad 0 \to H^0(\overset{\circ}{Z}_{(0,1)}) \to \bigoplus_{i \geq 0} H^0(D_i^*) \oplus \bigoplus_{[k<l]} H^0(B_{kl}) \to \bigoplus_{[k<l]} H^0({U_{kl}^k}^*) \oplus H^0({U_{kl}^k}^*)$$

$$\to H^1(\overset{\circ}{Z}_{(0,1)}) \to \bigoplus_{i \geq 0} H^1(D_i^*) \oplus \bigoplus_{[k<l]} H^1(B_{kl}) \to \bigoplus_{[k<l]} H^1({U_{kl}^k}^*) \oplus H^1({U_{kl}^k}^*) \to 0.$$

Now we compare the last terms in (2.3) with the last terms in (2.2). That is, we define isomorphisms φ, ψ such that

$$\begin{tikzcd}
H^1(B^\bullet) \arrow[r, "\Phi"] \arrow[d, "\varphi", "\cong"'] & H^1(C^\bullet) \arrow[r] \arrow[d, "\psi", "\cong"'] & H^2(A^\bullet) \arrow[r] \arrow[d] & 0 \\
\bigoplus_{i\geq 0} H^1(D_i^*) \oplus \bigoplus_{[k<l]} H^1(B_{kl}) \arrow[r] & \bigoplus_{[k<l]} H^1({U_{kl}^k}^*) \oplus H^1({U_{kl}^k}^*) \arrow[r] & 0
\end{tikzcd}$$

commutes. It is then a consequence of the 5-lemma that $H^2(A^\bullet) = 0$.

Denote by $\{\frac{dx}{x}\}$ (resp. $\{\frac{dy}{y}\}$) the cohomology class in $H^1(B_{kl})$, which has value $2\pi i$ on the positive generator of $H_1({U_{kl}^k}^*)$ (resp. $H_1({U_{kl}^l}^*)$) in $H_1(B_{kl})$. Then $\{\frac{dx}{x}\} = -\{\frac{dy}{y}\}$. We define an isomorphism

$$\begin{array}{llll}
\varphi : & H^1(B^\bullet) & \rightarrow & \bigoplus_{i\geq 0} H^1(D_i^*) \oplus \bigoplus_{[k<l]} H^1(B_{kl}) \\
& [\sum_{i\geq 0} \omega_i + \sum_{[k<l]} K\frac{dx}{x} + L\frac{dy}{y} + Hd\xi] & \mapsto & \sum_{i\geq 0} [\omega_i] + \sum_{[k<l]} K\{\frac{dx}{x}\} + L\{\frac{dy}{y}.\}
\end{array}$$

(Note that for closed $\sum_{i\geq 0} \omega_i + \sum_{[k<l]} K\frac{dx}{x} + L\frac{dy}{y} + Hd\xi$, the polynomial $K(\xi,u) - L(\xi,u)$ is constant, i. e. $K(\xi,u) - L(\xi,u) = K(0,0) - L(0,0) = K(1,0) - L(1,0)$. Therefore, $K\{\frac{dx}{x}\} + L\{\frac{dy}{y}\} = (K-L)\{\frac{dx}{x}\}$ makes sense.)

Define moreover the isomorphism

$$\begin{array}{llll}
\psi : & H^1(C^\bullet) & \rightarrow & \bigoplus_{[k<l]} H^1({U_{kl}^k}^*) \oplus H^1({U_{kl}^k}^*) \\
& \sum_{[k<l]} (a_k\frac{dx}{x} + b_k\frac{dy}{y}) + (a_l\frac{dx}{x} + b_l\frac{dy}{y}) & \mapsto & \sum_{[k<l]} (a_k - b_k)\{\frac{dx}{x}\} + (b_l - a_l)\{\frac{dy}{y}\}
\end{array}$$

and note that the above diagram commutes.

Finally, if we use the fact that the Euler characteristics in (2.2) on page 23 and (2.3) on page 33 are zero, we find:

$$\begin{aligned}
\dim_{\mathbb{C}} H^1(A^\bullet) = \dim_{\mathbb{C}} H^1(\overset{\circ}{Z}_{(0,1)}) &= 1 + \sum_{i\geq 0} \big(\dim_{\mathbb{C}} H^1(D_i^*) - 1\big) \\
&= r + 1 + \sum_{i\geq 0} \big(\dim_{\mathbb{C}} H^1(D_i) - 2\big) + \sum_{[k<l]} 2. \qquad \square
\end{aligned}$$

3. ITERATED INTEGRALS ON THE NEARBY FUNDAMENTAL GROUP

Let $I = \sum_{|J|\leq s} a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigoplus_{r=1}^{s} \bigotimes^r A^1$. As a generalization of the notion of closedness of an element in A^1 the notion of Chen-closedness for an element I like above will be given. Under the assumption that such an I is Chen-closed we will define an *iterated integral of I along a path γ over $\vec{v}$ (based at $\vec{w}$).*

First, for a sufficiently small $\varepsilon > 0$ and arbitrary $\varphi_1 \otimes \cdots \otimes \varphi_r \in \bigotimes^r A^1$, we define what we call the *ε-iterated integral* of $\varphi_1 \otimes \cdots \otimes \varphi_r$ along a path γ over $\vec{v}$ (based at $\vec{w}$) and will denote it by:

$$ {}^{\varepsilon}\!\!\int_{\gamma} \varphi_1 \cdots \varphi_r. $$

This definition depends on the choice of coordinates around the double points (and on ε). Moreover, in general the expression diverges when passing $\varepsilon \to 0$. But we can prove that for a *Chen-closed element* I and for a path γ over $\vec{v}$ based at $\vec{w}$ the limit

$$\lim_{\varepsilon\to 0} {}^{\varepsilon}\!\!\int_{\gamma} I = \lim_{\varepsilon\to 0} \sum_{|J|\leq s} a_J \; {}^{\varepsilon}\!\!\int_{\gamma} \varphi_{j_1}\cdots\varphi_{j_r}$$

exists and neither depends on the choice of coordinates around the double points nor on the choice of γ within a nearby homotopy class. This will be done in Section 4. The main tool for the proofs is a dga $A^\bullet_{kl}$, which we consider to be $A^\bullet$ localized in a sector at a double point p_{kl}.

These iterated integrals along paths over $\vec{v}$ based at $\vec{w}$ will give a way to describe $\mathrm{Hom}_{\mathbb{Z}}(\vec{J}/\vec{J}^{s+1},\mathbb{C})$ in terms of the dga $A^\bullet$, similar to Chen's π_1-De Rham theorem (cf. (4.4) in [Hai87c]). Finally, this leads to the definition of a MHS on $(\vec{J}/\vec{J}^{s+1})^*$ such that the short exact sequence dual to (2.1) becomes a short exact sequence of MHSs. For instance the reference [Hai87c] contains the necessary background on iterated (line) integrals on manifolds.

3.1. **Review on Differential Graded Algebras.** Here we are going to introduce a little of Chen's *Reduced Bar Construction*; only so much as we will need in the sequel. For a broader introduction to this subject we refer to [Che76] or to [Hai87a].

Let $\mathcal{A}^\bullet$ be a connected dga over $\mathbb{K}=\mathbb{R}$ or $\mathbb{C}$ with augmentation map $\epsilon:\mathcal{A}^\bullet\to\mathbb{K}$ and let $\mathcal{R}(\mathcal{A}^\bullet,\epsilon)$ be the sub-vector space of

$$\bigoplus_{r=0}^{\infty}\bigotimes^{r}\mathcal{A}^\bullet$$

generated by elements $R_i(u,f)$, $i=1,\dots,r$, which are defined for $f\in\mathcal{A}^0$ and $u=\varphi_1\otimes\cdots\otimes\varphi_r\in\bigotimes^r\mathcal{A}^\bullet$ by:

$$\begin{aligned}
R_1(u,f) :=& df\otimes\varphi_2\otimes\cdots\otimes\varphi_r-(f-\epsilon(f))\varphi_2\otimes\varphi_3\otimes\cdots\otimes\varphi_r, && (3.1)\\
R_i(u,f) :=& \varphi_1\otimes\cdots\otimes\varphi_{i-1}\otimes df\otimes\varphi_{i+1}\otimes\cdots\otimes\varphi_r \\
&+\varphi_1\otimes\cdots\otimes f\varphi_{i-1}\otimes\varphi_{i+1}\otimes\cdots\otimes\varphi_r \\
&-\varphi_1\otimes\cdots\otimes\varphi_{i-1}\otimes f\varphi_{i+1}\otimes\cdots\otimes\varphi_r \quad \text{for } 1<i<r, && (3.2)\\
R_r(u,f) :=& \varphi_1\otimes\cdots\otimes\varphi_{r-1}\otimes df+\varphi_1\otimes\cdots\otimes(f-\epsilon(f))\varphi_{r-1}. && (3.3)
\end{aligned}$$

We define two maps[7], the *internal* and the *combinatorial differential*,

$$d_I,\ d_C:\bigoplus_{r=0}^{\infty}\bigotimes^{r}\mathcal{A}^1\longrightarrow\bigoplus_{r=0}^{\infty}\bigotimes^{r}\mathcal{A}^\bullet$$

by giving their values on elements of the form $\varphi_1\otimes\cdots\otimes\varphi_s$ and extending $\mathbb{K}$-linearly:

$$\begin{aligned}
d_I(\varphi_1\otimes\cdots\otimes\varphi_s) &:= \sum_{i=1}^{s}\varphi_1\otimes\cdots\otimes\varphi_{i-1}\otimes d\varphi_i\otimes\varphi_{i+1}\otimes\cdots\otimes\varphi_s\,,\\
d_C(\varphi_1\otimes\cdots\otimes\varphi_s) &:= \sum_{i=1}^{s-1}\varphi_1\otimes\cdots\otimes\varphi_{i-1}\otimes\varphi_i\wedge\varphi_{i+1}\otimes\varphi_{i+2}\otimes\cdots\otimes\varphi_s.
\end{aligned}$$

Here the maps d_I and d_C shift the degrees in the following way:

$$d_I:\bigotimes^{r}\mathcal{A}^1\longrightarrow\left(\bigotimes^{r}\left[\mathcal{A}^1\oplus\mathcal{A}^2\right]\right)^{r+1},\ d_C:\bigotimes^{r}\mathcal{A}^1\longrightarrow\left(\bigotimes^{r-1}\left[\mathcal{A}^1\oplus\mathcal{A}^2\right]\right)^{r}.$$

---

[7] Define $\bigotimes^0\mathcal{A}^\bullet$ to be $\mathbb{K}$ and $d_C:\mathcal{A}^1\to\mathbb{K}$ to be the zero map.

Define

$$\mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon) := \left(\bigoplus\nolimits_{r=0}^{s} \bigotimes\nolimits^{r} \mathcal{A}^1\right) \cap \mathcal{R}(\mathcal{A}^\bullet,\epsilon)$$

and note that $(d_I + d_C)(\mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon)) \subseteq \mathcal{R}^s(\mathcal{A}^\bullet,\epsilon)$.

Definition 3.1. The sum $(d_I + d_C)$ induces a map

$$d_{\mathrm{Chen}} : \bigoplus_{r=0}^{\infty} \bigotimes^{r} \mathcal{A}^1 \Big/ \mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon) \longrightarrow \left(\bigoplus_{r=0}^{\infty} \bigotimes^{r} \mathcal{A}^\bullet\right) \Big/ \mathcal{R}(\mathcal{A}^\bullet,\epsilon),$$

the *Chen differential* and its kernel consists of what is called *Chen-closed elements.* We call an element $I \in \bigoplus_{r=0}^{\infty} \bigotimes^{r} \mathcal{A}^1$ *Chen-closed* if $d_{\mathrm{Chen}}([I]) = 0$. For each s the kernel of d_{Chen} gets a name

$$H^0B_s(\mathcal{A}^\bullet,\epsilon) := \ker\left\{ d_{\mathrm{Chen}} : \frac{(\bigoplus_{r=0}^{s} \bigotimes^{r} \mathcal{A}^1)}{\mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon)} \to \frac{(\bigoplus_{r=0}^{\infty} \bigotimes^{r} \mathcal{A}^\bullet)}{\mathcal{R}(\mathcal{A}^\bullet,\epsilon)} \right\}.$$

Let $H^0\bar{B}_s(\mathcal{A}^\bullet,\epsilon)$ consist of those elements in $H^0B_s(\mathcal{A}^\bullet,\epsilon)$ with zero constant term, i. e. define $H^0\bar{B}_s(\mathcal{A}^\bullet,\epsilon) := \{[I] \in H^0B_s(\mathcal{A}^\bullet,\epsilon) |\ I \in \bigoplus_{r=1}^{s} \bigotimes^{r} \mathcal{A}^1\}$. An element of $H^0B_s(\mathcal{A}^\bullet,\epsilon)$ respectively $H^0\bar{B}_s(\mathcal{A}^\bullet,\epsilon)$ represented by $\sum_J a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}$ is classically written as

$$\sum_J a_J\ [\varphi_{j_1}|\cdots|\varphi_{j_r}].$$

Observe that we have inclusions for all $s \geq 1$

$$H^0B_s(\mathcal{A}^\bullet,\epsilon) \subset H^0B_{s+1}(\mathcal{A}^\bullet,\epsilon) \quad \text{and} \quad H^0\bar{B}_s(\mathcal{A}^\bullet,\epsilon) \subset H^0\bar{B}_{s+1}(\mathcal{A}^\bullet,\epsilon).$$

Remark 3.2. In the general version of the 'Reduced Bar Construction' (cf. [Che76] or [Hai87a]) the vectorspaces $H^0B_s(\mathcal{A}^\bullet,\epsilon)$ and $H^0\bar{B}_s(\mathcal{A}^\bullet,\epsilon)$ appear as first cohomology groups of certain complexes, whence the notation.

The following four Lemmas will turn out to be useful later in Section 4. Let $\bar{\mathcal{A}}^\bullet$ be a connected sub-dga of $\mathcal{A}^\bullet$ such that $\bar{\mathcal{A}}^p = \mathcal{A}^p$ for $p > 1$ and

$$\mathcal{A}^1 = d\mathcal{A}^0 \oplus \bar{\mathcal{A}}^1. \tag{3.4}$$

The proof of the following two lemmas is similar to the proof of the Theorem on p.23 in [Che76].

Lemma 3.3. *Suppose* $I \in \bigoplus_{r=0}^{s} \bigotimes^{r} \mathcal{A}^1$. *Then there is an* $R \in \mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon)$ *and an* $\bar{I} \in \bigoplus_{r=0}^{s} \bigotimes^{r} \bar{\mathcal{A}}^1$ *such that* $I = \bar{I} + R$ *and*

$$d_{Chen}[I] = 0 \quad \Leftrightarrow \quad (d_I + d_C)(\bar{I}) = 0.$$

Proof: By finite induction on $\tilde{s}$ we prove that for every $\tilde{s}$ between s and 0 there is an $I^{\tilde{s}} \in \bigoplus_{r=0}^{\tilde{s}} \bigotimes^{r} \mathcal{A}^1$ and an $R_{\tilde{s}} \in \mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon)$ as well as an $\bar{I}_{\tilde{s}} \in \bigoplus_{r=0}^{s} \bigotimes^{r} \bar{\mathcal{A}}^1$ such that $I = \bar{I}_{\tilde{s}} + I^{\tilde{s}} + R_{\tilde{s}}$. First, for $\tilde{\mathbf{s}} = \mathbf{s}$ let $I^{\tilde{s}} = I$ and $\bar{I}_{\tilde{s}} = R_{\tilde{s}} = 0$.
$\tilde{\mathbf{s}} < \mathbf{s}$**:** Assume now that there is an $I^{\tilde{s}+1} \in \bigoplus_{r=0}^{\tilde{s}+1} \bigotimes^{r} \mathcal{A}^1$ and an $R_{\tilde{s}+1} \in \mathcal{R}_1^s(\mathcal{A}^\bullet,\epsilon)$ as well as an $\bar{I}_{\tilde{s}+1} \in \bigoplus_{r=0}^{s} \bigotimes^{r} \bar{\mathcal{A}}^1$ such that

$$I = \bar{I}_{\tilde{s}+1} + I^{\tilde{s}+1} + R_{\tilde{s}+1}.$$

Write $I^{\tilde{s}+1} = \sum_{|J|\leq \tilde{s}+1} a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}$ and let $\varphi_{j_\nu} = \bar{\varphi}_{j_\nu} + df_{j_\nu}$ be the decomposition according to (3.4). Then observe that

$$\begin{aligned} I^{\tilde{s}} := \sum_{|J|\leq \tilde{s}+1} a_J \quad & \{ df_{j_1} \otimes \varphi_{j_2} \otimes \cdots \otimes \varphi_{j_r} && - R_1(\varphi_{j_2} \otimes \cdots \otimes \varphi_{j_r}; f_{j_1}) \\ & + \bar{\varphi}_{j_1} \otimes df_{j_2} \otimes \cdots \otimes \varphi_{j_r} && - R_2(\bar{\varphi}_{j_1} \otimes \varphi_{j_3} \otimes \cdots \otimes \varphi_{j_r}; f_{j_2}) \\ & \qquad \vdots && \qquad \vdots \\ & + \bar{\varphi}_{j_1} \otimes \cdots \otimes \bar{\varphi}_{j_{r-1}} df_{j_r} && - R_r(\bar{\varphi}_{j_1} \otimes \cdots \otimes \bar{\varphi}_{j_{r-1}}; f_{j_r}) \} \end{aligned}$$

and $\bar{I}_{\tilde{s}} := \sum_{|J|\leq \tilde{s}+1} a_J\ \bar{\varphi}_{j_1} \otimes \cdots \otimes \bar{\varphi}_{j_r}$ as well as

$$\begin{aligned} R_{\tilde{s}} := \sum_{|J|\leq \tilde{s}+1} a_J \{ \quad & R_1(\varphi_{j_2} \otimes \cdots \otimes \varphi_{j_r}; f_{j_1}) \\ + \quad & R_2(\bar{\varphi}_{j_1} \otimes \varphi_{j_3} \otimes \cdots \otimes \varphi_{j_r}; f_{j_2}) \\ & \qquad \vdots \\ + \quad & R_r(\bar{\varphi}_{j_1} \otimes \cdots \otimes \bar{\varphi}_{j_{r-1}}; f_{j_r}) \} \end{aligned}$$

satisfy the induction hypothesis for $\tilde{s}$. This completes the induction. Finally let $\bar{I} := \bar{I}_0$ and note that $d_{\mathrm{Chen}}\left([\bar{I}\,]\right) = 0 \Leftrightarrow (d_I + d_C)\left(\bar{I}\,\right) = 0$. ☐

Lemma 3.4. *Suppose $I \in \bigoplus_{r=0}^{s} \bigotimes^{r} \mathcal{A}^1$ is Chen-closed. Then we can write*

$$I = \sum\nolimits_{|J|=r\leq s} a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r},$$

where for any J with $|J| = s$ and $a_J \neq 0$ either all $\varphi_{j_1}, \dots, \varphi_{j_s}$ are closed or at least one of them is exact.

Proof: Let $I \in \bigoplus_{r=0}^{s} \bigotimes^{r} \mathcal{A}^1$ be Chen-closed. By Lemma 3.3 we may assume without loss of generality that $I \in \bigoplus_{r=0}^{s} \bigotimes^{r} \bar{\mathcal{A}}^1$ and $(d_I + d_C)I = 0$. Write $I = \sum_J a'_J\ \varphi'_{j_1} \otimes \cdots \otimes \varphi'_{j_r}$ and consider the part of I, which is contained in the s-th tensor power of $\mathcal{A}^1$, say $T^s := \sum_{|J|=s} a'_J\ \varphi'_{j_1} \otimes \cdots \otimes \varphi'_{j_r}$, as element of $[\bigotimes^s \mathcal{A}^\bullet]^s$. The condition $(d_C + d_I)(\sum_J a'_J\ \varphi'_{j_1} \otimes \cdots \otimes \varphi'_{j_r}) = 0$ implies that T^s is a cocycle with respect to the differential of $(\bigotimes^s \mathcal{A}^\bullet,\ d)$. The Künneth formula ([Spa66], p. 228 and p. 247, Theorem 11) yields the short exact sequence (i. e. the isomorphism)

$$0 \longrightarrow \left[\bigotimes^{s} H^\bullet(\mathcal{A}^\bullet)\right]^s \longrightarrow H^s\left(\bigotimes^{s} \mathcal{A}^\bullet\right) \longrightarrow 0.$$

This is the reason why we can write T^s as $\sum_{|J|=s} a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}$, where for each J with $|J| = s$ either all forms $\varphi_{j_1}, \dots, \varphi_{j_s}$ are closed or at least one of them is exact. ☐

Suppose that $H^2(\mathcal{A}^\bullet) = 0$. Given closed elements $\varphi_1, \dots, \varphi_s \in \mathcal{A}^1$, the next proposition shows us, how we can find Chen-closed elements in $\bigoplus_{r=0}^{\infty} \bigotimes^r \mathcal{A}^1$ whose part in $\bigotimes^s \mathcal{A}^1$ is the tensor product $\varphi_1 \otimes \cdots \otimes \varphi_s$.

Lemma 3.5. *Suppose $H^2(\mathcal{A}^\bullet) = 0$ and assume $\varphi_1, \dots, \varphi_s$ are closed elements in $\mathcal{A}^1$. Then there exist elements $\varphi_{i_1,\dots,i_\nu} \in \mathcal{A}^1$ with $\nu > 1$ such that for all $\nu \geq 1$ (i. e. the forms $\varphi_1, \dots, \varphi_s$ included) and for any multi-index $(i_1, \dots, i_\nu) \in \{1, \dots, s\}^\nu$ holds:*

$$d\varphi_{i_1,\dots,i_\nu} + \sum_{k=i}^{\nu-1} \varphi_{i_1,\dots,i_k} \wedge \varphi_{i_{k+1},\dots,i_\nu} = 0.$$

The following elements in $\bigoplus_{r=1}^{\nu}\bigotimes^r\mathcal{A}^1$ *are closed under* (d_I+d_C)*:*

$$I=\sum_{r=1}^{\nu}\sum_{0<\alpha_1<\cdots<\alpha_{r-1}<\nu}\varphi_{i_1,\dots,i_{\alpha_1}}\otimes\varphi_{i_{\alpha_1+1},\dots,i_{\alpha_2}}\otimes\cdots\otimes\varphi_{i_{\alpha_{r-1}+1},\dots,i_s}.$$

Proof: The proof of the existence of $\varphi_{i_1,\dots,i_\nu}\in\mathcal{A}^1$ with $\nu>1$ goes by induction on ν. The computations are straightforward. □

3.2. **Definition of Iterated Integrals on the Nearby Fundamental Group.** Let us begin with the definition of an ε-iterated integral of a $\varphi_1\otimes\cdots\otimes\varphi_r\in\bigotimes^r A^1$ along a path over $\vec{v}$ (based at $\vec{w}$). Each of the $\varphi_\nu\in A^1$ can be written as

$$\varphi_\nu=\sum_{i\geq 0}\omega_i^{(\nu)}+\sum_{[k<l]}K_{kl}^{(\nu)}\frac{dx}{x}+L_{kl}^{(\nu)}\frac{dy}{y}+H_{kl}^{(\nu)}d\xi.$$

Choose around every double point p_{kl} coordinates $(x,y):W_{kl}=U_{kl}^k\times U_{kl}^l\to\mathbb{C}^2$ and a coordinate $t:\Delta\to\mathbb{C}$ such that $h_{|W_{kl}}(x,y)=x\cdot y$ and $\vec{v}=\frac{\partial}{\partial t}$. For the double point $p_0\in D_0$ we require moreover that the coordinate x (which we also called p) satisfies $\frac{\partial}{\partial x}=\frac{\partial}{\partial p}=\vec{w}$. Always, when we talk about ε-iterated integrals in this section, we will refer to this coordinate system.

As for the definition of line integrals in 2.2 we give the definition of iterated integrals in several steps.

(a): Let $\gamma:[a;b]\to Z_0$ be a path over $\vec{v}$, which meets the set of double points only once with parameter value $\tau_0\in]a;b[$, where it changes from D_k to D_l. Define $\gamma_x(\tau)$ (resp. $\gamma_y(\tau)$) to be $x(\gamma(\tau))$ (resp. $y(\gamma(\tau))$) for all $\tau\in[a;b]$ with $\gamma(\tau)\in U_{kl}^k$ (resp. $\gamma(\tau)\in U_{kl}^l$). To shorten the following formulas we abbreviate for $\varepsilon>0$ small enough:

$$\eta_\varepsilon^{(\nu)}:=H_{kl}^{(\nu)}\big(\xi,\log\left(\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)\right)\big)\,d\xi.$$

Then define [8]

$$\varepsilon\!\!\int_\gamma\varphi_1\cdots\varphi_r:=\sum_{0\leq\alpha\leq\beta\leq r}\int_{\gamma\leq\tau_0-\varepsilon}\omega_k^{(1)}\cdots\omega_k^{(\alpha)}\int_{[0;1]}\eta_\varepsilon^{(\alpha+1)}\cdots\eta_\varepsilon^{(\beta)}\int_{\gamma\geq\tau_0+\varepsilon}\omega_l^{(\beta+1)}\cdots\omega_l^{(r)}.$$

(b): Let $\gamma:[a;b]\to Z_0$ be a path over $\vec{v}$, which meets the set of double points only once with parameter value $\tau_0\in]a;b[$, where it stays in one component D_k. Then define:

$$\varepsilon\!\!\int_\gamma\varphi_1\cdots\varphi_r:=\sum_{0\leq\alpha\leq r}\int_{\gamma\leq\tau_0-\varepsilon}\omega_k^{(1)}\cdots\omega_k^{(\alpha)}\int_{\gamma\geq\tau_0+\varepsilon}\omega_k^{(\alpha+1)}\cdots\omega_k^{(r)}.$$

(c): Now let $\gamma:[a;b]\to Z_0$ be a path over $\vec{v}$ such that $\gamma(a)$ and $\gamma(b)$ are no double points. Then there is a finite number of parameter values $\tau_1,\dots,\tau_N$ with $a<\tau_1<\cdots<\tau_N<b$, which are mapped onto double points. Choose a $\tau_i^*\in]\tau_i;\tau_{i+1}[$ for $i=1,\dots,N-1$ and let $\tau_0^*:=a$ and $\tau_N^*:=b$. Then define

---

[8] Here and also later we make the convention that all the expressions with an integral sign, where from the upper indices the right one is smaller than the left one, like $\int_{[0;1]}\eta_\varepsilon^{(5)}\cdots\eta_\varepsilon^{(4)}$, take the value 1.

for $i = 1, \dots, N$ the paths $\gamma_i := \gamma\big|_{[\tau^*_{i-1}, \tau^*_i]}$ with which we define ${}^{\varepsilon}\!\!\int_{\gamma} \varphi_1 \cdots \varphi_r$ as

$$\sum_{0 \le \alpha_1 \le \cdots \le \alpha_{N-1} \le r} {}^{\varepsilon}\!\!\int\limits_{\gamma_1} \varphi_1 \cdots \varphi_{\alpha_1} {}^{\varepsilon}\!\!\int\limits_{\gamma_2} \varphi_{\alpha_1+1} \cdots \varphi_{\alpha_2} \cdots {}^{\varepsilon}\!\!\int\limits_{\gamma_N} \varphi_{\alpha_{N-1}+1} \cdots \varphi_{\alpha_r}.$$

(d): Finally, let $\gamma : [a; b] \to Z_0$ be a path in Z_0 over $\vec{v}$ based at $\vec{w}$. The coordinate $p : D_0 \to \mathbb{C}$ allows us to consider D_0 as part of the complex plane. There, in the complex plane, we have the differential form (and not the symbol) $\frac{dp}{p}$. Let like before σ be the straight path $\tau \mapsto (1 - \tau)$ in $\mathbb{C}$, then we define

$${}^{\varepsilon}\!\!\int\limits_{\gamma} \varphi_1 \cdots \varphi_r := {}^{\varepsilon}\!\!\int\limits_{\sigma \star \gamma \star \sigma^{-1}} \varphi_1 \cdots \varphi_r.$$

We leave it to the reader to prove the following proposition. It is a consequence of the corresponding formula for iterated integrals on manifolds (see [Hai87c], Prop. 2.9).

Proposition 3.6. *The definition (c) does not depend on the choice of the* τ^*_i. *Let* $\mathbf{p}, \mathbf{q}, \mathbf{r} \in Z_0 \backslash \bigcup_{[k<l]} \{p_{kl}\}$ *and let* $\alpha,\ \beta : [0; 1] \to Z_0$ *be paths over* $\vec{v}$, *where* $\alpha(0) = \mathbf{p}$, $\alpha(1) = \beta(0) = \mathbf{q}$ *and* $\beta(1) = \mathbf{r}$. *Then for forms* $\varphi_1, \dots, \varphi_r \in A^1$ *and* $\varepsilon > 0$ *small enough holds:*

$${}^{\varepsilon}\!\!\int\limits_{\alpha \star \beta} \varphi_1 \cdots \varphi_r = \sum_{0 \le m \le r} {}^{\varepsilon}\!\!\int\limits_{\alpha} \varphi_1 \cdots \varphi_m {}^{\varepsilon}\!\!\int\limits_{\beta} \varphi_{m+1} \cdots \varphi_r$$

□

This behaviour of iterated integrals under composition of paths will lead to the definition of $\mathfrak{T}$ in Section 4, motivated by Chen's theory of formal connections. The main result of Section 4 will be the proof the following theorem.

Theorem 4.1 *Let* $I = \sum_{|J| \le s} a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigoplus_{r=1}^{s} \bigotimes^r A^1$ *be a Chen-closed element and let* $\gamma : [a; b] \to Z_0$ *be a path over* $\vec{v}$ *based at* $\vec{w}$. *Then*

$$\lim_{\varepsilon \to 0} {}^{\varepsilon}\!\!\int\limits_{\gamma} \sum_{J} a_J\ \varphi_{j_1} \cdots \varphi_{j_r}$$

converges (4.13) and does not depend on the choice of coordinates (4.14). Moreover it does not depend upon the choice of γ *within a nearby homotopy class of paths over* $\vec{v}$ *based at* $\vec{w}$ *(4.16), i. e. we have a nearby homotopy functional:*

$$\int \sum_{J} a_J\ \varphi_{j_1} \cdots \varphi_{j_r} : \pi_1(Z_{\vec{v}}, \vec{w}) \to \mathbb{C}.$$

Finally, this homotopy functional is zero if $I \in \mathcal{R}^s_1(A^\bullet, a)$.

Theorem 4.1 assures that the following definition of iterated integrals of Chen-closed elements along paths over $\vec{v}$ makes sense.

Definition 3.7. Let $I = \sum_{|J| \leq s} a_J \; \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigoplus_{r=1}^{s} \bigotimes^{r} A^1$ be a Chen-closed element. Then *the iterated integral of* I *along a path* γ *over* $\vec{v}$ *based at* $\vec{w}$ is defined by:

$$\int\limits_{\gamma} \sum_{J} a_J \; \varphi_{j_1} \cdots \varphi_{j_r} := \lim_{\varepsilon \to 0} \left\{ \varepsilon\!\!\!\int\limits_{\gamma} \sum_{J} a_J \; \varphi_{j_1} \cdots \varphi_{j_r} \right\}.$$

3.3. **The π_1-De Rham Theorem for the Nearby Fundamental Group.** Denote by $P(Z_{\vec{v}}, \vec{w})$ the set of all paths over $\vec{v}$ based at $\vec{w}$. Note that we can consider an ε-iterated integral not just as a function on $P(Z_{\vec{v}}, \vec{w})$ but also on $\mathbb{C}P(Z_{\vec{v}}, \vec{w})$, the free abelian group generated by the elements of $P(Z_{\vec{v}}, \vec{w})$, by extending it linearly. Note that elements in $P(Z_{\vec{v}}, \vec{w})$ can be composed and this operation extends to $\mathbb{C}P(Z_{\vec{v}}, \vec{w})$.

Also the following proposition corresponds to a rule for iterated integrals on manifolds (see (2.10), (2.13) in [Hai87c]).

Proposition 3.8. *Let* $\alpha_1, \ldots, \alpha_s$ *be paths over* $\vec{v}$ *based at* $\vec{w}$ *and let* $\varphi_1, \ldots, \varphi_r \in A^1$ *with* $s \geq r$. *Then*

$$\varepsilon\!\!\!\!\int\limits_{(\alpha_1-1)\cdots(\alpha_s-1)} \varphi_1 \cdots \varphi_r = \begin{cases} \varepsilon\!\!\int_{\alpha_1} \varphi_1 \cdots \varepsilon\!\!\int_{\alpha_s} \varphi_s + V(\varepsilon), & \text{if } r = s \\ W(\varepsilon), & \text{if } r < s. \end{cases}$$

where $V(\varepsilon)$, $W(\varepsilon)$ *are functions* $V, W : \mathbb{R}^{>0} \to \mathbb{C}$ *converging to* 0 *for* $\varepsilon \to 0$.

Proof: We write $A(\varepsilon) \equiv B(\varepsilon)$ if $A(\varepsilon) - B(\varepsilon) = V(\varepsilon)$ for a function $V(\varepsilon)$ with $\lim_{\varepsilon \to 0} V(\varepsilon) = 0$. One can prove (and this is easy) the following formula by induction on k:

$$\varepsilon\!\!\!\!\int\limits_{(\alpha_1-1)\cdots(\alpha_k-1)\alpha_{k+1}\cdots\alpha_s} \varphi_1 \cdots \varphi_r$$

$$\equiv \sum_{\substack{k \leq N \leq s \\ 0<\nu_1<\cdots<\nu_k\leq\nu_{k+1}\leq\cdots\leq\nu_{N-1}\leq\nu_N=r}} \varepsilon\!\!\!\int\limits_{\alpha_1} \varphi_1 \cdots \varphi_{\nu_1} \varepsilon\!\!\!\int\limits_{\alpha_2} \varphi_{\nu_1+1} \cdots \varphi_{\nu_2} \cdots \varepsilon\!\!\!\int\limits_{\alpha_N} \varphi_{\nu_{N-1}+1} \cdots \varphi_{\nu_N}.$$

For $k > r$, the sum is zero and for $k = r = s$ the sum is: $\varepsilon\!\!\int_{\alpha_1} \varphi_1 \cdots \varepsilon\!\!\int_{\alpha_s} \varphi_s$. ☐

Recall that we defined an augmentation $a : A^\bullet \to \mathbb{C}$ (see 2.1). Therefore Definition 3.1 makes sense for the pair $(A^\bullet, a)$, whence we have the complex vector spaces

$$H^0 \bar{B}_s(A^\bullet, a) \subset H^0 B_s(A^\bullet, a).$$

Note in particular: $H^0 \bar{B}_1(A^\bullet, a) = H^1(A^\bullet)$. Theorem 4.1 and Proposition 3.8 yield a well-defined integration map

$$\int : H^0 B_s(A^\bullet, a) \longrightarrow \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w}) \big/ \vec{J}^{s+1}; \mathbb{C}).$$

The analogue for the nearby fundamental group of Chen's π_1-De Rham Theorem states that this map is an isomorphism.

Also the group ring $\mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w})$ comes with an augmentation map $\mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w}) \to \mathbb{Z}$. This augmentation has a natural section $\sigma : n \mapsto n\mathbf{1}$ which makes the short exact sequence

$$0 \to \vec{J} \big/ \vec{J}^{s+1} \to \mathbb{Z}\pi_1(Z_{\vec{v}}, \vec{w}) \big/ \vec{J}^{s+1} \overset{\overset{\sigma}{\dashleftarrow}}{\longrightarrow} \mathbb{Z} \to 0$$

split. This splitting gives a natural isomorphism

$$\mathrm{Hom}_{\mathbb{Z}}\left(\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})/\vec{J}^{s+1};\mathbb{Z}\right)\cong\mathrm{Hom}_{\mathbb{Z}}\left(\vec{J}/\vec{J}^{s+1};\mathbb{Z}\right)\oplus\mathbb{Z}$$

Via the integration map this decomposition corresponds with

$$H^0B_s(A^\bullet,a)\cong H^0\bar{B}_s(A^\bullet,a)\oplus\mathbb{C}.$$

Therefore it suffices to consider $\mathrm{Hom}_{\mathbb{Z}}(\vec{J}/\vec{J}^{s+1},\mathbb{Z})$ in order to describe the dual of $\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})/\vec{J}^{s+1}$.

Theorem 3.9. *Integration of Chen-closed elements in* $\bigoplus_{r=1}^{s}\bigotimes^{r}A^1$ *along paths over* $\vec{v}$ *based at* $\vec{w}$ *defines an isomorphism of complex vector spaces*

$$H^0B_s(A^\bullet,a)\xrightarrow{\cong}\mathrm{Hom}_{\mathbb{Z}}\left(\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})/\vec{J}^{s+1};\mathbb{C}\right).$$

Proof: We first indicate that for each $s\geq 1$ there is a short exact sequence

$$0\to H^0\bar{B}_{s-1}(A^\bullet,a)\to H^0\bar{B}_s(A^\bullet,a)\to\bigotimes^{s}H^1(A^1)\to 0.$$

In particular, the map

$$\begin{array}{ccc} H^0\bar{B}_s(A^\bullet,a) & \to & \bigotimes^s H^1(A^1)\\ \sum_J a_J[\varphi_{j_1}|\cdots|\varphi_{j_r}] & \mapsto & \sum_J a_J[\varphi_{j_1}]\otimes\cdots\otimes[\varphi_{j_r}]\end{array}$$

is well-defined and is surjective by Proposition 3.5. Its kernel is $H^0\bar{B}_{s-1}(A^\bullet,a)$ by Lemma 3.4. Then by Proposition 3.8 we have the commutative diagram:[9]

$$\begin{tikzcd}
0 \arrow[r] & H^0\bar{B}_{s-1}(A^\bullet,a) \arrow[r] \arrow[d] & H^0\bar{B}_s(A^\bullet,a) \arrow[r] \arrow[d] & \bigotimes^s H^1(A^1) \arrow[r] \arrow[d] & 0\\
0 \arrow[r] & (\vec{J}/\vec{J}^s)^* \arrow[r] & (\vec{J}/\vec{J}^{s+1})^* \arrow[r] & \bigotimes^s(\vec{J}/\vec{J}^{\,2})^* \arrow[r] & 0
\end{tikzcd}$$

The theorem can now be proved by induction on s and the use of the 5-lemma. The beginning of the induction is given by Theorem 2.8. □

Warning 3.10. Assume that we are given two closed 1-forms φ, $\psi\in A^1$,

$$\varphi=\sum_{i\geq 0}0+\sum_{[k<l]}c_{kl}d\xi\qquad\text{and}\qquad\psi=\sum_{i\geq 0}\omega_i+\sum_{[k<l]}c'_{kl}d\xi,$$

where all $\omega_i\in E^1(D_i)$. Then $\varphi\wedge\psi=0$ and hence is $\int\varphi\psi$ a nearby homotopy functional. Define $\psi_0=\sum_{i\geq 0}0+\sum_{[k<l]}c'_{kl}d\xi$. When we multiply φ and ψ (componentwise), we get the same as if we multiply φ and ψ_0. But nonetheless, in general holds: $\int\varphi\psi\neq\int\varphi\psi_0=0$. In general, there exists such φ and ψ for which there is a closed path α in one of the D_i's such that $\int_\alpha\omega_i\neq 0$ and a path u over $\vec{v}$ starting at $\vec{w}$ and ending at $\alpha(0)=\alpha(1)$ such that $\int_u\varphi\neq 0$. Then

$$\begin{aligned}\int_{u\star\alpha\star u^{-1}}\varphi\psi&=\int_u\varphi\int_\alpha\psi-\int_\alpha\varphi\int_u\psi+\int_\alpha\varphi\psi\\&=\int_u\varphi\int_\alpha\psi\end{aligned}\qquad\diamond$$

---

[9] The map $(\vec{J}^s/\vec{J}^{s+1})^*\to\bigotimes^s(\vec{J}/\vec{J}^2)^*$, which is given by the ring structure on $\mathbb{Z}\pi_1(Z_{\vec{v}},\vec{w})$, is surjective since we have the isomorphism $\pi_1(Z_{\vec{v}},\vec{w})\cong\pi_1(Z_t,\sigma_p(t))$ for some $t\in\Delta^*$ and $p\in D_0^*$. In $\pi_1(Z_t,\sigma_p(t))$ it is true that $(J^s/J^{s+1})^*\to\bigotimes^s(J/J^{\,2})^*$ is surjective.

3.4. **The Mixed Hodge Structure on $\vec{J}/\vec{J}^{s+1}$.** After having developed this theory of iterated integrals along paths over $\vec{v}$, we are ready now to define a mixed Hodge Structure on $\vec{J}/\vec{J}^{s+1}$. With all what we know about iterated integrals along paths over $\vec{v}$ yet, this is just an application of standard techniques in mixed Hodge theory.

3.4.1. *The Hodge and the Weight Filtration.* Let us use Theorem 3.9 to define a weight - and a Hodge filtration on $(\vec{J}/\vec{J}^{s+1})^*$.

On $\bigoplus_{r=1}^{s} \bigotimes^{r} A^1$ the *weight filtration* $W_\bullet$ is defined by

$$W_l \left(\bigoplus_{r=1}^{s} \bigotimes^{r} A^1 \right) := \bigoplus_{r=1}^{s} \bigoplus_{l_1+\cdots+l_r+r\leq l} W_{l_1}A^1 \otimes \cdots \otimes W_{l_r}A^1$$

and the *Hodge filtration* $F^\bullet$ by

$$F^p \left(\bigoplus_{r=1}^{s} \bigotimes^{r} A^1 \right) := \bigoplus_{r=1}^{s} \bigoplus_{p_1+\cdots+p_r\geq p} F^{p_1}A^1 \otimes \cdots \otimes F^{p_r}A^1.$$

These filtrations $W_\bullet$ and $F^\bullet$ on $\bigoplus_{r=1}^{s} \bigotimes^{r} A^1$ induce filtrations $W_\bullet$ and $F^\bullet$ on the subspace $\ker(d_I + d_C)$ and on $H^0\bar{B}_s(A^\bullet, a) \subset H^0B_s(A^\bullet, a)$ by:

$$\begin{aligned} W_l H^0 B_s(A^\bullet, a) &:= \operatorname{im}\left\{ W_l \ker(d_I + d_C) \to H^0\bar{B}_s(A^\bullet, a) \right\} \\ F^p H^0 B_s(A^\bullet, a) &:= \operatorname{im}\left\{ F^p \ker(d_I + d_C) \to H^0\bar{B}_s(A^\bullet, a) \right\}. \end{aligned}$$

Proposition 3.11. *The short exact sequences (with $s \geq 2$)*

$$0 \to H^0\bar{B}_{s-1}(A^\bullet, a) \xrightarrow{\iota} H^0\bar{B}_s(A^\bullet, a) \xrightarrow{pr} \bigotimes^{s} H^1(A^1) \to 0$$

are strict with respect to $W_\bullet$ and $F^\bullet$.

Proof: It is easy to see that pr is strict. We shall prove that ι is strict with respect to both filtrations.

Hodge filtration: Let $\bar{I} = \sum_J a_J\,[\varphi_{j_1}|\cdots|\varphi_{j_r}] \in F^pH^0\bar{B}_s(A^\bullet, a) \cap H^0\bar{B}_{s-1}(A^\bullet, a)$. Then there is a Chen-closed $I = \sum_{|J|\leq s} a_J \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in F^p \bigoplus_{r=1}^{s} \bigotimes^{r} A^1$ such that $\int I = \int \bar{I}$. Since $\bar{I} \in H^0\bar{B}_{s-1}(A^\bullet, a)$ we may assume by Lemma 3.4, Theorem 4.1 and Proposition 3.8 that for J with $|J| = s$ all the $\varphi_{j_1}, \ldots, \varphi_{j_s}$ are exact: $\varphi_{j_m} = df_{j_m}$ and $f_{j_m}(p_{01}) = 0$. We have by definition: $\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_s} \in F^p \bigotimes^{s} A^1$ $:\Leftrightarrow \exists (p_1, \ldots, p_s) \in \mathbb{Z}^s$ with $p_1 + \cdots + p_s \geq p : \varphi_{j_1} \in F^{p_1}, \ldots, \varphi_{j_s} \in F^{p_s}$. Now there is either one $p_m \geq 1$ (which implies $\varphi_{j_m} = 0$ as $A^0 \xrightarrow{d} A^1$ is strict with respect to $F^\bullet$, i. e. $dA^0 \cap F^1A^1 = 0$) or $p_1 = \cdots = p_s = 0$. Then

$$(f_{j_1} - a(f_{j_1}))\, \varphi_{j_2} \otimes \varphi_{j_3} \otimes \cdots \otimes \varphi_{j_s} \in F^p \bigotimes^{s-1} A^1.$$

Therefore, there is a Chen-closed $\tilde{I} \in F^p \bigoplus_{r=1}^{s-1} \bigotimes^{r} A^1$ such that $\int \bar{I} = \int \tilde{I}$.

Weight filtration: Let $\bar{I} \in W_lH^0\bar{B}_s(A^\bullet, a) \cap H^0\bar{B}_{s-1}(A^\bullet, a)$. Then there is a Chen-closed $I = \sum_{|J|\leq s} a_J \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in W_l \bigoplus_{r=1}^{s} \bigotimes^{r} A^1$ such that $\int I = \int \bar{I}$. Because of $\bar{I} \in H^0\bar{B}_{s-1}(A^\bullet, a)$ we may again assume by Lemma 3.4, Theorem 4.1 and Proposition 3.8 that for J with $|J| = s$ all the $\varphi_{j_1}, \ldots, \varphi_{j_s}$ are exact: $\varphi_{j_m} = df_{j_m}$ and $f_{j_m}(p_{01}) = 0$.

Note that if $df \in W_\ell A^1$ then $f \in W_{\ell+1}A^0$. This can be seen as follows. Write $f = \sum_{i\geq 0} g_i + \sum_{[k<l]} P_0 + P_1u + \cdots P_mu^m$. Now $f \in W_{\ell+1} \Leftrightarrow 2m \leq \ell + 1$ and

$df \in W_\ell \Leftrightarrow 2m-1 \le \ell$. We have by definition: $\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_s} \in W_l \bigotimes^s A^1 :\Leftrightarrow \exists (l_1, \dots, l_s) \in \mathbb{Z}^s$ with $l_1 + \cdots + l_s + s \le l : \varphi_{j_1} \in W_{l_1}, \dots, \varphi_{j_s} \in W_{l_s}$. Then

$$(f_{j_1} - a(f_{j_1}))\, \varphi_{j_2} \otimes \varphi_{j_3} \otimes \cdots \otimes \varphi_{j_s} \in W_l \bigotimes^{s-1} A^1 .$$

Hence, there is a Chen-closed $\tilde{I} \in W_l \bigoplus_{r=1}^{s-1} \bigotimes^r A^1$ such that $\int \tilde{I} = \int \bar{I}$. □

The main theorem of this section is the following.

Theorem 3.12. *For any $s \ge 1$ is*

$$(\vec{J}/\vec{J}^{s+1})^* := \Big((\vec{J}/\vec{J}^{s+1})^*_{\mathbb{Z}},\ ((\vec{J}/\vec{J}^{s+1})^*_{\mathbb{Q}}, W_\bullet)\ ((\vec{J}/\vec{J}^{s+1})^*_{\mathbb{C}}, W_\bullet, F^\bullet) \Big)$$

a mixed Hodge structure and the short exact sequence

$$0 \to (\vec{J}/\vec{J}^{s})^* \to (\vec{J}/\vec{J}^{s+1})^* \to H^1(A^\bullet)^{\otimes s} \to 0 \tag{3.5}$$

is a short exact sequence of MHSs.

Proof: Using Proposition 3.11 and 1.16 in [GS75], the proof of the theorem can be done by induction on s. □

3.5. **The Monodromy on $\vec{J}/\vec{J}^{s+1}$.** Recall that we defined in 2.4 the nilpotent chain-morphism

$$\begin{array}{rccc} N: & A^\bullet & \to & A^\bullet \\ & \phi & \mapsto & \frac{d}{d(-u)}\phi = -\frac{d}{du}\phi\ . \end{array}$$

with the properties $N(W_{l+1}A^\bullet) \subset W_{l-1}A^\bullet$ and $N(F^pA^\bullet) \subset F^{p-1}A^\bullet$.

Now we define a second nilpotent chain-morphism:

$$M : A^\bullet \to A^\bullet$$

by defining it first on $B^\bullet$ in the following way: Let M be the zero map on

$$\bigoplus_{i \le r-1} \bigwedge\nolimits_i^\bullet \Big(\frac{dp}{p}\Big) \oplus \bigoplus_{i > r-1} E^\bullet(D_i \log P_i) \oplus \bigoplus_{[0<k<l]} E^\bullet(\Delta^1) \otimes \Lambda^\bullet .$$

Only on the one component $E^\bullet(\Delta^1) \otimes \Lambda^\bullet$ for $(k,l) = (0,1)$ it is, like N, defined by

$$\begin{array}{rccc} M: & E^\bullet(\Delta^1) \otimes \Lambda^\bullet & \to & E^\bullet(\Delta^1) \otimes \Lambda^\bullet \\ & \Xi & \mapsto & \frac{d}{d(-u)}\Xi = -\frac{d}{du}\Xi\ . \end{array}$$

Also M satisfies $M(W_{l+1}A^\bullet) \subset W_{l-1}A^\bullet$ and $M(F^pA^\bullet) \subset F^{p-1}A^\bullet$.

Remark 3.13. Note that for a $Q(u) \in E^\bullet(\Delta^1) \otimes \bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})[u]$ and for a $\lambda \in \mathbb{C}^*$ always holds:

$$\lambda^N Q(u) = Q(u + \log \lambda).$$

Similar for M. Moreover, N and M define linear maps

$$N,\ M : \bigoplus_{r=1}^{s} \bigotimes^{r} A^\bullet \to \bigoplus_{r=1}^{s} \bigotimes^{r} A^\bullet, \tag{3.6}$$

which are defined on a generator $\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}$ by the Leibniz rule:

$$N(\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}) = \sum_{\nu=1}^{r} \varphi_{j_1} \otimes \cdots \otimes N\varphi_{j_\nu} \otimes \cdots \otimes \varphi_{j_r} \qquad \text{resp.}$$

$$M(\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}) = \sum_{\nu=1}^{r} \varphi_{j_1} \otimes \cdots \otimes M\varphi_{j_\nu} \otimes \cdots \otimes \varphi_{j_r}.$$

The proof of the following proposition is a standard computation:

Proposition 3.14. *For* $\lambda \in \mathbb{C}^*$ *and* $\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigotimes^r A^\bullet$ *holds:*

$$\lambda^N (\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}) = (\lambda^N \varphi_{j_1}) \otimes \cdots \otimes (\lambda^N \varphi_{j_r}) \quad \textit{and}$$
$$\lambda^M (\varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}) = (\lambda^M \varphi_{j_1}) \otimes \cdots \otimes (\lambda^M \varphi_{j_r}).$$

□

Both, N and M in (3.6) induce maps $N,\ M : H^0 \bar{B}_s(A^\bullet, a) \to H^0 \bar{B}_s(A^\bullet, a)$. The complex vector space $H^0 \bar{B}_s(A^\bullet, a)$ does neither depend on $\vec{v}$ nor on $\vec{w}$. Therefore, we can study, how the lattice

$$(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}})^*_{\mathbb{Z}} \subset H^0 \bar{B}_s(A^\bullet, a)$$

behaves, when we move $\vec{v} \in (T_0\Delta)^*$ and $\vec{w} \in (T_{p_0}D_0)^*$. Let T and S be the monodromies of the local systems

$$\left\{ \left(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}} \right)^*_{\mathbb{Z}} \right\}_{\vec{v}\in(T_0\Delta)^*} \text{ and } \left\{ \left(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}} \right)^*_{\mathbb{Z}} \right\}_{\vec{w}\in(T_{p_0}D_0)^*}$$

respectively. The main result of this subsection is the following theorem.

Theorem 3.15. $e^{-2\pi i N} = T$ *and* $e^{2\pi i M} = S$. *For* $\lambda,\ \mu \in \mathbb{C}^*$ *holds:*

$$\left(J_{\lambda\vec{v},\mu\vec{w}}/J^{s+1}_{\lambda\vec{v},\mu\vec{w}} \right)^*_{\mathbb{Z}} = \lambda^{-N} \mu^M \left(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}} \right)^*_{\mathbb{Z}} \subset H^0 \bar{B}_s(A^\bullet, a).$$

The proof of this theorem is given in 4.4. As a consequence of Theorem 3.15 we obtain the following.

Theorem 3.16. *For any* $s \geq 1$ *the family of mixed Hodge structures*

$$\left\{ \left(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}} \right)^* \right\}_{\vec{v}\in(T_0\Delta)^*}$$

is a nilpotent orbit of mixed Hodge structure. □

3.6. **The MHS on the Fundamental Group of the Central Fiber.** Let $\pi_1(Z_0, p_0)$ be the fundamental group of the central fiber and let $\mathbb{Z}\pi_1(Z_0, p_0)$ be its group ring with augmentation ideal J_0. In this subsection we define a MHS on the fundamental group of the central fiber. Apart from the contribution coming from the non compact disks D_i with $i = 1, \ldots, r-1$, the construction of this MHS is a special case of the general construction of a MHS on the fundamental group of a complex algebraic varietiy as introduced by Hain [Hai87a]. The main result here in this subsection is that the obvious group homomorphism

$$c : \pi_1(Z_{\vec{v}}, \vec{w}) \to \pi_1(Z_0, p_0)$$

induces inclusions of MHSs:

$$c^* : \left(J_0/J_0^{s+1} \right)^* \to \left(\vec{J}/\vec{J}^{s+1} \right)^*.$$

Most proofs are very similar to the proofs of the corresponding statements for $\pi_1(Z_{\vec{v}}, \vec{w})$ and are therefore left to the reader.

3.6.1. *A DGA of Differential Forms on the Central Fiber.* Now let us construct a dga $A_0^\bullet$ which computes the cohomology of Z_0 and allows to define the MHS on $\pi_1(Z_0, p_0)$. Define

$$B_0^\bullet := \mathbb{C} \oplus \bigoplus_{i>0} E^\bullet(D_i) \oplus \bigoplus_{[k<l]} E^\bullet(p_{kl}) \otimes E^\bullet(\Delta^1).$$

By setting $\mathbb{C} = \bigwedge^\bullet(\frac{dp}{p})$ and $E^\bullet(p_{kl}) = \Lambda^0$ we may consider $B_0^\bullet$ as sub-dga of the complex $B^\bullet$ as it was defined in 2.1. Now define the dga:[10]

$$A_0^\bullet := B_0^\bullet \cap A^\bullet.$$

Note that: $A_0^\bullet = W_0 A^\bullet$, i. e. $A_0^\bullet$ consists of exactly those elements in $A^\bullet$, which have no poles and no "$\log t = u$". The dga $A_0^\bullet$ inherits a Hodge- and a weight filtration $F^\bullet$ and $W_\bullet$ as well as an augmentation map $a : A_0^\bullet \to \mathbb{C}$ from $A^\bullet$.

Similarly to Theorem 2.2 one can prove:

Theorem 3.17. $H^\bullet(A_0^\bullet) \cong H^\bullet(Z_0; \mathbb{C})$. □

3.6.2. *Collapsing the Vanishing Cycles.* A path $\gamma : [0;1] \to Z_0$ over $\vec{v}$ based at $\vec{w}$ is in particular a path in D^+ based at p_0. We leave it to the reader to prove:

Proposition 3.18. *The obvious map*

$$c : \pi_1(Z_{\vec{v}}, \vec{w}) \to \pi_1(Z_0, p_0).$$

is a well-defined surjective group homomorphism.

3.6.3. *Iterated Integrals on the Central Fiber.* Since $A_0^\bullet$ is a sub-dga of $A^\bullet$ we may define:

Definition 3.19. Let $I = \sum_{|J| \le s} a_J \; \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigoplus_{r=1}^s \bigotimes^r A_0^1$ be Chen-closed and let $[\gamma] \in \pi_1(Z_0, p_0)$ be such that it is represented by a path γ over $\vec{v}$ based at $\vec{w}$. Then define

$$\int_{[c(\gamma)]} I := \int_\gamma I.$$

Now let J_0 be the augmentation ideal in the group ring $\mathbb{Z}\pi_1(Z_0, p_0)$. Similarly to Theorem 2.8 on page 27 one proves:

Theorem 3.20. *The integration of closed forms in A_0^1 along elements of $\pi_1(Z_0, p_0)$ defines an isomorphism*

$$H^1(A_0^\bullet) \xrightarrow{\cong} \operatorname{Hom}_{\mathbb{Z}} \left(J_0 / J_0^2; \mathbb{C} \right).$$

Since by definition the integration maps are compatible with $c : \pi_1(Z_{\vec{v}}, \vec{w}) \to \pi_1(Z_0, p_0)$, we have an inclusion

$$H^1(A_0^\bullet) \hookrightarrow H^1(A^\bullet).$$

Define for $R = \mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$ or $\mathbb{C}$:

$$H^1(A_0^\bullet)_R := H^1(A^\bullet)_R \cap H^1(A_0^\bullet) \text{ and } \left(J_0 / J_0^{s+1} \right)_R^* := \operatorname{Hom}_{\mathbb{Z}} \left(J_0 / J_0^{s+1}; R \right).$$

It follows from the proof of Theorem 2.11 on page 28 that holds:

---

[10] We avoid to introduce the language of simplicial and cosimplicial objects here. However except for the contribution from $D_0, \ldots, D_{r-1}$, the complex $A_0^\bullet$ is isomorphic to the de Rham complex of the (complex part of the) cosimplicial mixed Hodge complex on the simplicial variety Z_0.

Theorem 3.21.

$$(J_0/J_0^2)^* := \big(H^1(A_0^\bullet)_{\mathbb{Z}},\ (H^1(A_0^\bullet)_{\mathbb{Q}}, W_\bullet),\ (H^1(A_0^\bullet), W_\bullet, F^\bullet)\big) = W_1(\vec{J}/\vec{J}^2)^*$$

In particular, $(J_0/J_0^2)^*$ *is a* $\mathbb{Z}$*-MHS of possible weights 0 and 1.*

Moreover, one can prove (similarly to Theorem 3.9):

Theorem 3.22. *Integration of Chen-closed elements in* $\bigoplus_{r=1}^s \bigotimes^s A_0^1$ *along elements of* $\pi_1(Z_0, p_0)$ *defines an isomorphism of complex vector spaces*

$$H^0\bar{B}_s(A_0^\bullet, a) \xrightarrow{\cong} \operatorname{Hom}_{\mathbb{Z}}\left(J_0/J_0^{s+1}; \mathbb{C}\right).$$

□

Since the integration maps are compatible with c and since $\vec{J}/\vec{J}^{s+1} \to J_0/J_0^{s+1}$ is surjective, we find (the path σ in the definition **(d)** in 3.2 does not contribute anything to the integral since there are no residues):

Corollary 3.23. $H^0\bar{B}_s(A_0^\bullet, a) \hookrightarrow H^0\bar{B}_s(A^\bullet, a)$. □

Similarly to Theorem 3.12 one proves:

Theorem 3.24. *For any* $s \geq 1$ *is*

$$\left(J_0/J_0^{s+1}\right)^* := \left(\left(J_0/J_0^{s+1}\right)^*_{\mathbb{Z}},\ \left(\left(J_0/J_0^{s+1}\right)^*_{\mathbb{Q}}, W_\bullet\right)\ \left(\left(J_0/J_0^{s+1}\right)^*_{\mathbb{C}}, W_\bullet, F^\bullet\right)\right)$$

a mixed Hodge structure and the short exact sequence

$$0 \to \left(J_0/J_0^s\right)^* \to \left(J_0/J_0^{s+1}\right)^* \to H^1(A_0^\bullet)^{\otimes s} \to 0 \tag{3.7}$$

is a short exact sequence of MHSs.

Finally, let us state the following easy to prove theorem.

Theorem 3.25. *The map* $c : \pi_1(Z_{\vec{v}}, \vec{w}) \to \pi_1(Z_0, p_0)$ *induces an inclusion of MHSs*

$$\left(J_0/J_0^{s+1}\right)^* \hookrightarrow \left(\vec{J}/\vec{J}^{s+1}\right)^*.$$

□

4. Well-Definedness and Formal Connections

The main results of this section can be summarized in the following theorem. [11]

Theorem 4.1. *Let* $I = \sum_{|J|\leq s} a_J\ \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigoplus_{r=1}^s \bigotimes^r A^1$ *be a Chen-closed element and let* $\gamma : [a; b] \to Z_0$ *be a path over* $\vec{v}$ *based at* $\vec{w}$*. Then*

$$\lim_{\varepsilon \to 0} {}^{\varepsilon}\!\!\int_\gamma \sum_J a_J\ \varphi_{j_1} \cdots \varphi_{j_r}$$

converges (4.13) and does not depend on the choice of coordinates (4.14). Moreover it does not depend upon the choice of γ *within a nearby homotopy class of paths over* $\vec{v}$ *based at* $\vec{w}$ *(4.16), i. e. we have a nearby homotopy functional:*

$$\int \sum_J a_J\ \varphi_{j_1} \cdots \varphi_{j_r} : \pi_1(Z_{\vec{v}}, \vec{w}) \to \mathbb{C}.$$

Finally, this homotopy functional is zero if $I \in \mathcal{R}_1^s(A^\bullet, a)$*.*

---

[11] The numbers in brackets refer to the place, where the respective assertions are proved.

In general on a manifold one can express iterated integrals over Chen-closed linear combinations of tensor products of smooth 1-forms in three different ways as line integrals: First, one can pull back the iterated integral to the parameter interval according to the definition of iterated integrals. Second, one can express the iterated integrals as line integrals on the universal covering. And third, one can split up the iterated integral into pieces over parts of the path, which lie in simply connected open sets. In all three cases one transforms the general iterated integral into an iterated integral on a space, whose dga of differential forms is acyclic.

The idea to prove 4.1 is to express iterated integrals along paths over $\vec{v}$ in a similar way locally as line integrals along paths over $\vec{v}$. Away from the double points in Z_0 this can be done like described above. But around a double point the problem arises that if we restrict our dga $A^\bullet$ in the naive way to a dga living on an open neighbourhood of the double point, still the result will not be an acyclic complex. After all the dga $A^\bullet$ restricted to a double point computes the cohomology of the vanishing cycle at that double point.

Since a path over $\vec{v}$ locally at a double point is forced to approach the double point in a particular sector we can find simply connected open sets in Z_0, which contain all of this (local) path apart from the double point itself. If we are given a closed form φ in A^1 then we can find primitives of this form at least on those open sets. But since φ may have a simple pole in the double point, these primitives can have logarithmic singularities there.

Recall that reducing the length of an iterated integral over a Chen-closed element with differential forms in an acyclic complex can be done as follows: Choose a primitive of a closed form and multiply this function in an appropriate way with the other forms (cf. [Hai87c], Prop. (1.3), p. 252). When one takes an iterated integral over a composition of two paths, one can take primitives of this closed form over each of the paths and has to take care that their values on the junction point match.

For instance in the case of a path which passes only through one double point p_{kl} and changes from one component to another, we would like to decompose this path into three parts: two parts where it stays in one of the components and a third part which is represented by the 1-simplex Δ^1. (Observe that also in the definition (a) on page 23 we thought of Δ^1 as being part of the path γ.) But, when we now choose primitives on each of the components of a closed compatible 1-form, in general these primitives have logarithmic singularities at the junction points of the three parts of the path. Hence, we cannot make their values match. It will turn out that although we cannot make their values match, we can make their formal shapes or the type of their singularity match. This leads to a kind of *higher compatibility* and finally to the definition of the dga $A^\bullet_{kl}$. In this way we can then indeed locally express an iterated integral along a path γ over $\vec{v}$ as line integral of a closed form in A^1_{kl} along γ. This construction allows us to prove in the local case convergence, independence of local coordinates around the double points and invariance under nearby homotopies between paths over $\vec{v}$ of iterated integrals of Chen-closed elements.

A crucial role in all convergence considerations is played by the following classical fact (cf. [Cou34] III.2, p. 192).

Fact 4.2. For $\varepsilon_0 > 0$ let $h :]-\varepsilon_0, \varepsilon_0[\to \mathbb{R}$ be a smooth function that vanishes at 0. Then for any $m \in \mathbb{N}$ holds: $\lim_{\varepsilon \searrow 0} h(\varepsilon) \log^m \varepsilon = 0$.

Since iterated integrals are not additive on paths in the naive sense, these results do not carry over immediately to the corresponding assertions in the global situation. Here we first have to introduce an analogue to Chen's theory of *formal power series connections* in order to describe the relation between iterated integrals on a composition of two paths over $\vec{v}$ and iterated integrals over these paths seperately.

4.1. **Local Considerations.**

We start out by considering iterated integrals along paths which are defined locally in a neighbourhood of a double point.

4.1.1. *A Localization of $A^\bullet$ in a Sector at a Double Point.* Consider a double point p_{kl} in Z_0 with $[k < l]$. Recall that there are coordinates $(x, y) : W_{kl} = U^k_{kl} \times U^l_{kl} \to \mathbb{C}^2$ around this double point. Using these coordinates, we want to construct a dga $A^\bullet_{kl}$ for this double point p_{kl}.

Define $U^{k-}_{kl} := U^k_{kl} \setminus \mathbb{R}^{\le 0}$ and $U^{l-}_{kl} := U^l_{kl} \setminus \mathbb{R}^{\le 0}$. Note that the function $\log x$ (resp. $\log y$) can be defined univalently on U^{k-}_{kl} (resp. U^{l-}_{kl}). Now let $E^\bullet(U^{k-}_{kl} \log p_{kl})$ (resp. $E^\bullet(U^{l-}_{kl} \log p_{kl})$) be the sub-dga of $E^\bullet(U^{k-}_{kl})$ (resp. $E^\bullet(U^{l-}_{kl})$), which is generated as $E^\bullet(U^k_{kl})$ (resp. $E^\bullet(U^l_{kl})$)–modules by powers of $\log x$ (resp. $\log y$) and their derivatives. That is, we have embeddings of dga's

$$E^\bullet(U^k_{kl}) \otimes_{\mathbb{C}} \bigwedge\nolimits^\bullet(\frac{dx}{x})[\log x] \hookrightarrow E^\bullet(U^{k-}_{kl})$$

and

$$E^\bullet(U^l_{kl}) \otimes_{\mathbb{C}} \bigwedge\nolimits^\bullet(\frac{dy}{y})[\log y] \hookrightarrow E^\bullet(U^{l-}_{kl}),$$

whose images are $E^\bullet(U^{k-}_{kl} \log p_{kl})$ and $E^\bullet(U^{l-}_{kl} \log p_{kl})$. [12]

Since $E^\bullet(U^k_{kl})$ (resp. $E^\bullet(U^l_{kl})$) and $\bigwedge^\bullet(\frac{dx}{x})[\log x]$ (resp. $\bigwedge^\bullet(\frac{dy}{y})[\log y]$) have cohomology only in degree 0, the Künneth formula tells us that also $E^\bullet(U^{k-}_{kl} \log p_{kl})$ (resp. $E^\bullet(U^{l-}_{kl} \log p_{kl})$) have no cohomology except in degree 0. If $\{k, l\} = \{0, l\}$ then define

$$E^\bullet(U^{0-}_{0l} \log p_{01}) := \bigwedge\nolimits^\bullet(\frac{dp}{p})[\log p],$$

which are the polynomials in the variable $\log p$ and coefficients in $\bigwedge^\bullet(\frac{dp}{p})$, where the differential is given by the Leibniz rule, $d\,\frac{dp}{p} = 0$ and $d\,\log p = \frac{dp}{p}$.

Now let $\Lambda^\bullet_{kl} := \bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})[v, w]$ be the dga, which is defined as follows: Let $\Lambda^\bullet := \bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})[v, w]$ be the dga $\bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})[v, w]$ of polynomials in two variables v and w with coefficients in $\bigwedge^\bullet(\frac{dx}{x}, \frac{dy}{y})$, where the differential d is given by the Leibniz rule, $d\left(\frac{dx}{x}\right) = d\left(\frac{dy}{y}\right) = 0$ and

$$d\,v := \frac{dx}{x} \qquad \text{and} \qquad d\,w := \frac{dy}{y}.$$

Observe that there is an embedding

$$\bigwedge\nolimits^\bullet(\frac{dx}{x}, \frac{dy}{y})[u] \hookrightarrow \bigwedge\nolimits^\bullet(\frac{dx}{x}, \frac{dy}{y})[v, w]$$

---

[12] These maps are embeddings for the following reason: If $\sum_{\nu=0}^m g_\nu(x) \log^\nu x \equiv 0$ with $g_\nu \in E^0(U^k_{kl})$ it follows that all $g_\nu \equiv 0$ for $\nu = 0, \dots, m$, because for any $x_0 \in U^k_{kl}$ the polynomial $\sum_{\nu=0}^m g_\nu(x_0) w^\nu \in \mathbb{C}[w]$ has infinitely many zeroes: $\{\log x_0 + 2\pi i n | n \in \mathbb{Z}\}$.

by setting $u = v + w$. Note that $(\Lambda^\bullet_{kl}, d)$ only has cohomology in degree 0.

Similar like in the definition of $A^\bullet$ we define the dga $A^\bullet_{kl}$ as sub-dga of

$$B^\bullet_{kl} := E^\bullet(U^{k-}_{kl} \log p_{kl}) \oplus E^\bullet(U^{l-}_{kl} \log p_{kl}) \oplus E^\bullet(\Delta^1) \otimes_{\mathbb{C}} \Lambda^\bullet_{kl},$$

which contains all those elements that are subject to the following compatibility conditions generalizing the compatibility conditions in the definition of $A^\bullet$. Observe that $B^n_{kl} = 0$ for $n \geq 4$.

We shall use P_{kl}, K_{kl}, L_{kl}, H_{kl}, R_{kl}, S_{kl} and T_{kl} to denote elements in $\mathbb{C}[\xi, v, w]$. If the context is clear, we omit the indices k, l.

$\mathbf{A^0_{kl}}$: An element

$$f = \sum_{\nu=0}^m g_{k,\nu}(x) \log^\nu x + \sum_{\nu=0}^m g_{l,\nu}(y) \log^\nu y + P_{kl}(\xi, v, w) \in B^0_{kl}$$

is called *compatible*, iff holds:

$$P_{kl}(0, v, w) = \sum_{\nu=0}^m g_{k,\nu}(0) v^\nu \text{ and } P_{kl}(1, v, w) = \sum_{\nu=0}^m g_{l,\nu}(0) w^\nu.$$

$\mathbf{A^1_{kl}}$: We call an element in B^1_{kl},

$$\varphi = \sum_{\nu=0}^m \omega_{k,\nu} \log^\nu x + \sum_{\nu=0}^m \omega_{l,\nu} \log^\nu y + K \frac{dx}{x} + L \frac{dy}{y} + H\, d\xi$$

a *compatible* element iff it satifies:

$$\begin{array}{rcccl} K(0, v, w) = & \sum_{\nu=0}^m \operatorname{Res}_{p_{kl}} \omega_{k,\nu}\, v^\nu & , & L(0, v, w) = 0, \\ K(1, v, w) = & 0 & , & L(1, v, w) = \displaystyle\sum_{\nu=0}^m \operatorname{Res}_{p_{kl}} \omega_{l,\nu}\, w^\nu. \end{array}$$

$\mathbf{A^2_{kl}}$: Let us call an element

$$\phi = \Omega^{(k)} + \Omega^{(l)} + R\, d\xi \wedge \frac{dx}{x} + S\, d\xi \wedge \frac{dy}{y} + T \frac{dx}{x} \wedge \frac{dy}{y} \in B^2_{kl}$$

compatible iff holds:

$$T(0, v, w) = T(1, v, w) = 0.$$

$\mathbf{A^3_{kl}}$: And finally let $A^3_{kl} := B^3_{kl}$.

Note that $dA^i_{kl} \subset A^{i+1}_{kl}$ for $i \geq 0$ and that $A^\bullet_{kl}$ is a dga. Again, similar like in the definition of $A^\bullet$, we can give an alternative definition of $A^\bullet_{kl}$ as follows. Consider the surjective map of complexes

$$\Phi_{kl} : B^\bullet_{kl} \longrightarrow \Lambda^\bullet_{kl} \oplus \Lambda^\bullet_{kl},$$

which sends the element $\sum_{\nu=0}^m g_{k,\nu}(x) \log^\nu x + \sum_{\nu=0}^m g_{l,\nu}(y) \log^\nu y + P_{kl}(\xi, v, w) \in B^0_{kl}$ to $(P_{kl}(0, v, w) - \sum_{\nu=0}^m g_{k,\nu}(0) v^\nu) \oplus (P_{kl}(1, v, w) - \sum_{\nu=0}^m g_{l,\nu}(0) w^\nu)$ and

$$\sum_{\nu=0}^m \omega_{k,\nu} \log^\nu x + \sum_{\nu=0}^m \omega_{l,\nu} \log^\nu y + K(\xi, v, w) \frac{dx}{x} + L(\xi, v, w) \frac{dy}{y} + H(\xi, v, w) d\xi$$

in B^1_{kl} to
$$\left[\left(K(0,v,w)-\sum_{\nu=0}^{m}\operatorname{Res}_{p_{kl}}\omega_{k,\nu}\,v^{\nu}\right)\frac{dx}{x}+L(0,v,w)\frac{dy}{y}\right]$$
$$\oplus\left[K(1,v,w)\frac{dx}{x}+\left(L(1,v,w)-\sum_{\nu=0}^{m}\operatorname{Res}_{p_{kl}}\omega_{l,\nu}\,w^{\nu}\right)\frac{dy}{y}\right].$$
Moreover, Φ_{kl} maps
$$\Omega^{(k)}+\Omega^{(l)}+R(\xi,v,w)\,d\xi\wedge\frac{dx}{x}+S(\xi,v,w)\,d\xi\wedge\frac{dy}{y}+T(\xi,v,w)\,\frac{dx}{x}\wedge\frac{dy}{y}\in B^2_{kl}$$
to $T(0,v,w)\frac{dx}{x}\wedge\frac{dy}{y}\oplus T(1,v,w)\frac{dx}{x}\wedge\frac{dy}{y}$. It is easy to check that Φ is indeed a surjective map of complexes. Then we find:
$$A^{\bullet}_{kl}=\ker\Phi_{kl}$$
and with $C^{\bullet}_{kl}:=\Lambda^{\bullet}_{kl}\oplus\Lambda^{\bullet}_{kl}$ we get the short exact sequence
$$0\longrightarrow A^{\bullet}_{kl}\longrightarrow B^{\bullet}_{kl}\longrightarrow C^{\bullet}_{kl}\longrightarrow 0$$
and may conclude from the long exact cohomology sequence using $H^0(A^{\bullet}_{kl})=\mathbb{C}$ that $H^i(A^{\bullet}_{kl})=0$ for $i\geq 1$.

Note that there is an obvious dga-morphism: $A^{\bullet}\to A^{\bullet}_{kl}$.

4.1.2. *A Path Locally at a Double Point.* Consider a path over $\vec{v}$, which lies entirely in $U^{k-}_{kl}\cup\{p_{kl}\}\cup U^{l-}_{kl}$ for a double point p_{kl}. We want to extend the definitions (a), (b), (c) and (d) in 3.2 to the definition of ε-iterated integrals with forms in A^1_{kl} along this path. Again like before we define these ε-iterated integrals in several steps. At the end we let the iterated integral of a Chen-closed element with these forms along such a path be the limit of the sum of ε-iterated integrals.

Let $\varphi_1\otimes\cdots\otimes\varphi_r$ in $\bigotimes^r A^1_{kl}$, where we write each
$$\varphi_j=\tilde{\omega}^{(j)}_k+\tilde{\omega}^{(j)}_l+\Xi^{(j)}$$
with
$$\tilde{\omega}^{(j)}_k=\sum_{\nu=0}^{m}\omega^{(j)}_{k,\nu}\,\log^{\nu}x\qquad\text{and}\qquad\tilde{\omega}^{(j)}_l=\sum_{\nu=0}^{m}\omega^{(j)}_{l,\nu}\,\log^{\nu}y$$
and
$$\Xi^{(j)}=K^{(j)}\frac{dx}{x}+L^{(j)}\frac{dy}{y}+H^{(j)}d\xi.$$
We define now the iterated integral of $\varphi_1\otimes\cdots\otimes\varphi_r$ along a path over $\vec{v}$ in $U^{k-}_{kl}\cup\{p_{kl}\}\cup U^{l-}_{kl}$ as follows.

(a): Let $\gamma:[a;b]\to U^{k-}_{kl}\cup\{p_{kl}\}\cup U^{l-}_{kl}$ be a path over $\vec{v}$, which meets the set of double points only once with parameter value $\tau_0\in]a;b[$, where it changes from D_k to D_l. Define $\gamma_x(\tau)$ (resp. $\gamma_y(\tau)$) to be $x(\gamma(\tau))$ (resp. $y(\gamma(\tau))$). We abbreviate for $\varepsilon>0$ small enough:
$$\eta^{(j)}_{\varepsilon}:=H^{(j)}\big(\xi,\log\left(\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)\right)\big)\,d\xi.$$
Then define
$${}^{\varepsilon}\!\!\int_{\gamma}\varphi_1\cdots\varphi_r:=\sum_{0\leq\alpha\leq\beta\leq r}\int_{\gamma_{\leq\tau_0-\varepsilon}}\tilde{\omega}^{(1)}_k\cdots\tilde{\omega}^{(\alpha)}_k\int_{[0;1]}\eta^{(\alpha+1)}_{\varepsilon}\cdots\eta^{(\beta)}_{\varepsilon}\int_{\gamma_{\geq\tau_0+\varepsilon}}\tilde{\omega}^{(\beta+1)}_l\cdots\tilde{\omega}^{(r)}_l.$$

(b): Let $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ be a path over $\vec{v}$, which meets the set of double points only once with parameter value $\tau_0 \in]a;b[$, where it stays in one component D_k. Then define:

$$\varepsilon\!\!\int\limits_{\gamma} \varphi_1 \cdots \varphi_r := \sum_{0 \le \alpha \le r} \int\limits_{\gamma \le \tau_0 - \varepsilon} \tilde{\omega}_k^{(1)} \cdots \tilde{\omega}_k^{(\alpha)} \int\limits_{\gamma \ge \tau_0 + \varepsilon} \tilde{\omega}_k^{(\alpha+1)} \cdots \tilde{\omega}_k^{(r)}.$$

(c): Now let $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ be a path over $\vec{v}$ such that $\gamma(a)$ and $\gamma(b)$ are different from p_{kl}. Then there is a finite number of parameter values $\tau_1, \ldots, \tau_N$ with $a < \tau_1 < \cdots < \tau_N < b$, which are mapped onto p_{kl}. Choose a $\tau_i^* \in]\tau_i; \tau_{i+1}[$ for $i = 1, \ldots, N-1$ and let $\tau_0^* := a$ and $\tau_N^* := b$. Then we define $\gamma_i := \gamma\big|_{[\tau_{i-1}^*, \tau_i^*]}$ for $i = 1, \ldots, N$ and let: $\varepsilon\!\!\int_\gamma \varphi_1 \cdots \varphi_r :=$

$$\sum_{0 \le \alpha_1 \le \cdots \le \alpha_{N-1} \le r} \varepsilon\!\!\int\limits_{\gamma_1} \varphi_1 \cdots \varphi_{\alpha_1} \varepsilon\!\!\int\limits_{\gamma_2} \varphi_{\alpha_1+1} \cdots \varphi_{\alpha_2} \cdots \varepsilon\!\!\int\limits_{\gamma_N} \varphi_{\alpha_{N-1}+1} \cdots \varphi_{\alpha_r}.$$

(d): Finally assume that $p_{kl} = p_0$. Let $\gamma : [a;b] \to \{p_{01}\} \cup U_{01}^{1-}$ be a path over $\vec{v}$ starting (or ending) at $\vec{w}$. Like before, the coordinate $p : D_0 \to \mathbb{C}$ allows us to consider D_0 as part of the complex plane. There, in $\mathbb{C}$, we have the differential form (and not the symbol) $\frac{dp}{p}$. Let σ be the straight path $\tau \mapsto (1-\tau)$ in $\mathbb{C}$, then we define $\varepsilon > 0$ small enough

$$\varepsilon\!\!\int\limits_{\gamma} \varphi_1 \cdots \varphi_r := \varepsilon\!\!\int\limits_{\sigma \star \gamma} \varphi_1 \cdots \varphi_r \quad \left(\text{or} \ \varepsilon\!\!\int\limits_{\gamma \star \sigma^{-1}} \varphi_1 \cdots \varphi_r \right).$$

Note that if all the φ_j are even in the image of the obvious dga morphism $A^\bullet \to A_{kl}^\bullet$, then clearly these definitions coincide with the definitions (a), (b), (c) and (d) in 3.2.

Proposition 4.3. *Let $f \in A_{kl}^0$. Then for any path $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ over $\vec{v}$ holds:*

$$\lim_{\varepsilon \to 0} \varepsilon\!\!\int\limits_{\gamma} df = f(\gamma(b)) - f(\gamma(a)).$$

Proof: Let $\varphi = \tilde{\omega}_k + \tilde{\omega}_l + K\frac{dx}{x} + L\frac{dy}{y} + Hd\xi = df = d\left(G_k(x) + G_l(y) + P(\xi, v, w)\right)$ and write

$$\tilde{\omega}_k = \sum_{\nu=0}^{m} \omega_{k,\nu} \log^\nu x \qquad \text{and} \qquad \tilde{\omega}_l = \sum_{\nu=0}^{m} \omega_{l,\nu} \log^\nu y$$

as well as

$$G_k(x) = \sum_{\nu=0}^{m} g_{k,\nu}(x) \log^\nu x \qquad \text{and} \qquad G_l = \sum_{\nu=0}^{m} g_{l,\nu}(y) \log^\nu y.$$

We can decompose γ into paths γ_i, for $i = 1, \ldots, N$, i. e. $\gamma = \gamma_1 \star \cdots \star \gamma_N$ such that each of the paths γ_i meets the double point p_{kl} only once. Because line integrals are additive we have

$$\int_\gamma \varphi = \sum_{i=1}^{N} \int_{\gamma_i} \varphi,$$

provided that all the integrals on the right hand side converge. Therefore we may assume without loss of generality that γ meets the double point p_{kl} only once with parameter value $\tau_0 \in]a;b[$. We distinguish two cases.

A path traversing the double point: Suppose that γ changes components, say, it runs from D_k to D_l. Observe first that holds:

$$\int_{[0;1]} H(\xi, v, w) d\xi = P(1, v, w) - P(0, v, w).$$

Then we may compute – using the compatibility:

$$\begin{aligned}
&\int_{\gamma^{\leq \tau_0 - \varepsilon}} dG_k - P(0, \log \gamma_x(\tau_0 - \varepsilon), \log \gamma_y(\tau_0 + \varepsilon)) \\
=&G_k(\gamma_x(\tau_0 - \varepsilon)) - \sum_{\nu=0}^{m} g_{k,\nu}(0) \log^\nu \gamma_x(\tau_0 - \varepsilon) - G_k(\gamma_x(a)) \\
=&\sum_{\nu=0}^{m} (g_{k,\nu}(\gamma_x(\tau_0 - \varepsilon)) - g_{k,\nu}(0)) \log^\nu \gamma_x(\tau_0 - \varepsilon) - G_k(\gamma_x(a)).
\end{aligned}$$

With Fact 4.2 we derive

$$\lim_{\varepsilon \to 0} \int\limits_{\gamma^{\leq \tau_0 - \varepsilon}} \sum_{\nu=0}^{m} \omega_{k,\nu} \log^\nu x - P(0, \log \gamma_x(\tau_0 - \varepsilon), \log \gamma_y(\tau_0 + \varepsilon)) = -G_k(\gamma_x(a))$$

and similarly we find:

$$\lim_{\varepsilon \to 0} \int\limits_{\gamma^{\geq \tau_0 + \varepsilon}} \sum_{\nu=0}^{m} \omega_{l,\nu} \log^\nu y - P(1, \log \gamma_x(\tau_0 - \varepsilon), \log \gamma_y(\tau_0 + \varepsilon)) = G_l(\gamma_y(b)).$$

A path colliding with the double point: Suppose now that γ stays in one component D_k. Here we compute:

$$\int\limits_{\gamma^{\leq \tau_0 - \varepsilon}} \tilde{\omega}_k + \int\limits_{\gamma^{\geq \tau_0 + \varepsilon}} \tilde{\omega}_k = (G_k(\gamma_x(\tau_0 - \varepsilon)) - G_k(\gamma_y(\tau_0 + \varepsilon))) + (G_k(\gamma_x(b)) - G_k(\gamma_x(a)))$$

and the first summand tends to 0 as ε becomes small. □

Before we formulate and prove a generalization of the preceeding Proposition to iterated integrals of arbitrary length we need some preparation. To perform all the convergence considerations in the proof of Theorem 4.6 we need the following technical but elementary Lemma.

Lemma 4.4. *Let $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ be a path over $\vec{v}$ that meets the double point once with parameter value τ_0 and changes from D_k to D_l. Let $\tilde{\omega}_k^{(1)}, \ldots, \tilde{\omega}_k^{(r)} \in E^1(U_{kl}^{k-} \log p_{kl})$ and let $\tilde{\omega}_l^{(1)}, \ldots, \tilde{\omega}_l^{(r)} \in E^1(U_{kl}^{l-} \log p_{kl})$. Define for the polynomials $h^{(1)}, \ldots, h^{(r)} \in \mathbb{C}[\xi, v, w]$:*

$$\eta_\varepsilon^{(j)} := h^{(j)}\big(\xi, \log \gamma_x(\tau_0 - \varepsilon), \log \gamma_y(\tau_0 + \varepsilon)\big) d\xi.$$

Then there is a positive number C and a natural number N such that for all $\varepsilon > 0$ small enough holds:

$$\left|\int_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)}\cdots\tilde{\omega}_k^{(r)}\right| \leq C\left|\log^N \varepsilon\right| \tag{4.1}$$

$$\left|\int_{[0;1]} \eta_\varepsilon^{(1)}\cdots\eta_\varepsilon^{(r)}\right| \leq C\left|\log^N \varepsilon\right| \tag{4.2}$$

$$\left|\int_{\gamma\geq\tau_0-\varepsilon} \tilde{\omega}_l^{(1)}\cdots\tilde{\omega}_l^{(r)}\right| \leq C\left|\log^N \varepsilon\right|. \tag{4.3}$$

Also the corresponding assertion holds for a path γ over $\vec{v}$, which meets the double point once but stays in one component.

Proof: For the proof of (4.2) in Lemma 4.4, observe that

$$\int_{[0;1]} \left(h^{(1)}(\xi,v,w)d\xi\right)\cdots\left(h^{(r)}(\xi,v,w)d\xi\right)$$

is a polynomial, say $Q(v,w)$, in $\mathbb{C}[v,w]$ such that

$$\int_{[0;1]} \eta_\varepsilon^{(1)}\cdots\eta_\varepsilon^{(r)} = Q(\log\gamma_x(\tau_0-\varepsilon), \log\gamma_y(\tau_0+\varepsilon)).$$

Write $Q(v,w)=\sum_{m,n} a_{mn}v^m w^n$. Then we find $Q(\log\gamma_x(\tau_0-\varepsilon), \log\gamma_y(\tau_0+\varepsilon))$

$$=\sum_{m,n} a_{mn}\left(\log(\frac{\gamma_x(\tau_0-\varepsilon)}{\varepsilon})+\log\varepsilon\right)^m\left(\log(\frac{\gamma_y(\tau_0+\varepsilon)}{\varepsilon})+\log\varepsilon\right)^n.$$

Assertion (4.2) is true since:

$$\lim_{\varepsilon\to 0}\log\left(\frac{\gamma_x(\tau_0-\varepsilon)}{\varepsilon}\right) = -\log\dot{\gamma}_x(\tau_0) \quad\text{and}\quad \lim_{\varepsilon\to 0}\log\left(\frac{\gamma_y(\tau_0+\varepsilon)}{\varepsilon}\right) = \log\dot{\gamma}_y(\tau_0).$$

(4.3) and (4.1) are quite similar to prove. We show (4.1) and leave it to the reader to prove (4.3). Let $f_j(\tau)d\tau = \gamma_x^*\tilde{\omega}_k^{(i)}$ for $i=1,\dots,r$. Then

$$\begin{aligned}
&\left|\int\limits_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)}\cdots\tilde{\omega}_k^{(r)}\right| \\
\leq\ &\left|\int\limits_{a\leq\tau_1\leq\cdots\leq\tau_r\leq\tau_0-\varepsilon}\cdots\int f_1(\tau_1)\cdots f_r(\tau_r)d\tau_1\cdots d\tau_r\right| \\
\leq\ &\int_a^{\tau_0-\varepsilon}|f_1(\tau_1)|d\tau_1\cdot\cdots\cdot\int_a^{\tau_0-\varepsilon}|f_r(\tau_r)|d\tau_r
\end{aligned}$$

and if $\tilde{\omega}_k(j)=\sum_{\nu=0}^m A_\nu(x)\log^\nu x\frac{dx}{x}+B_\nu(x)\log^\nu xd\bar{x}$, then we find

$$f_j(\tau)=\sum_{\nu=0}^m\left\{\frac{A_\nu(\gamma_x(\tau))\log^\nu\gamma_x(\tau)}{\gamma_x(\tau)}\dot{\gamma}_x(\tau)+B_\nu(\gamma_x(\tau))\log^\nu\gamma_x(\tau)\overline{\dot{\gamma}_x(\tau)}\right\}.$$

There are A_ν, $B_\nu > 0$ such that $\sup_x |A_\nu(x)| \leq A_\nu$ and $\sup_x |B_\nu(x)| \leq b_\nu$. Moreover let $C_\nu := \int_a^{\tau_0} \left|\overline{\dot{\gamma}_x(\tau)}\right| d\tau$. Then

$$\begin{aligned}
&\int_a^{\tau_0-\varepsilon} |f_j(\tau)| d\tau \\
\leq\ & \sum_{\nu=0}^{m} \left\{ A_\nu \int\limits_a^{\tau_0-\varepsilon} \left| \frac{\log^\nu \gamma_x(\tau)}{\gamma_x(\tau)} \dot{\gamma}_x(\tau) \right| d\tau + B_\nu \int_a^{\tau_0-\varepsilon} \left| \log^\nu \gamma_x(\tau) \overline{\dot{\gamma}_x(\tau)} \right| \right\} \\
\leq\ & \sum_{\nu=0}^{m} \left\{ A_\nu \left| \frac{1}{\nu+1} \left(\log^{\nu+1} \gamma_x(\tau_0-\varepsilon) - \log^{\nu+1} \gamma_x(a) \right) \right| + B_\nu \left| \log^\nu \gamma_x(\tau_0-\varepsilon) \right| C_\nu \right\}
\end{aligned}$$

The assertion follows now from the differentiability of γ_x. □

The following proposition is an analogue to the corresponding laws for iterated integrals on manifolds. There it is an easy consequence of partial integration. Here we have to take the limit $\varepsilon \to 0$ into account.

Proposition 4.5. *Assume now that* $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ *is a path over* $\vec{v}$. *Let* $\varphi_1, \ldots, \varphi_s \in A_{kl}^1$ *be such that at least one of them is exact.*

If $\varphi_1 = df_1$, *then*

$$\lim_{\varepsilon\to 0} \left\{ {}^{\varepsilon}\!\!\int_\gamma df_1\varphi_2 \ldots \varphi_s - {}^{\varepsilon}\!\!\int_\gamma \big((f_1 - f_1(\gamma(a))\varphi_2\big)\varphi_3 \cdots \varphi_s \right\} = 0,$$

if $\varphi_\lambda = df_\lambda$ *for* $2 \leq \lambda \leq s$, *then*

$$\lim_{\varepsilon\to 0} \left\{ {}^{\varepsilon}\!\!\int_\gamma \varphi_1 \cdots df_\lambda \cdots \varphi_s - {}^{\varepsilon}\!\!\int_\gamma \varphi_1 \cdots (f_\lambda \varphi_{\lambda+1}) \cdots \varphi_s + {}^{\varepsilon}\!\!\int_\gamma \varphi_1 \cdots (f_\lambda \varphi_{\lambda-1}) \cdots \varphi_s \right\} = 0,$$

if $\varphi_s = df_s$, *then*

$$\lim_{\varepsilon\to 0} \left\{ {}^{\varepsilon}\!\!\int_\gamma \varphi_1 \cdots \varphi_{s-1} df_s - {}^{\varepsilon}\!\!\int_\gamma \varphi_1 \cdots \varphi_{s-2} \big(f_s - f_s(\gamma(b))\varphi_{s-1}\big) \right\} = 0.$$

Proof: The computations for the proof of these laws are straightforward but tedious. The idea is to express the iterated integrals of the proposition in terms of iterated integrals along paths lying completely in either one of the components or on the 1-simplex Δ^1. For the latter iterated integrals the corresponding laws hold. The complete calculations can be found in [Kae97], pp. 121-129.

Let us here just prove the first of the three laws. The proof of the others is comparable. Write

$$\varphi_j = \tilde{\omega}_k^{(j)} + \tilde{\omega}_l^{(j)} + K^{(j)} \frac{dx}{x} + L^{(j)} \frac{dy}{y} + H^{(j)} d\xi$$

and

$$f_1 = f = G_k(x) + G_l(y) + P(\xi, v, w),$$

where

$$\tilde{\omega}_k = \sum_{\nu=0}^{m} \omega_{k,\nu} \log^\nu x \qquad \text{and} \qquad \tilde{\omega}_l = \sum_{\nu=0}^{m} \omega_{l,\nu} \log^\nu y$$

and

$$G_k(x) = \sum_{\nu=0}^{m} g_{k,\nu}(x) \log^{\nu} x \qquad \text{and} \qquad G_l(y) = \sum_{\nu=0}^{m} g_{l,\nu}(y) \log^{\nu} y.$$

Before we show the first law for general paths γ over $\vec{v}$ in $U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ we consider the local situation at a double point first.

A path traversing the double point once: Suppose that $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ meets the double point p_{kl} once with parameter value$\tau_0 \in]a;b[$, where it changes from D_k to D_l. Again, we make use of the notation:

$$P_\varepsilon(\xi) := P(\xi, \log\gamma_x(\tau_0 - \varepsilon), \log\gamma_y(\tau_0 + \varepsilon))$$

and

$$\eta_\varepsilon^{(j)} := H^{(j)}(\xi, \log\gamma_x(\tau_0 - \varepsilon), \log\gamma_y(\tau_0 + \varepsilon))d\xi.$$

Note that: $dP_\varepsilon(\xi) = \eta_\varepsilon^{(\lambda)}$ in $E^\bullet(\Delta^1)$. Assume without loss of generality that $f(p) = f(\gamma(a)) = 0$. Then we compute with $\varphi_1 = df$:

$$\begin{aligned}
&\varepsilon\!\!\!\int\limits_{\gamma} (f\varphi_2)\varphi_3 \cdots \varphi_s \\
&= \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} (G_l\tilde{\omega}_l^{(2)})\tilde{\omega}_l^{(3)} \cdots \tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} \int\limits_{[0;1]} (P_\varepsilon\eta_\varepsilon^{(2)})\eta_\varepsilon^{(3)} \cdots \eta_\varepsilon^{(\beta)} \int_{\gamma^{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(\beta+1)} \cdots \tilde{\omega}_l^{(s)} \\
&+ \sum_{1<\alpha\leq\beta\leq s} \int\limits_{\gamma_{\leq\tau_0-\varepsilon}} (G_k\tilde{\omega}_k^{(2)})\tilde{\omega}_k^{(3)} \cdots \tilde{\omega}_k^{(\alpha)} \int\limits_{[0;1]} \eta_\varepsilon^{(\alpha+1)} \cdots \eta_\varepsilon^{(\beta)} \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(\beta+1)} \cdots \tilde{\omega}_l^{(s)} \\
&= \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(1)} \cdots \tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} \int\limits_{[0;1]} \eta_\varepsilon^{(1)} \cdots \eta_\varepsilon^{(\beta)} \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(\beta+1)} \cdots \tilde{\omega}_l^{(s)} \\
&+ \sum_{1<\alpha\leq\beta\leq s} \int\limits_{\gamma^{\leq\tau_0-\varepsilon}} \tilde{\omega}_k^{(1)} \cdots \tilde{\omega}_k^{(\alpha)} \int\limits_{[0;1]} \eta_\varepsilon^{(\alpha+1)} \cdots \eta_\varepsilon^{(\beta)} \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(\beta+1)} \cdots \tilde{\omega}_l^{(s)} \\
&+ G_l(\gamma_y(\tau_0+\varepsilon)) \int\limits_{\gamma_{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(2)} \cdots \tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} P_\varepsilon(0) \int\limits_{[0;1]} \eta_\varepsilon^{(2)})\eta_\varepsilon^{(3)} \cdots \eta_\varepsilon^{(\beta)} \int\limits_{\gamma^{\geq\tau_0+\varepsilon}} \tilde{\omega}_l^{(\beta+1)} \cdots \tilde{\omega}_l^{(s)}
\end{aligned}$$

$$
\begin{aligned}
&= \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(1)}\cdots\tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} \int\limits_{[0;1]} \eta_\varepsilon^{(1)}\cdots\eta_\varepsilon^{(\beta)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)} \\
&+ \sum_{1<\alpha\leq\beta\leq s} \int\limits_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)}\cdots\tilde{\omega}_k^{(\alpha)} \int\limits_{[0;1]} \eta_\varepsilon^{(\alpha+1)}\cdots\eta_\varepsilon^{(\beta)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)} \\
&+ \int\limits_{[0,1]} \eta_\varepsilon^{(1)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(2)}\cdots\tilde{\omega}_l^{(s)} + \int\limits_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(2)}\cdots\tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} \int\limits_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)} \int\limits_{[0;1]} \eta_\varepsilon^{(2)}\cdots\eta_\varepsilon^{(\beta)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)} \\
&+ [(G_l(\gamma_y(\tau_0+\varepsilon)) - P_\varepsilon(1)) + (P_\varepsilon(0) - G_k(\gamma_x(\tau_0-\varepsilon)))] \\
&\quad \cdot \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(2)}\cdots\tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} (P_\varepsilon(0) - G_k(\log\gamma_x(\tau_0-\varepsilon)) \\
&\quad \cdot \int\limits_{[0;1]} \eta_\varepsilon^{(2)}\cdots\eta_\varepsilon^{(\beta)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)} \\
&= \varepsilon\!\!\int\limits_{\gamma} \varphi_1\cdots\varphi_s \\
&+ [(G_l(\gamma_y(\tau_0+\varepsilon)) - P_\varepsilon(1)) + (P_\varepsilon(0) - G_k(\gamma_x(\tau_0-\varepsilon)))] \cdot \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(2)}\cdots\tilde{\omega}_l^{(s)} \\
&+ \sum_{1=\alpha<\beta\leq s} (P_\varepsilon(0) - G_k(\log\gamma_x(\tau_0-\varepsilon)) \cdot \int\limits_{[0;1]} \eta_\varepsilon^{(2)}\cdots\eta_\varepsilon^{(\beta)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)}
\end{aligned}
$$

With Fact 4.2 and Lemma 4.4 we find that

$$
\lim_{\varepsilon\to 0}\left\{ \varepsilon\!\!\int\limits_{\gamma} (f\varphi_2)\varphi_3\cdots\varphi_s - \varepsilon\!\!\int\limits_{\gamma} \varphi_1\ldots\varphi_s \right\} = 0.
$$

A path colliding once with the double point. Now let $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ be a path over $\vec{v}$ that meets the double point p_{kl} once with parameter value $\tau_0 \in]a;b[$ and stays in one component D_k. Assume again $f(p) = f(\gamma(a)) = 0$. Then we compute as follows.

$$
\varepsilon\!\!\int\limits_{\gamma} (f\varphi_2)\varphi_3\cdots\varphi_s
$$

$$
\begin{aligned}
&= \int\limits_{\gamma\geq\tau_0+\varepsilon} (G_k\tilde{\omega}_k^{(2)})\tilde{\omega}_k^{(3)}\cdots\tilde{\omega}_k^{(s)} \\
&+ \sum_{1\leq\alpha\leq s} \int\limits_{\gamma\leq\tau_0-\varepsilon} (G_k\tilde{\omega}_k^{(2)})\tilde{\omega}_k^{(3)}\cdots\tilde{\omega}_k^{(\alpha)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_k^{(\alpha+1)}\cdots\tilde{\omega}_k^{(s)} \\
&= \sum_{0\leq\alpha\leq s} \int\limits_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)}\cdots\tilde{\omega}_k^{(\alpha)} \int\limits_{\gamma\geq\tau_0+\varepsilon} \tilde{\omega}_k^{(\alpha+1)}\cdots\tilde{\omega}_k^{(s)} \\
&+ \left(G_k(\gamma_x(\tau_0+\varepsilon)) - \int\limits_{\gamma\leq\tau_0-\varepsilon} \tilde{\omega}_k^{(1)} \right) \\
&= \varepsilon\!\!\int\limits_{\gamma} (f\varphi_2)\varphi_3\cdots\varphi_s + \left(G_k(\gamma_x(\tau_0+\varepsilon)) - G_k(\gamma(\tau_0-\varepsilon))\right).
\end{aligned}
$$

This shows by Fact 4.2 and Lemma 4.4:

$$
\lim_{\varepsilon\to 0}\left\{ \varepsilon\!\!\int\limits_{\gamma} (f\varphi_2)\varphi_3\cdots\varphi_s - \varepsilon\!\!\int\limits_{\gamma} \varphi_1\ldots\varphi_s \right\} = 0.
$$

A path locally at a double point Assume now that $\gamma : [a;b] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ is a path over $\vec{v}$. Note that we can decompose γ into pieces each of which meets the double point once: $\gamma = \gamma_1 \star \cdots \star \gamma_N$. We saw already that for each of the paths γ_i with $i = 1, \ldots, N$ the assertion of the Proposition holds.

It remains to show that if for a path $\alpha : [0;1] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ over $\vec{v}$ and for a path $\beta : [0;1] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ over $\vec{v}$ with $\alpha(1) = \beta(0)$ the assertion of the proposition holds, that then the assertion is true for their composition $\alpha \star \beta$. This is a straightforward calculation similar to those above, which finally makes again use of 4.2 and 4.4. □

Theorem 4.6. *Let p, $q \in U_{kl}^{k-} \cup U_{kl}^{l-}$ and let $I_{kl} \in \bigoplus_{r=1}^{s} \bigotimes^{s} A_{kl}^{1}$ be a Chen-closed element. Then there is a "function" $f_{pq} \in A_{kl}^{0}$ such that for any path $\gamma : [a;b] \to Z_0$ over $\vec{v}$, which lies in $U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ and which starts in p and ends in q, holds:*

$$
\lim_{\varepsilon\to 0} \varepsilon\!\!\int\limits_{\gamma} I_{kl} = f_{pq}(q) - f_{pq}(p).
$$

Theorem 4.6 has the following immediate consequence.

Corollary 4.7. $\lim_{\varepsilon\to 0} \varepsilon\!\!\int_{\gamma} I_{kl}$ converges under the assumptions of Theorem 4.6.

Proof of 4.6: We will prove the theorem by induction on the maximal length s of tensor products in I_{kl}. For $s = 1$ this is the assertion of Proposition 4.3.
$s \geq 2$: Let $I_{kl} = \sum_J a_J\, \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r}$ be Chen-closed. According to Lemma 3.4 there is for each J with $|J| = s$ at least one closed and hence exact form $\varphi_{j_\lambda} = df_{j_\lambda}$.

We distinguish three cases:

(i) If $\lambda = 1$, then define $\tilde{I}_{j_1,\ldots,j_s} := (f_{j_1} - f_{j_1}(p))\, \varphi_{j_2} \otimes \cdots \otimes \varphi_{j_s} \in \bigotimes^{s-1} A_{kl}^{1}$.
(ii) If $1 < \lambda < s$ define $\tilde{I}_{j_1,\ldots,j_s} \in \bigotimes^{s-1} A_{kl}^{1}$ to be:

$$
\varphi_{j_1} \otimes \cdots \otimes f_{j_\lambda}\varphi_{j_{\lambda+1}} \otimes \cdots \otimes \varphi_{j_s} - \varphi_{j_1} \otimes \cdots \otimes f_{j_\lambda}\varphi_{j_{\lambda-1}} \otimes \cdots \otimes \varphi_{j_s}.
$$

(iii) Set $\tilde{I}_{j_1,\dots,j_s} := -\varphi_{j_1}\otimes\cdots\otimes(f_{j_s}-f_{j_s}(q))\,\varphi_{j_{s-1}}\in\bigotimes^{s-1}A^1_{kl}$, if $\lambda = s$.

Note that the definition of the $\tilde{I}_{j_1,\dots,j_s}$ only depends on p and q. Now

$$\tilde{I}_{kl} := \sum_{|J|=s} a_J\,\tilde{I}_{j_1,\dots,j_s} + \sum_{|J|\le s-1} a_J\,\varphi_{j_1}\otimes\cdots\otimes\varphi_{j_r}\in\bigoplus_{r=1}^{s}\bigotimes^{r}A^1_{kl}$$

is Chen-closed and by induction hypothesis there is a $f_{kl}\in A^0_{kl}$ such that $\int_\gamma \tilde{I}_{kl} = f_{pq}(q)-f_{pq}(p)$. Proposition 4.5 tells us: $\lim_{\varepsilon\to 0}\{\varepsilon\!\!\int_\gamma \varphi_{j_1}\dots\varphi_{j_s} - \varepsilon\!\!\int_\gamma \tilde{I}_{j_1,\dots,j_s}\} = 0$. And this implies $\int_\gamma I_{kl} = \int_\gamma \tilde{I}_{kl}$. □

4.2. **Formal Power Series Connections.** Iterated integrals are not additive on paths in the naive sense as we saw already in Proposition 3.6. However, once we know they satisfy rules (see 4.8) similar to those in Proposition 4.5, it is possible to give Proposition 3.6 an interpretation of a homomorphism. The appropriate language to do this is Chen's theory of *formal power series connections* (cf. [Che77a], [Che73] and [Che75a]). Here we introduce an anologous language for integration with forms in $A^\bullet_{kl}$ and $A^\bullet$.

But first let us state the global version of Proposition 4.5. This is necessary in order to apply the reduced bar construction.

Proposition 4.8. *Assume that $\gamma : [a;b]\to Z_0$ is a path over $\vec{v}$ (based at $\vec{w}$). Let $\varphi_1,\dots,\varphi_s\in A^1$ be such that at least one of them is exact.*

If $\varphi_1 = df_1$, then

$$\lim_{\varepsilon\to 0}\left\{\varepsilon\!\!\int_\gamma df_1\varphi_2\dots\varphi_s - \varepsilon\!\!\int_\gamma \big((f_1-f_1(\gamma(a))\varphi_2\big)\varphi_3\cdots\varphi_s\right\} = 0,$$

if $\varphi_\lambda = df_\lambda$ for $2\le\lambda\le s$, then

$$\lim_{\varepsilon\to 0}\left\{\varepsilon\!\!\int_\gamma \varphi_1\cdots df_\lambda\cdots\varphi_s - \varepsilon\!\!\int_\gamma \varphi_1\cdots(f_\lambda\varphi_{\lambda+1})\cdots\varphi_s + \varepsilon\!\!\int_\gamma \varphi_1\cdots(f_\lambda\varphi_{\lambda-1})\cdots\varphi_s\right\} = 0,$$

if $\varphi_s = df_s$, then

$$\lim_{\varepsilon\to 0}\left\{\varepsilon\!\!\int_\gamma \varphi_1\cdots\varphi_{s-1}df_s - \varepsilon\!\!\int_\gamma \varphi_1\cdots\varphi_{s-2}\big(f_s-f_s(\gamma(b))\varphi_{s-1}\big)\right\} = 0.$$

Proof: Assume now that $\gamma : [a;b]\to Z_0$ is a path over $\vec{v}$ (based at $\vec{w}$). Note that according to Proposition 4.5 we can decompose γ into piece such that for each of them the assertion of Proposition 4.8 holds: $\gamma = \gamma_1\star\cdots\star\gamma_N$. One has to show that if for a path $\alpha : [0;1]\to Z_0$ over $\vec{v}$ (based at $\vec{w}$) and for a path $\beta : [0;1]\to Z_0$ over $\vec{v}$ (based at $\vec{w}$) with $\alpha(1) = \beta(0)$ the assertion of Proposition 4.8 holds, that then the assertion is true for their composition $\alpha\star\beta$. This is a straightforward calculation similar to those in the proof of Proposition 4.5. □

4.2.1. *Formal Power Series Connections.* Let φ_i, φ_{ij}, φ_{ijk}, ...; i, j, $k = 1, 2, \dots, s$ be elements in A^1. Let $X_1,\dots,X_s$ be noncommutative indeterminates.

The formal power series

$$\omega = \sum\varphi_i X_i + \sum\varphi_{ij}X_iX_j + \sum\varphi_{ijk}X_iX_jX_k + \cdots$$

will be called a *formal power series connection for $A^\bullet$*. We refer to the number s as to the *length* of ω. The *curvature* κ of ω is defined by:

$$\kappa = d\omega + \omega\wedge\omega.$$

Associated to the given power series connection ω and a path $\gamma : [a;b] \to Z_0$ over $\vec{v}$ (based at $\vec{w}$) as well as $\varepsilon > 0$ small enough there is a formal power series with coefficients in $\mathbb{C}$, *the ε-transport of ω applied to γ*

$$ {}^{\varepsilon}\mathrm{T}(\gamma) = 1 + {}^{\varepsilon}\!\!\int\limits_{\gamma} \omega + {}^{\varepsilon}\!\!\int\limits_{\gamma} \omega\omega + {}^{\varepsilon}\!\!\int\limits_{\gamma} \omega\omega\omega + \cdots , $$

whose coefficients are given by

$$ {}^{\varepsilon}\mathrm{T}_i(\gamma) = {}^{\varepsilon}\!\!\int\limits_{\gamma} \varphi_i, \qquad {}^{\varepsilon}\mathrm{T}_{ij}(\gamma) = {}^{\varepsilon}\!\!\int\limits_{\gamma} \varphi_i\varphi_j + \varphi_{ij} $$

$$ {}^{\varepsilon}\mathrm{T}_{ijk}(\gamma) = {}^{\varepsilon}\!\!\int\limits_{\gamma} \varphi_i\varphi_j\varphi_k + \varphi_{ij}\varphi_k + \varphi_i\varphi_{jk} + \varphi_{ijk} $$

and in general

$$ {}^{\varepsilon}\mathrm{T}_{i_1,\dots,i_r}(\gamma) = \sum_{0<\alpha_1,\cdots<\alpha_{r-1}<r} {}^{\varepsilon}\!\!\int\limits_{\gamma} \varphi_{i_1\cdots i_{\alpha_1}} \cdots {}^{\varepsilon}\!\!\int\limits_{\gamma} \varphi_{i_{\alpha_{r-1}+1}\cdots i_r}. $$

The reason for us to make these definitions is the following Proposition.

Proposition 4.9. *Let* $\mathbf{p}$, $\mathbf{q}$, $\mathbf{r} \in Z_0 \setminus \bigcup_{[k<l]} \{p_{kl}\}$ *and let* α, $\beta : [0;1] \to Z_0$ *be paths over* $\vec{v}$ *(based at* $\vec{w}$*), where* α *goes from* $\mathbf{p}$ *to* $\mathbf{q}$ *and* β *from* $\mathbf{q}$ *to* $\mathbf{r}$*. Let* ω *be a power series connection. Then for* $\varepsilon > 0$ *small enough:*

$$ {}^{\varepsilon}\mathrm{T}(\alpha \star \beta) = {}^{\varepsilon}\mathrm{T}(\alpha) \cdot {}^{\varepsilon}\mathrm{T}(\beta). $$

Proof: The proof follows from the following rule

$$ {}^{\varepsilon}\!\!\int\limits_{\alpha\star\beta} \underbrace{\omega\cdots\omega}_{n} = \sum_{k=0}^{n} {}^{\varepsilon}\!\!\int\limits_{\alpha} \underbrace{\omega\cdots\omega}_{k} \;\; {}^{\varepsilon}\!\!\int\limits_{\beta} \underbrace{\omega\cdots\omega}_{n-k}, $$

which in turn can be seen to be a consequence of Proposition 3.6. □

Not difficult to prove is the following proposition when one uses Lemma 3.5.

Proposition 4.10. *Let* $\omega = \sum \varphi_i X_i + \sum \varphi_{ij} X_i X_j + \sum \varphi_{ijk} X_i X_j X_k + \cdots$ *be a formal power series connection for* $A^\bullet$*. Then* $\kappa = d\omega + \omega \wedge \omega = 0$ *if and only if the coefficients of the formal power series* $1 + \omega + \omega \otimes \omega + \omega \otimes \omega \otimes \omega + \cdots$ *are closed under* $(d_I + d_C)$*.* □

The next proposition allows us to use formal power series connections in order to prove things about Chen-closed elements in $\bigoplus_{r=0}^{s} \bigotimes^{r} A^1$.

Proposition 4.11. *Let* $I \in \bigoplus_{r=0}^{s} \bigotimes^{r} A^1$ *be Chen-closed. Then there exists a finite number of formal power series connections* $\omega_1, \dots, \omega_N$ *of length* s *with curvature 0 such that up to an element in* $\mathcal{R}_1^s(A^\bullet, a)$ I *is the coefficient of* $X_1 \cdots X_s$ *in:*

$$ \sum_{i=1}^{N} \left(1 + \omega_i + \omega_i \otimes \omega_i + \omega_i \otimes \omega_i \otimes \omega_i + \cdots \right). $$

Proof: Induction on s: $s = 1$ is trivial. So $s > 1$. By Lemma 3.3 and Lemma 3.5 we find that there are fomal power series connections (fpsc's) $\omega_1^{(s)}, \dots, \omega_K^{(s)}$ of length s such that the difference $I^{(s)}$ between I and the coefficient of $X_1 \cdots X_s$ in $\sum_{i=1}^{K} \left(1 + \omega_i^{(s)} + \omega_i^{(s)} \otimes \omega_i^{(s)} + \cdots \right)$ is of length $s-1$ and closed under $(d_I + d_C)$.

By induction hypothesis there exist fpsc's $\tilde{\omega}_1^{(s-1)}, \dots, \tilde{\omega}_L^{(s-1)}$ of length $s-1$ such that $I^{(s-1)}$ is the coefficient of $X_1 \cdots X_{s-1}$ in $\sum_{i=1}^L \left(1 + \tilde{\omega}_i^{(s-1)} + \tilde{\omega}_i^{(s-1)} \otimes \tilde{\omega}_i^{(s-1)} + \cdots\right)$. Define the fpsc's $\omega_i^{(s-1)}$ to be $\tilde{\omega}_i^{(s-1)}$ except that the indeterminate X_{s-1} is replaced by $X_{s-1}X_s$. The $\omega_i^{(s-1)}$ still have zero curvature! and I is the coefficient of $X_1 \cdots X_s$ in

$$\sum_{i=1}^{K} \left(1 + \omega_i^{(s)} + \omega_i^{(s)} \otimes \omega_i^{(s)} + \cdots\right) + \sum_{i=1}^{L} \left(1 + \omega_i^{(s-1)} + \omega_i^{(s-1)} \otimes \omega_i^{(s-1)} + \cdots\right).$$

□

Proposition 4.9 enables us to prove that the iterated integrals along paths over $\vec{v}$ converge not just locally but also globally.

Theorem 4.12. *Let ω be a formal power series connection for $A^\bullet$ with curvature $\kappa = 0$, let $\gamma : [0;1] \to Z_0$ be a path over $\vec{v}$ (based at $\vec{w}$) and let ${}^{\varepsilon}\mathrm{T}(\gamma)$ be the associated ε-transport of ω applied to γ. Then*

$$\mathrm{T}(\gamma) := \lim_{\varepsilon \to 0} {}^{\varepsilon}\mathrm{T}(\gamma)$$

converges and is called the transport of ω applied to γ. Moreover if $\alpha,\ \beta : [0;1] \to Z_0$ are paths over $\vec{v}$ (based at $\vec{w}$), which can be composed to a connected path over $\vec{v}$, then

$$\mathrm{T}(\alpha \star \beta) = \mathrm{T}(\alpha) \cdot \mathrm{T}(\beta).$$

Proof: Let $\gamma : [a;b] \to Z_0$ be a path over $\vec{v}$ (based at $\vec{w}$) that meets the set of double points $\{p_{kl}\}$ with parameter values $a = \tau_0 < \tau_1 < \cdots < \tau_m < \tau_{m+1} = b$. At the double point $p_0 \in D_0$, we have the embedding $x = p : D_0 \to \mathbb{C}$. Like before, let σ be the straight path $t \mapsto (1-t)$ from 1 to $0 \in D_0$. Decompose $\sigma \star \gamma \star \sigma^{-1} = \gamma_0 \star \cdots \star \gamma_{m+1}$ such that for each $\gamma_\kappa : [a_\kappa; b_\kappa] \to Z_0$, the interval $[a_\kappa; b_\kappa]$ contains only the parameter value τ_κ in its interior. There is an $\varepsilon_0 > 0$ with which we can decompose each of the paths γ_κ into $\gamma_\kappa = \alpha_\kappa \star \delta_\kappa \star \beta_\kappa$ such that α_κ resp. β_κ lie entirely in one component and each δ_κ is a map

$$\delta_\kappa : [\tau_\kappa - \varepsilon_0; \tau_\kappa + \varepsilon_0] \to U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$$

for some $[k < l]$.

By Theorem 4.6 we know that if ϑ is one of the paths $\alpha_\kappa,\ \delta_\kappa,\ \beta_\kappa$ then ${}^{\varepsilon}\mathrm{T}(\vartheta)$ converges. The proof can now be accomplished by applying Proposition 4.9. □

Hence we may conclude the following corollary.

Corollary 4.13. Let $I \in \bigoplus_{r=0}^{s} \bigotimes^r A^1$ be a Chen-closed element and let $\gamma : [0;1] \to Z_0$ be a path over $\vec{v}$ (based at $\vec{w}$). Then $\lim_{\varepsilon \to 0} {}^{\varepsilon}\!\!\int_\gamma I$ converges. If $\gamma : [0;1] \to Z_0$ is a path over $\vec{v}$ based at $\vec{w}$, then $\int_\gamma (\mathcal{R}_1^s(A^\bullet, a)) = 0$.

Proof: The assertions in the corollary are consequences of Proposition 4.11, Proposition 4.8 and Corollary 4.12. □

Theorem 4.14. *Let $I \in \bigoplus_{r=0}^{s} \bigotimes^r A^1$ be Chen-closed and let $\gamma : [0;1] \to Z_0$ be a path over $\vec{v}$ (based at $\vec{w}$). Then $\int_\gamma I$ does not depend on the choice of coordinates (x,y) around the double points.*

Remark 4.15. Theorem 4.6 may not be applied directly to prove Theorem 4.14 since a priori the definition of the dga $A_{kl}^\bullet$ depends on coordinates.

Proof of 4.14: By Lemma 4.11, Proposition 4.8 and Corollary 4.12 it is enough to prove the theorem for the following local situation. Assume without loss of generality that the path $\gamma : [0;1] \to Z_0$ over $\vec{v}$ lies entirely in a $U_{kl}^{k-} \cup \{p_{kl}\} \cup U_{kl}^{l-}$ for $[k < l]$ and meets the double point p_{kl} once with parameter value τ_0. Like earlier we distinguish two cases.

A path traversing the double point. We have

$$\int_\gamma I = \lim_{\varepsilon\to 0}\left\{\sum_J \varepsilon\!\!\!\int_\gamma \varphi_{j_1}\cdots\varphi_{j_r}\right\}$$

Recall that for each J there are polynomials $Q^J_{\alpha\beta} \in \mathbb{C}[u]$ such that $\varepsilon\!\!\!\int_\gamma \varphi_{j_1}\cdots\varphi_{j_r}$

$$= \sum_{0\le\alpha\le\beta\le r} \int_{\gamma\le\tau_0-\varepsilon} \tilde{\omega}_k^{(1)}\cdots\tilde{\omega}_k^{(\alpha)} Q^J_{\alpha\beta}\left(\log\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)\right) \int_{\gamma\ge\tau_0+\varepsilon} \tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)}.$$

Furthermore remark:

$$\left(\log\left(\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)\right)\right)^m = \left(\log\left(\frac{\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)}{\varepsilon^2}\right) + \log\varepsilon^2\right)^m$$

$$= \left(\log\varepsilon^2\right)^m + \left\{\log\left(\frac{\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)}{\varepsilon^2}\right)\right.$$
$$\left.\cdot\left(\sum_{k=1}^m \binom{m}{k}\log^{k-1}\left(\frac{\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)}{\varepsilon^2}\right)\left(\log\varepsilon^2\right)^{m-k}\right)\right\}.$$

The function $\log\left(\frac{\gamma_x(\tau_0-\varepsilon)\cdot\gamma_y(\tau_0+\varepsilon)}{\varepsilon^2}\right)$ is a differentiable function in ε on an intervall $(-\varepsilon_0;\varepsilon_0)$ for some ε_0, which vanishes in 0. Hence the vanishing of this function is stronger than any logarithmic growth (Use Fact 4.2 on page 47). This indicates that the following limit is zero:

$$\lim_{\varepsilon\to 0}\left\{\varepsilon\!\!\!\int_\gamma \varphi_{j_1}\cdots\varphi_{j_r} - \sum_{0\le\alpha\le\beta\le r}\int_{\gamma\le\tau_0-\varepsilon}\tilde{\omega}_k^{(1)}\cdots\tilde{\omega}_k^{(\alpha)}Q^J_{\alpha\beta}\left(\log\varepsilon^2\right)\int_{\gamma\ge\tau_0+\varepsilon}\tilde{\omega}_l^{(\beta+1)}\cdots\tilde{\omega}_l^{(s)}\right\}.$$

We find that $\int_\gamma I$ does not depend on coordinates.

A path colliding with a double point

Let $p := \gamma(0)$ and $q := \gamma(1)$. We know by 4.6 that there exists a closed and hence exact $\varphi_{pq} = df_{pq} \in A^1_{kl}$ such that

$$\int_\gamma I = \int_\gamma \varphi_{pq} = f_{pq}(q) - f_{pq}(p).$$

Let $\check{\gamma} : [a;b] \to U^{k-}_{kl}$ be a path (over $\vec{v}$), which is nearby homotopic relative to the end points with γ, but does not touch the double point. According to Proposition 4.3 we observe:

$$\int_\gamma \varphi_{pq} = f_{pq}(q) - f_{pq}(p) = \int_{\check{\gamma}} \varphi_{pq}$$

The integral on the right is independent of the choice of coordinates. $\square$

4.3. **Nearby Homotopy Functionals.** In this subsection we want to prove that iterated integrals of Chen-closed elements in $\bigoplus_{r=1}^{s}\bigotimes^{r}A^1$ are invariant under a homotopy over $\vec{v}$ based at $\vec{w}$. Hence, they are *nearby homotopy functionals*. There is no converse of this assertion since in general the iterated integrals only converge for Chen-closed elements in $\bigoplus_{r=1}^{s}\bigotimes^{r}A^1$.

Theorem 4.16. *Let $I \in \bigoplus_{r=0}^{s}\bigotimes^{r}A^1$ be Chen-closed. Then for two nearby homotopic paths $\gamma_0, \gamma_1 : [0;1] \to Z_0$ over $\vec{v}$ based at $\vec{w}$ holds:*

$$\int_{\gamma_0} I = \int_{\gamma_1} I.$$

Proof: By Lemma 4.11 and Proposition 4.8 it is enough to prove the corresponding theorem for a formal power series connection ω with curvature $\kappa = 0$. Let T be the associated transport. We want to prove: $\mathrm{T}(\gamma_0) = \mathrm{T}(\gamma_1)$

Let $H : [0;1] \times [0;1] \to Z_0$ be the nearby homotopy. That is, $H(\cdot, 0) = \gamma_0$ and $H(\cdot, \gamma_1) = \gamma_1$ and for any $\lambda \in [0;1]$ is $H(\cdot, \lambda)$ is a path over $\vec{v}$ based at $\vec{w}$. Consider the function:

$$\begin{array}{cccc} C: & [0;1] & \longrightarrow & \mathbb{C} \\ & \lambda & \longmapsto & \mathrm{T}(H(\cdot,\lambda)). \end{array}$$

We will show that C is constant by showing that it is locally constant.

Let $\lambda_0 \in [0;1]$. We show that C is locally constant at λ_0. Write $\eta := H(\cdot, \lambda_0)$ and denote by $0 = \tau_0 < \tau_1 < \cdots < \tau_N < \tau_{n+1} = 1$ the parameter values, where η meets the set of double points: η defines a map

$$\mu : \{0, \dots, N+1\} \longrightarrow \left\{ (k,l) \middle| \, [k < l] \right\},$$

by $\eta(\tau_i) = p_{\mu(i)}$ for $i = 0, \dots, N+1$.

For any $\varepsilon > 0$ we define $L_0^{\varepsilon} := [0;\varepsilon]$, $L_i^{\varepsilon} := [\tau_i - \varepsilon; \tau_i + \varepsilon]$ for $i = 1, \dots, N$ and $L_{N+1}^{\varepsilon} := [1-\varepsilon; 1]$. Choose a positive $\varepsilon_1 < \min\limits_i \frac{|\tau_{i+1} - \tau_i|}{2}$ (then all the $L_i^{\varepsilon_1}$ are mutually disjoint) such that: $\eta\left(L_i^{\varepsilon_1}\right) \subset U_{\mu(i)}^{k-} \cup \{p_{\mu(i)}\} \cup U_{\mu(i)}^{l-}$.

By the continuity of H and the fact that all $H(\cdot, \lambda)$ are paths over $\vec{v}$ we can find a $\delta > 0$ and an $\varepsilon_0 > 0$ (with $\varepsilon_0 < \varepsilon_1$) such that for all $i = 0, \dots, N+1$ holds:

$$H\left(L_i^{\varepsilon_0} \times [\lambda_0 - \delta; \lambda_0 + \delta]\right) \subset U_{\mu(i)}^{k-} \cup \{p_{\mu(i)}\} \cup U_{\mu(i)}^{l-}.$$

and the inverse image of the double points under $H\big|_{[0;1]\times[\lambda_0-\delta,\lambda_0+\delta]}$ lies entirely in the interior of the set $\bigcup_i L_i^{\varepsilon_0} \times [\lambda_0 - \delta, \lambda_0 + \delta]$. (If $\lambda_0 = 0$ resp. $\lambda_0 = 1$ then replace $[\lambda_0 - \delta, \lambda_0 + \delta]$ by $[0, \delta]$ resp. $[1 - \delta, 1]$.)

Fix a $\lambda \in [\lambda_0 - \delta, \lambda_0 + \delta]$. It is a consequence of Theorem 4.6 that T takes the same value on any two paths $H \circ \theta_i$ ($i = 1, 2$) over $\vec{v}$, where the θ_i are both paths with the same end points within $L_i^{\varepsilon_0} \times [\lambda_0 - \delta; \lambda_0 + \delta]$, whose end points are two of the four points: $\{(\tau_i \pm \varepsilon_0, \lambda_0),\ (\tau_i \pm \varepsilon_0, \lambda)\}$.

Moreover $\int I$ takes the same value on any two paths $H \circ \theta_i$ ($i = 1, 2$) over $\vec{v}$, where the θ_i are both paths with the same end points within $[\tau_i + \varepsilon_0, \tau_{i+1} - \varepsilon_0] \times [\lambda_0 - \delta; \lambda_0 + \delta]$, whose end points are two of the four points: $\{(\tau_i + \varepsilon_0, \lambda_0),\ (\tau_{i+1} - \varepsilon_0, \lambda_0),\ (\tau_i + \varepsilon_0, \lambda),\ (\tau_{i+1} - \varepsilon_0, \lambda)\}$, likewise by Theorem 4.6.

Now one easily constructs a finite sequence of paths $\alpha_0, \alpha_1, \dots, \alpha_M$ in $[0;1] \times [\lambda_0 - \delta; \lambda_0 + \delta]$ with the following properties:

- $H \circ \alpha_0 = H(\cdot, \lambda_0) = \eta$ and $H \circ \alpha_M = H(\cdot, \lambda)$,
- each $H \circ \alpha_i$ is a path over $\vec{v}$ based at $\vec{w}$,



FIGURE 8. Illustration of the domain of H, the path η and the construction of the sequence of paths $(\alpha_i)_{i=0,\dots,M}$.

- two successive paths α_i, α_{i+1} coincide everywhere except either on an interval $L_i^{\varepsilon_0} \times [\lambda_0 - \delta; \lambda_0 + \delta]$ or on an interval $[\tau_i + \varepsilon_0, \tau_{i+1} - \varepsilon_0] \times [\lambda_0 - \delta; \lambda_0 + \delta]$.

In other words we may write $\alpha_i = \zeta \star \theta \star \vartheta$ and $\alpha_{i+1} = \zeta \star \theta' \star \vartheta$. The operator T takes the same value on θ and θ'. Apply Corollary 4.12 to conclude $\mathrm{T}(H \circ \alpha_i) = \mathrm{T}(H \circ \alpha_{i+1})$ □

4.4. **Nilpotent Orbits.** Here we will apply the formal power series connections to prove Theorem 3.15. Since the operators M and N commute, we may prove the theorem by proving the following two equations separately:

(i) $\left(J_{\lambda\vec{v},\vec{w}}/J^{s+1}_{\lambda\vec{v},\vec{w}}\right)^*_{\mathbb{Z}} = \lambda^{-N}\left(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}}\right)^*_{\mathbb{Z}} \subset H^0\bar{B}_s(A^\bullet, a)$,

(ii) $\left(J_{\vec{v},\mu\vec{w}}/J^{s+1}_{\vec{v},\mu\vec{w}}\right)^*_{\mathbb{Z}} = \mu^{M}\left(J_{\vec{v},\vec{w}}/J^{s+1}_{\vec{v},\vec{w}}\right)^*_{\mathbb{Z}} \subset H^0\bar{B}_s(A^\bullet, a)$.

Proof of Theorem 3.15: We will prove (i) and leave (ii) for the reader.

Let $\lambda : [0;1] \to \mathbb{C} \setminus \mathbb{R}^{\leq 0}$ be a path with $\lambda(0) = 1$. Let λ_0 be a positive real number such that $\lambda_0 < |\lambda(\varsigma)|$ for all $\varsigma \in [0;1]$.

Like always, we fix coordinates $(x, y) : W_{kl} = U^k_{kl} \times U^l_{kl} \to \mathbb{C}^2$ with respect to $\vec{v}$, that means $\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y} \rangle = \vec{v}$. For an appropriate $\rho > 0$ let $\tilde{y}_\varsigma = \tilde{y} : U^l_{kl} \to \mathbb{C}$ be another coordinate that depends continuously on $\lambda(\varsigma)$ resp. ς and that satisfies:

$$\begin{aligned} \tilde{y}_\varsigma &= y && \text{for } |y| \geq \frac{\rho}{\lambda_0}, \\ \tilde{y}_\varsigma &= \frac{1}{\lambda(\varsigma)} y && \text{for } |y| \leq \rho \text{ and} \\ \tilde{y}_0 &= y && \text{for all } y. \end{aligned}$$

For example, take a smoothing of the map that is given in polar coordinates by:

$$\begin{pmatrix} \tilde{r} \\ \tilde{\varphi} \end{pmatrix} = \begin{pmatrix} (r-\rho)\frac{|\lambda|-\lambda_0}{\lambda-|\lambda|\lambda_0} + \frac{\rho}{|\lambda|} \\ \varphi + \arg(\lambda)\left(\frac{\lambda_0 r - \rho}{\rho - \rho\lambda_0}\right) \end{pmatrix} \quad \text{for } \rho \leq r \leq \frac{\rho}{\lambda_0}.$$

These coordinates $(x, \tilde{y}_\varsigma)$ are made such that holds: $\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial \tilde{y}_\varsigma} \rangle = \lambda(\varsigma)\vec{v}$.

The coordinate changes $\tilde{y}_\varsigma^{-1} \circ y$ at any U^l_{kl} can be put together to an isotopy

$$\phi_\varsigma : Z_0 \to Z_0,$$

which is defined to be $\tilde{y}_\varsigma^{-1} \circ y$ on any U^l_{kl} and the identity elsewhere. Note that $\phi_0 = id$ and $\phi_\varsigma(y) = \lambda(\varsigma)y$ for $y \in U^l_{kl}$ with $|y| \leq \rho$. Given a path $\gamma : [a; b] \to Z_0$ over $\vec{v}$ based at $\vec{w}$, we can define a homotopy H by $H(t, \varsigma) := \phi_\varsigma \circ \gamma(t)$. This $H(\cdot, \varsigma)$ is a path over $\lambda(\varsigma)\vec{v}$ based at $\vec{w}$.

Now take a Chen-closed $I = \sum_J a_J \; \varphi_{j_1} \otimes \cdots \otimes \varphi_{j_r} \in \bigoplus_{r=1}^s \bigotimes^r A^1$. We obtain by the definition of iterated integrals along paths over $\lambda(\varsigma)\vec{v}$ based at $\vec{w}$:

$$\int_{H(\cdot,\varsigma)} I = \int_\gamma \phi_\varsigma^* I. \tag{4.4}$$

Here the dga-automorphism $\phi_\varsigma^* : A^\bullet \to A^\bullet$ is defined by

$$(\phi_\varsigma\big|_{D_i})^* : E^\bullet(D_i \log P_i) \to E^\bullet(D_i \log P_i) \text{ and}$$
$$id : E^\bullet(\Delta^1) \otimes \Lambda^\bullet \to E^\bullet(\Delta^1) \otimes \Lambda^\bullet.$$

ϕ_ς^* induces an automorphism $\bigoplus_{r=1}^s \bigotimes^r \phi_\varsigma^*$ of $\bigoplus_{r=1}^s \bigotimes^r A^\bullet$, which we also denote by ϕ_ς^*. This is the ϕ_ς^* in (4.4).

We are done with the proof of the theorem if we can show for all paths $\gamma \in \pi_1(Z_{\vec{v}}, \vec{w})$:

$$\int_\gamma \phi_\varsigma^* I = \int_\gamma \lambda(\varsigma)^N I.$$

Also this statement will be proved by first looking at the local and then at the global situation.

Local Situation: Assume here that $\vartheta : [a; b] \to U^{k-}_{kl} \cup \{p_{kl}\} \cup U^{l-}_{kl}$ is a path over $\vec{v}$ that meets the double point p_{kl} once with parameter value τ_0, where it changes from D_k to D_l. (Since $\phi_\varsigma^* I$ and $\lambda(\varsigma)^N I$ are nearby homotopy functionals, we do not have to worry about paths that *collide with a double point.*) Suppose the endpoint $\vartheta(b)$ of ϑ be such that $|y(\vartheta(b))| > \frac{\rho}{\lambda_0}$.

Notice that the two coordinate systems (x, y) and $(x, \tilde{y}_\varsigma)$ around p_{kl} give two different *localizations in a sector at the double point* p_{kl}

$$A^\bullet \to A^\bullet_{kl} \qquad \text{and} \qquad A^\bullet \to \tilde{A}^\bullet_{kl}$$

respectively. Denote the images of I under these natural maps by I_{kl} resp. $\tilde{I}_{kl}$. Observe moreover that ϕ_ς induces

$$\phi_\varsigma : U^{k-}_{kl} = \{y \not\in \mathbb{R}^{\leq 0}\} \xrightarrow{\approx} \tilde{U}^{k-}_{kl} = \{\tilde{y}_\varsigma \not\in \mathbb{R}^{\leq 0}\}.$$

Then we can extend the dga-automorphisms ϕ_ς^* and $\lambda(\varsigma)^N$ of $A^\bullet$ to dga-isomorphisms with the same names, ϕ_ς^* and $\lambda(\varsigma)^N$, from $\tilde{A}^\bullet_{kl}$ to $A^\bullet_{kl}$ like indicated in the following two commutative diagrams.

$$\begin{tikzcd}
A^\bullet \arrow[r] \arrow[d, "\phi_\varsigma^*"'] & \tilde{A}^\bullet_{kl} \arrow[d, "\phi_\varsigma^*"] & \log \tilde{y} \arrow[d] & w \arrow[d] \\
A^\bullet \arrow[r] & A^\bullet_{kl} & \log \tilde{y} \circ \phi_\varsigma(y) & w,
\end{tikzcd}$$

I. e. ϕ_ς on $\tilde{A}^\bullet_{kl}$ is the pull-back with ϕ^*_ς. Extend $\lambda(\varsigma)^N$ by

$$\begin{tikzcd}
A^\bullet \arrow[r] \arrow[d, "\lambda(\varsigma)^N"'] & \tilde{A}^\bullet_{kl} \arrow[d, "\lambda(\varsigma)^N"] & \log \tilde{y} \arrow[d] & w \arrow[d] \\
A^\bullet \arrow[r] & A^\bullet_{kl} & \log \tilde{y}(y) & (w - \log \lambda).
\end{tikzcd}$$

We know by Theorem 4.6 that for $\tilde{I}_{kl}$ and the points $\vartheta(b)$, $\vartheta(a)$ there is a function $\tilde{f}_{kl} \in \tilde{A}^0_{kl}$ such that holds:

$$\int_\vartheta I = \int_\vartheta \tilde{I}_{kl} = \tilde{f}_{kl}\big(\tilde{y} \circ \vartheta(b)\big) - \tilde{f}_{kl}\big(\tilde{y} \circ \vartheta(a)\big).$$

Since both, ϕ^*_ς and $\lambda(\varsigma)^N$, are isomorphisms of dga's, which preserve the values of functions on $\vartheta(b)$ and $\vartheta(a)$, the pull-back $\phi^*_\varsigma \tilde{f}_{kl}$ is such a function for $\phi^*_\varsigma \tilde{I}_{kl}$ and $\lambda(\varsigma)^N \tilde{f}_{kl}$ is such a function for $\lambda(\varsigma)^N \tilde{I}_{kl}$. That is to say:

$$\int_\vartheta \phi^*_\varsigma I = \int_\vartheta \phi^*_\varsigma \tilde{I}_{kl} = \left(\phi^*_\varsigma \tilde{f}_{kl}\right) (\tilde{y} \circ \vartheta(b)) - \left(\phi^*_\varsigma \tilde{f}_{kl}\right) (\tilde{y} \circ \vartheta(a))$$

and

$$\int_\vartheta \lambda(\varsigma)^N I = \int_\vartheta \lambda(\varsigma)^N \tilde{I}_{kl} = \left(\lambda(\varsigma)^N \tilde{f}_{kl}\right) (\tilde{y} \circ \vartheta(b)) - \left(\lambda(\varsigma)^N \tilde{f}_{kl}\right) (\tilde{y} \circ \vartheta(a)).$$

If $\tilde{f}_{kl}$ is given by

$$\tilde{f}_{kl} = \sum_{\nu=0}^{m} g_{k,\nu}(x) \log^\nu x + \sum_{\nu=0}^{m} g_{l,\nu}(\tilde{y}) \log^\nu \tilde{y} + P(\xi, v, w)$$

then

$$\phi^*_\varsigma \tilde{f}_{kl} = \sum_{\nu=0}^{m} g_{k,\nu}(x) \log^\nu x + \sum_{\nu=0}^{m} g_{l,\nu}(y) \log^\nu y + P(\xi, v, w)$$

and

$$\lambda(\varsigma)^N \tilde{f}_{kl} = \sum_{\nu=0}^{m} g_{k,\nu}(x) \log^\nu x + \sum_{\nu=0}^{m} g_{l,\nu}(\tilde{y}(y)) \log^\nu \big(\tilde{y}(y)\big) + P(\xi, v, w - \log \lambda).$$

Because of $\tilde{y}\big(y \circ \vartheta(b)\big) = y \circ \vartheta(b)$ as $|y \circ \vartheta(b)| > \frac{\rho}{\lambda_0}$, we finally find:

$$\int_\vartheta \phi^*_\varsigma I = \int_\vartheta \lambda(\varsigma)^N I.$$

Global Situation: Let $\gamma : [a; b] \to Z_0$ be a path over $\vec{v}$ based at $\vec{w}$. At the double point $p_0 \in D_0$, we have the embedding $x = p : D_0 \to \mathbb{C}$. Like before, let σ be the straight path $t \mapsto (1-t)$ from 1 to $0 \in D_0$. Decompose like in the proof of Theorem 4.12 the path $\sigma \star \gamma \star \sigma^{-1} = \gamma_0 \star \cdots \star \gamma_{m+1}$ such that for a Chen-closed I each of the paths γ_ν satisfies: $\int_{\gamma_\nu} \phi^*_\varsigma I = \int_{\gamma_\nu} \lambda(\varsigma)^N I$.

Let ω be a formal power series connection with curvature zero. Then we can apply ϕ^*_ς and $\lambda(\varsigma)^N$ to ω by applying them to each coefficient. We know that the transport associated to $\phi^*_\varsigma \omega$ and the transport associated to $\lambda(\varsigma)^N \omega$ take the same value on all paths γ_ν. Apply Theorem 4.12 to prove the theorem. $\square$

5. Plane Curve Singularities

In this last section, we finally want to show, how the setting of 1.1 arises in the case of plane curve singularities. Throughout the whole section let

$$f = f_0 \cdots f_{r-1} : \big(\mathbb{C}^2, 0\big) \to (\mathbb{C}, 0)$$

be a plane curve singularity with r branches.

First we will construct a map $h : (Z, D^+) \to (\Delta, 0)$ with the properties described earlier in 1.1 and which is related to f in a natural way. We obtain this map $h : (Z, D^+) \to (\Delta, 0)$ by a construction that is described in [Ste77]. We refer to this process as *semistable reduction.*

Then we construct orbits of tangent vectors in $T_0\Delta$ and in $T_{p_0}D_0$ under the action of certain roots of unity. All these objects are defined in such a way that any right-equivalence between two plane curve singularities transforms these data into each other. That is, what the word *natural* refers to in this context.

Finally, we will define *the mixed Hodge structure on the nearby fundamental group of a plane curve singularity* and will finish with an example.

5.1. **Semistable Reduction.** Consider the plane curve singularity $f = f_0 \cdots f_{r-1}$ with r branches and let $\pi : (\tilde{X}, E^+) \to (\mathbb{C}^2, 0)$ be its embedded (minimal) resolution. Let E denote the divisor $(f \circ \pi)^{-1}(0)$ with components $E = \bigcup_{i \geq 0} E_i$, where E_i has multiplicity e_i. Let $E_0 \cup \cdots \cup E_{r-1}$ be *the strict preimage* of $f^{-1}(0)$ in $\tilde{X}$; that is the closure of $(f \circ \pi)^{-1}(0) \setminus \pi^{-1}(0)$ in $\tilde{X}$. Assume that the component E_i corresponds to the factor f_i in f. Define E^+, *the exceptional divisor*, to be $\bigcup_{i > r-1} E_i$. The E_i's for $i > r-1$ are all rational curves. Call $q_0, \ldots, q_{r-1}$ the intersection points $E_i \cap E^+$ for $i = 0, \ldots, r-1$.

Let $d = \operatorname{lcm}_i e_i$ and let $\theta : \Delta \to \mathbb{C}$ be the map $t \mapsto t^d$. Now consider the normalization of the fibered product of $f \circ \pi$ and θ:

$$Z' := \widetilde{\tilde{X} \times_{\mathbb{C}} \Delta}.$$

There are projections $h' : Z' \to \Delta'$ and $P' : Z' \to \tilde{X}$. Denote the closure of $(f \circ \pi \circ P')^{-1}(0) \setminus (\pi \circ P')^{-1}(0)$ in Z' by $D_0' \cup \cdots \cup D_{r-1}'$. It is not difficult to see that due to the fact that $e_0 = \cdots = e_{r-1} = 1$ the projection P' maps $D_0' \cup \cdots \cup D_{r-1}'$ to $E_0 \cup \cdots \cup E_{r-1}$. Assume the notation to be such that D_i' is mapped to E_i.

Proposition 5.1. *Z' has possibly cyclic quotient singularities. More precisely, for each singular point $p \in Z'$ there exists a natural number $n_p > 0$ and an isomorphism*

$$\varphi_p : \mathcal{O}_{Z',p} \xrightarrow{\cong} \frac{\mathbb{C}\{\alpha, \beta, \eta\}}{(\alpha\beta - \eta^{n_p})}$$

that maps h' to η.

For a proof of this proposition we refer to [Ste77] or [Kae97].

Now we resolve these singularities of Z' according to a well-known procedure (cf. for instance [BK91]). For a singularity of the type: $\alpha\beta - \eta^n$ we replace the singular point by a chain of $(n-1)$ copies of $\mathbb{P}^1$ in the following way.

Consider n copies of $\mathbb{C}^2$ with coordinates (u_i, v_i) for $i = 1, \ldots, n$. These spaces are glued together by the following glueing maps:

$$\begin{array}{rccc} \varphi_i : & \mathbb{C}^2 \setminus \{v_i = 0\} & \to & \mathbb{C}^2 \setminus \{u_{i+1} = 0\} \\ & (u_i, v_i) & \mapsto & \big(\frac{1}{v_i},\ u_i v_i^2\big) = (u_{i+1}, v_{i+1})\,. \end{array}$$

Call the result M and observe that M is smooth. Now the resolution of the variety $\{\alpha\beta - \eta^n = 0\}$ is given by the map $M \to \{\alpha\beta - \eta^n = 0\}$, that is defined on the coordinates (u_i, v_i) by $(u_i, v_i) \mapsto \big(u_i^i v_i^{i-1};\ u_i^{n-i} v_i^{n+1-i};\ u_i v_i\big) = (\alpha, \beta, \eta)$. Notice that this map is well-defined. In this way we can resolve all the singularities of Z' and find a map $h : (Z, D^+) \to (\Delta, 0)$ with the desired properties. We have a commutative diagram of space germs

$$\begin{tikzcd}
(Z, D^+) \arrow[r, "P"] \arrow[d, "h"'] & (\tilde{X}, {D'}^+) \arrow[d, "f\circ\pi"] \\
(\Delta, 0) \arrow[r] & (\mathbb{C}, 0) \\
t \arrow[r, mapsto] & t^d \, ,
\end{tikzcd} \tag{5.1}$$

such that P induces an isomorphism over Δ^* between the spaces $Z \setminus h^{-1}(0)$ and $(\mathbb{C}^2 \setminus f^{-1}(0)) \times_{\mathbb{C}^*} \Delta^*$. With the notation of 1.1 we are ready to prove:

Proposition 5.2. *Let f, $\tilde{f} : \big(\mathbb{C}^2, 0\big) \to (\mathbb{C}, 0)$ be two right-equivalent plane curve singularities, i. e. there is a biholomorphism $\phi : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ such that $\tilde{f} \circ \phi = f$. Let then $h : (Z, D^+) \to (\Delta, 0)$ and $\tilde{h} : (\tilde{Z}, \tilde{D}^+) \to (\Delta, 0)$ be the maps constructed by the procdure above for f and $\tilde{f}$ respectively. Then ϕ lifts to a biholomorphism $\Phi : (Z, D^+) \to (\tilde{Z}, \tilde{D}^+)$ such that holds $\tilde{h} \circ \Phi = h$.*

Proof: The assertion follows from the fact that both copies of $(\mathbb{C}^2, 0)$ have undergone the same blowing-up procedures, which of course do not depend on coordinates. □

5.2. **Canonical Tangent Vectors.** In this subsection we first recall some well-known facts about the map $t \mapsto t^d$. These observations will be used to construct an orbit under the action of the d-th roots of unity on $(T_0\Delta)^*$.

If we distinguish one branch f_0 of the singularity f then a careful study of the Puiseux expansion of f_0 will show subsequently that there is a natural orbit under the action of certain roots of unity on $(T_{p_0} D_0)^*$, which we call *the monstrance* of (f, f_0).

5.2.1. *Pulling Back Tangent Vectors.* Let Δ and Δ' be two disks in $\mathbb{C}$ around 0. Always, when we are given a holomorphic map $\varphi : \Delta \to \Delta'$ of multiplicity m, then the differential of φ in 0 is zero, when $m > 1$. However, a tangent vector $\vec{v}\,' \in (T_0\Delta')^*$ defines in a natural way m tangent vectors $\zeta_m^\nu \vec{v} \in (T_0\Delta)^*$ for $\nu = 0, \dots, m-1$.

Let us refer to the set $\big\{\zeta_m^\nu \vec{v} \in (T_0\Delta)^* \,\big|\, \nu = 0, \dots, m-1\big\}$ as to *the inverse star of $\vec{v}\,'$ under φ*. We will give two descriptions of this phenomenon: a concrete one and an abstract one.

The concrete description is as follows. Choose a coordinate z' on $(\Delta', 0)$ such that $\vec{v}\,' = \frac{\partial}{\partial z'}$. Then there is up to a choice of an m-th root of unity only one coordinate z on Δ such that: $z' \circ \varphi = z^m$. Define the m tangent vectors by $\zeta_m^\nu \frac{\partial}{\partial z} \in (T_0\Delta)^*$, where $\nu = 0, \dots, m-1$. Notice that this definition does not depend on the choice of z'.

The abstract description is the following. Let $\mathfrak{m}$ resp. $\mathfrak{m}'$ be the maximal ideal in $\mathcal{O}_{\Delta,0}$ resp. $\mathcal{O}_{\Delta',0}$ and let $\mathfrak{m}/\mathfrak{m}^2$ resp. $\mathfrak{m}'/{\mathfrak{m}'}^2$ be the corresponding Zariski cotangent spaces. Observe that by the duality

$$\begin{array}{ccc} T_0\Delta \otimes \big(\mathfrak{m}/\mathfrak{m}^2\big) & \to & \mathbb{C} \\ (X \otimes g) & \mapsto & X(g)(0) \end{array}$$

a choice of a tangent vector in $T_0\Delta$ corresponds to the choice of an element in $\mathfrak{m}/\mathfrak{m}^2$. Now φ induces a map

$$\begin{array}{cccc} \varphi^* : & \mathfrak{m}'/\mathfrak{m}'^2 & \to & \mathfrak{m}^m/\mathfrak{m}^{m+1} \\ & g' & \mapsto & g' \circ \varphi. \end{array}$$

On the other hand we have the composition of natural maps:

$$\begin{array}{ccccc} \mathfrak{m}/\mathfrak{m}^2 & \to & \big(\mathfrak{m}/\mathfrak{m}^2\big)^{\otimes m} & \to & \mathfrak{m}^m/\mathfrak{m}^{m+1} \\ g & \mapsto & g \otimes \cdots \otimes g & & \\ & & a_1 \otimes \cdots \otimes a_m & \mapsto & a_1 \cdots a_m. \end{array}$$

Each non-zero element in $\mathfrak{m}^m/\mathfrak{m}^{m+1}$ has m inverse images in $\mathfrak{m}/\mathfrak{m}^2$ under this composition of maps. This shows that a tanget vector $\vec{v}\,' \in (T_0\Delta')^*$ determines m tangent vectors in $(T_0\Delta)^*$ in a natural way.

For plane curve singularities $f : \big(\mathbb{C}^2, 0\big) \to (\mathbb{C}, 0)$, the tangent vector $\frac{\partial}{\partial t} \in (T_0\mathbb{C})^*$ is preserved under right-equivalences. Denote by

$$\mathcal{S}(\frac{\partial}{\partial t}) = \{\zeta_d^\nu \vec{v} \in (T_0\Delta)^* | \nu = 0, \dots, d-1\}$$

the inverse star of $\frac{\partial}{\partial t}$ under the map $\theta : t \mapsto t^d$.

5.2.2. *The Monstrance of* (f, f_0). Let f_0 be a distinguished branch of f with Puiseux pairs (m_1, n_1), ... , (m_g, n_g). With the definitions $m := m_1 \cdots m_g$ and $k := n_1 m_2 \cdots m_g$, we will define a natural orbit of $m \cdot k$ tangent vectors in $(T_{p_0}E_0)^*$ under the induced action of the group $\mu_{m\cdot k}$ of mk-th roots of unity. We will refer to this orbit as to *the monstrance* of (f, f_0) and $\frac{\partial}{\partial t}$. We denote it by $\mathcal{M}(f_0; \frac{\partial}{\partial t})$. Its elements will serve as *base vectors* like the $\vec{w}$ in the discussion in the previous sections.

The idea for the construction is the following. Note that the map P in (5.1) maps E_0 biholomorphically to D_0. We know that if we have a parametrization $\tau : (\mathbb{C}, 0) \to (E_0, q_0)$ of the strict preimage, then this defines a tangent vector in $T_{q_0}E_0$ respectively $T_{p_0}D_0$, namely: $\frac{\partial}{\partial \tau}$. The Puiseux expansion is a parametrization $p : (\mathbb{C}, 0) \to \big(f_0^{-1}(0), 0\big)$ with the property that it can be lifted to a smooth parametrization $\tilde{p} : (\mathbb{C}, 0) \to (E_0, q_0)$. In this way, a Puiseux expansion yields a tangent vector in $T_{q_0}E_0$. Now for the irreducible plane curve singularity f_0, we can always find coordinates (x, y) of $\mathbb{C}^2$ such that

$$f_0(x, y) = 1 \cdot x^m + \cdots + 1 \cdot y^k + \cdots ,$$

where x^m and y^k are the smallest pure powers of x and y respectively. Up to first order, there are $m \cdot k$ such coordinate systems. This requirement on the coordinates with respect to f_0 determines the Puiseux expansions up to first order. Up to first order, we get $m\cdot k$ canonical parametrizations of (E_0, q_0) and therefore $m\cdot k$ canonical tangent vectors.

But let us give a geometric meaning to this construction. Since for the moment we do not want to fix coordinates on $\mathbb{C}^2$, we let $(X, x_0) := (\mathbb{C}^2, 0)$.

Definition 5.3. Let $\iota : (L, x_0) \hookrightarrow (X, x_0)$ be a 1-dimensional smooth subspace of (X, x_0). We will call a map $\mathbf{P} : (X, x_0) \to (L, x_0)$ *a projection* of (X, x_0) onto (L, x_0), if the induced map $T_{x_0}L \xrightarrow{D\iota} T_{x_0}X \xrightarrow{D\mathbf{P}} T_{x_0}L$ is the identity.

Now let $\iota : (R, x_0) \hookrightarrow (X, x_0)$ be the tangent to $(f_0^{-1}(0), x_0) \subset (X, x_0)$. We formulate the following theorem.

Theorem 5.4. *Let* $f = f_0 \cdots f_{r-1} : (X, x_0) \to (\mathbb{C}, 0)$ *be a plane curve singularity with one distinguished branch* f_0. *Let* $(m_1, n_1), \ldots, (m_g, n_g)$ *be the Puiseux pairs of* f_0. *Set* $m := m_1 \cdots m_g$ *and* $k := n_1 m_2 \cdots m_g$.

Let $\iota : (R, x_0) \hookrightarrow (X, x_0)$ *be the tangent and* $\pi : (\tilde{X}, E^+) \to (X, x_0)$ *be the embedded resolution of* f. *Then for any projection* $\mathbf{P} : (X, x_0) \to (R, x_0)$, *the composition* ϕ *of maps*

$$(E_0, q_0) \xrightarrow{\pi_{|(E_0,q_0)}} (X, x_0) \xrightarrow{\mathbf{P}} (R, x_0) \xrightarrow{f_{0|(R,x_0)}} (\Delta, 0)$$

has multiplicity $m \cdot k$ *and the inverse star of the tangent vector* $\frac{\partial}{\partial t}$ *under* ϕ *does not depend on the choice of the projection* $\mathbf{P}$.

Proof: Choose coordinates (x, y) on (X, x_0) such that the following two conditions are satisfied:

- The tangent line (R, x_0) of $(f_0^{-1}(0), x_0)$ is given by $x = 0$.
- $f_0(x, y) = 1 \cdot x^m + \cdots + 1 \cdot y^k + \cdots$, where x^m and y^k are the smallest pure powers of x and y respectively.

Then the Puiseux expansion of f_0 with respect to (x, y) has the following form, where $m < k$ (f_0 is irreducible):

$$\begin{aligned} x(\tau) &= a_k \tau^k + a_{k+1} \tau^{k+1} + \cdots \\ y(\tau) &= \tau^m \end{aligned}$$

and $a_k^m = -1$. Moreover we can find f_0 back by:

$$f(x, y) = \prod_{\nu=0}^{m-1} \left(x - x(\zeta_m^\nu y^{\frac{1}{m}}) \right).$$

τ can be used as a coordinate on the strict transform (E, q_0). We may take $\tilde{y} := y_{\big|(R,x_0)}$ as coordinate on the tangent line (R, x_0).

Any projection $\mathbf{P} : (X, x_0) \to (R, x_0)$ is a priori given by $\tilde{y} \circ \mathbf{P}(x, y) = y(\mathbf{P}(x, y)) = Ax + By + \mathbf{h}(x, y)$ and $\mathbf{h}(x, y) \in \mathfrak{m}^2_{(X,x_0)}$. The condition $D\mathbf{P}_{\big|T_{x_0}R} = id$ means here: $B = 1$. Hence, we find for a unit $u(\tau) \in \mathbb{C}\{\tau\} \setminus \mathfrak{m}$ with $u(0) = \zeta_m^\nu$:

$$\begin{aligned} \tilde{y} \circ \mathbf{P}\left(x(\tau), y(\tau)\right) &= \tau^m + \tau^{m+1}\left(\cdots\right) \\ &= \tau^m \cdot u(\tau)^m. \end{aligned}$$

Therefore

$$\begin{aligned} \phi(\tau) &= f_0\left(0, \tilde{y} \circ \mathbf{P}\left(x(\tau),\ y(\tau)\right)\right) \\ &= (-1)^m \prod_{\nu=0}^{m-1} x\left(\zeta_m^\nu \tau \cdot u(\tau)\right) \\ &= (-1)^m \prod_{\nu=0}^{m-1} \left(a_k (\zeta_m^\nu \tau \cdot u)^k + a_{k+1} (\zeta_m^\nu \tau \cdot u)^{k+1} + \cdots \right) \\ &= (-1)^{m+k} a_k^m \tau^{m \cdot k} + \tau^{m \cdot k + 1}\big(\cdots\big). \end{aligned}$$

This shows that $\phi(\tau)$ has multiplicity $m \cdot k$ and the term of lowest order does not depend on the choice of $\mathbf{P}$. $\square$

Definition 5.5. Define *the monstrance of* (f, f_0) to be the inverse star of $\frac{\partial}{\partial t}$ under a map ϕ like in Theorem 5.4 and denote it by

$$\mathcal{M}(f_0; \frac{\partial}{\partial t}) = \{\zeta_{mk}^{\nu} \vec{w} \in (T_{q_0} E_0)^* |\ \nu = 0, \dots , mk - 1\}.$$

Remark 5.6. $m \cdot k$ does not divide d in general. For instance in the example of Section 5.4 is $m = 4$, $k = 6$ and $d = 156$. However, if $f = f_0$ and f_0 has only one Puiseux pair $(m, k) = (m_1, n_1)$, then $m \cdot k$ divides d as one easily sees by blowing up twice.

5.3. **The Invariant.** Here finally we are going to associate to a plane curve singularity an invariant of its right-equivalence class. In fact it is an invariant of the following seemingly coarser equivalence relation.

Definition 5.7. We say that two hypersurface singularities

$$f,\, g : (\mathbb{C}^{n+1}, 0) \to (\mathbb{C}, 0)$$

are *right-equivalent to first order* if there are isomorphisms $\phi : (\mathbb{C}^{n+1}, 0) \to (\mathbb{C}^{n+1}, 0)$ and $\varphi : (\mathbb{C}, 0) \to (\mathbb{C}, 0)$ such that $g \circ \phi = \varphi \circ f$ and $\varphi_* : T_0\mathbb{C} \to T_0\mathbb{C}$ is the identity.

The following theorem[13] might be interesting in this context.

Theorem 5.8. *Two plane curve singularities are right-equivalent if they are right-equivalent to first order.*

Proof: Let $f : (\mathbb{C}^2, 0) \to (\mathbb{C}, 0)$ be a plane curve singularity and let $h(t) = t + a_2t^2 + a_3t^3 + \cdots$ be in $\mathbb{C}\{t\}$. We have to show that f and $h \circ f$ are right-equivalent. Let $\mathfrak{m}$ be the maximal ideal in $\mathbb{C}\{x, y\}$. According to a theorem of Briançon-Skoda [BS74] (see also [LT81]) $f^2 \in \mathfrak{m}\mathbb{C}\{x, y\}$. Using this result one can show by a standard construction (see for instance [Lam75] p. 189, V§4, Prop. 4) that all f_t in $f_t(x, y) := f(x, y) + t\left(a_2f^2(x, y) + a_3f^3(x, y) + \cdots\right)$ are mutually right-equivalent. $\square$

Definition 5.9. Let $f = f_0 \cdots f_{r-1} : (\mathbb{C}^2, 0) \to (\mathbb{C}, 0)$ be a plane curve singularity. For all $s \geq 1$ we define

$$\mathfrak{P}^s(f, \tfrac{\partial}{\partial t}) := \bigoplus_{i=0}^{r-1} \left\{ \bigoplus_{\vec{v} \in \mathcal{S}(\frac{\partial}{\partial t})} \quad \bigoplus_{\vec{w}_i \in \mathcal{M}(f_i; \frac{\partial}{\partial t})} \left(J_{\vec{v}, \vec{w}_i} / J_{\vec{v}, \vec{w}_i}^{s+1} \right)^* \right\} \tag{5.2}$$

and we call the direct limit

$$\mathfrak{P}(f, \tfrac{\partial}{\partial t}) := \varinjlim_{s}\ \mathfrak{P}^s(f, \tfrac{\partial}{\partial t})$$

the mixed Hodge structure on the nearby fundamental group of the plane curve singularity f.

Note that μ_d acts on $\mathfrak{P}(f, \frac{\partial}{\partial t})$. More precisely, $\zeta_d^\nu = e^{\frac{2\pi i}{d}\nu} \in \mu_d$ acts as the map:

$$(\zeta_d^\nu)^{-N} = e^{2\pi i \nu \frac{N}{d}} : \left(J_{\vec{v}, \vec{w}_i} / J_{\vec{v}, \vec{w}_i}^{s+1} \right)_{\mathbb{Z}}^* \to \left(J_{\zeta_d^\nu \vec{v}, \vec{w}_i} / J_{\zeta_d^\nu \vec{v}, \vec{w}_i}^{s+1} \right)_{\mathbb{Z}}^* . \tag{5.3}$$

Let $(m_1^{(i)}, n_1^{(i)}), \dots , (m_{g_i}^{(i)}, n_{g_i}^{(i)})$ be the Puiseux pairs of the branch f_i and define $m^{(i)} := m_1^{(i)} \cdots m_{g_i}^{(i)}$ and $k^{(i)} := n_1^{(i)} m_2^{(i)} \cdots m_{g_i}^{(i)}$. Then each of the groups $\mu_{m^{(i)}k^{(i)}}$

---

[13] This theorem was pointed out to us by Claus Hertling.

acts on $\mathfrak{P}(f, \frac{\partial}{\partial t})$. Let $M^{(i)}$ be the map M of 3.15 associated to the distinguished branch f_i. Then $\zeta^{\nu}_{m^{(i)}k^{(i)}}$ induces the map

$$\left(\zeta^{\nu}_{m^{(i)}k^{(i)}}\right)^{M^{(i)}} : \left(J_{\vec{v},\vec{w}_i}/J^{s+1}_{\vec{v},\vec{w}_i}\right)^*_{\mathbb{Z}} \to \left(J_{\zeta_d^{\nu}\vec{v},\zeta^{\nu}_{m^{(i)}k^{(i)}}\vec{w}_i}/J^{s+1}_{\zeta_d^{\nu}\vec{v},\zeta^{\nu}_{m^{(i)}k^{(i)}}\vec{w}_i}\right)^*_{\mathbb{Z}}. \tag{5.4}$$

By Theorem 3.15 all the maps in (5.3) and (5.4) are isomorphisms of MHSs over $\mathbb{Z}$. Let $\mathcal{T}$ be the monodromy of the local system

$$\mathfrak{P}(f, \tfrac{\partial}{\partial t})_{\mathbb{Z}} = \bigoplus_{i=0}^{r-1} \left\{ \bigoplus_{\vec{v}\in\mathcal{S}(\frac{\partial}{\partial t})} \quad \bigoplus_{\vec{w}_i\in\mathcal{M}(f_i;\frac{\partial}{\partial t})} \left(J_{\vec{v},\vec{w}_i}/J^{s+1}_{\vec{v},\vec{w}_i}\right)^*_{\mathbb{Z}} \right\}.$$

Define $L := (\frac{1}{d}N - \sum_{i=0}^{r-1}\frac{1}{m^{(i)}k^{(i)}}M^{(i)}) : H^0\bar{B}(A^\bullet, a) \to H^0\bar{B}(A^\bullet, a)$ and moreover

$$\mathcal{L} := \bigoplus_{i,\vec{v},\vec{w}_i} L : \bigoplus_{i,\vec{v},\vec{w}_i} H^0\bar{B}(A^\bullet, a) \to \bigoplus_{i,\zeta_d\vec{v},\zeta_{m^{(i)}k^{(i)}}\vec{w}_i} H^0\bar{B}(A^\bullet, a).$$

The map $\mathcal{L}$ sends the summand with index $(i, \vec{v}, \vec{w}_i)$ to the summand corresponding to $(i, \zeta_d\vec{v}, \zeta_{m^{(i)}k^{(i)}}\vec{w}_i)$. By virtue of Theorem 3.15 we find for $\vartheta \in \mathbb{C}^*$:

$$\mathcal{T} = e^{-2\pi i\mathcal{L}} \qquad \text{and} \qquad \mathfrak{P}^s(f, \vartheta\tfrac{\partial}{\partial t})_{\mathbb{Z}} = \vartheta^{-\mathcal{L}}\,\mathfrak{P}^s(f, \tfrac{\partial}{\partial t})_{\mathbb{Z}},$$

from where we may conclude that the family of MHSs

$$\{\,\mathfrak{P}(f, \tfrac{\partial}{\partial t})\}_{\frac{\partial}{\partial t}\in T_0\mathbb{C}}$$

is a nilpotent orbit of MHS. Note that on the components of the direct sum (5.2) the monodromy $\mathcal{T}$ is given by

$$e^{-2\pi iL} : \left(J_{\vec{v},\vec{w}_i}/J^{s+1}_{\vec{v},\vec{w}_i}\right)^*_{\mathbb{Z}} \to \left(J_{\zeta_d\vec{v},\zeta_{m^{(i)}k^{(i)}}\vec{w}_i}/J^{s+1}_{\zeta_d\vec{v},\zeta_{m^{(i)}k^{(i)}}\vec{w}_i}\right)^*_{\mathbb{Z}}.$$

We consider both lattices as sub-lattices of $H^0\bar{B}(A^\bullet, a)$.

5.4. **An Example.** The only purpose of the example we give in this subsection is to indicate that *the mixed Hodge structure on the nearby fundamental group of a plane curve singularity* detects moduli that are hidden for the MHS on the vanishing cohomology. We focus especially on effects coming from the monodromy on $\vec{J}/\vec{J}^4$. But in general, we also expect interesting information from higher periods – even on $\vec{J}/\vec{J}^3$. A systematic investigation of the invariant on classes of singularities still remains to be done.

Here we investigate the example (0.1) that was mentioned in the introduction:

$$f_\lambda(x, y) = \left(y^2 - x^3\right)^2 - \left(4\lambda x^5 y + \lambda^2 x^7\right), \quad \lambda \neq 0.$$

All f_λ have Milnor number 16 and the resolution of f_λ is shown in Figure 9. From this resolution and from the Milnor number it is not difficult (cf.[BK86]) to derive that the divisor with normal crossings $D \subset Z$ has the shape that is illustrated in Figure 10. Moreover we see $d = 156$. Notice also that $m = 4$ and $k = 6$.

Observe that the automorphism of $(\mathbb{C}^2, 0)$ which is given by

$$\phi : \begin{pmatrix}\tilde{x}\\ \tilde{y}\end{pmatrix} = \begin{pmatrix}\vartheta^2 x\\ \vartheta^3 y\end{pmatrix} \tag{5.5}$$



FIGURE 9. The resolution of $f_\lambda(x,y) = \big(y^2 - x^3\big)^2 - \big(4\lambda x^5 y + \lambda^2 x^7\big)$.

has the property that: $f_\lambda(\tilde{x},\tilde{y}) = \vartheta^{12} f_{\vartheta\lambda}(x,y)$. Or likewise, ϕ yields a commutative diagram:

$$\begin{tikzcd}
(\mathbb{C}^2,0) \arrow[r, "\phi"] \arrow[d, "f_{\vartheta\lambda}"'] & (\mathbb{C}^2,0) \arrow[d, "f_\lambda"] \\
(\mathbb{C},0) \arrow[r, "\cdot\vartheta^{12}"] & (\mathbb{C},0).
\end{tikzcd}$$

Hence, the MHS on the nearby fundamental group of the pair $(f_{\vartheta\lambda}, \frac{\partial}{\partial t})$ is isomorphic to the one of $(f_\lambda, \vartheta^{12}\frac{\partial}{\partial t})$, i. e. $\mathfrak{P}(f_{\vartheta\lambda}, \frac{\partial}{\partial t}) \cong \mathfrak{P}(f_\lambda, \vartheta^{12}\frac{\partial}{\partial t})$.

Therefore, we fix one f_λ and consider how $\mathfrak{P}(f_\lambda, \vartheta^{12}\frac{\partial}{\partial t})$ varies, when we change $\vartheta^{12} =: \theta$. We know that for any $s \geq 1$, the variation of MHS $\{\mathfrak{P}^s(f_\lambda, \theta\frac{\partial}{\partial t})\}_{\theta\in\mathbb{C}^*}$ is the nilpotent orbit of MHS associated to the pair $(\mathfrak{P}^s(f_\lambda, \theta\frac{\partial}{\partial t}), \mathcal{L})$, where $\mathcal{L} = \bigoplus_{\vec{v},\vec{w}} L$ is the nilpotent endomorphism given by $L = \frac{1}{156}N - \frac{1}{24}M$.

Denote by $h_\lambda : (Z_\lambda, D_\lambda^+) \to (\Delta, 0)$ the map that is obtained by semistable reduction from f_λ. By Theorem 5.2 we have a commutative diagram

$$\begin{tikzcd}
(Z_\lambda, D_\lambda^+) \arrow[r] \arrow[d, "h_\lambda"'] & (\mathbb{C},0) \arrow[d, "f_\lambda"] \\
(\Delta,0) \arrow[r] & (\mathbb{C},0) \\
t \arrow[r, mapsto] & t^{156}.
\end{tikzcd}$$

To h_λ we associate the dga $A^\bullet$ and note by Corollary 2.20 on page 33 that $H^1(A^\bullet)$ is a pure Hodge structure of weight 1. Since $(\vec{J}/\vec{J}^3)^*$ has weights 1 and 2 and since N and M both lower the weights by 2, we see that N and M are both zero on $(\vec{J}/\vec{J}^3)^*$. This shows that the nilpotent orbit of MHS

$$\{\mathfrak{P}^2(f_\lambda, \theta\tfrac{\partial}{\partial t})\}_{\theta\in\mathbb{C}^*}$$

is constant. Our goal now is to show:

Proposition 5.10. *The nilpotent orbit of MHS*

$$\{\mathfrak{P}^3(f_\lambda, \theta\tfrac{\partial}{\partial t})\}_{\theta\in\mathbb{C}^*}$$

is not constant.



FIGURE 10. The divisor D for $f_\lambda(x,y) = \big(y^2 - x^3\big)^2 - \big(4\lambda x^5 y + \lambda^2 x^7\big)$.

Proof: In order to prove this proposition we just have to show that $L \neq 0$. Choose $\vec{v} \in \mathcal{S}(\frac{\partial}{\partial t})$ and $\vec{w} \in \mathcal{M}(\frac{\partial}{\partial t})$. On the elliptic curve D_7, choose a point $* \in D_7$ and a closed path α based at $*$. Let $\omega^{(7)}$ be a holomorphic 1-form on D_7 such that

$$\int_\alpha \omega^{(7)} = 1.$$

There exists a $\mu^{(7)} \in E^1(D_7 \log p_{67})$ such that $\omega^{(7)} \wedge \bar{\omega}^{(7)} + d\mu^{(7)} = 0$ and $\rho := \operatorname{Res}_{p_{67}} \mu^{(7)} \neq 0$. Now let $\eta_1^{(7)}, \dots, \eta_6^{(7)}$ be meromorphic forms on $D_1, \dots, D_6$ respectively and $\eta_0^{(7)} = \rho_{01} \frac{dp}{p} \in \bigwedge^1(\frac{dp}{p})$ such that:

$$\rho_{i,i+1} = \operatorname{Res}_{p_{i,i+1}} \eta_i^{(7)} = -\operatorname{Res}_{p_{i-1,i}} \eta_i^{(7)} = -\rho.$$

Now consider the Chen-closed element in $(A^1 \otimes A^1) \oplus A^1$ that is given by:

$$\Omega = \omega^{(7)} \otimes \omega^{(7)} \otimes \bar{\omega}^{(7)} + \omega^{(7)} \otimes \Big(\mu^{(7)} + \eta_6^{(7)} + \dots + \eta_0^{(7)} + \rho_{67}\Theta + \dots + \rho_{01}\Theta\Big).$$

And note that $N(\Omega) = \omega^{(7)} \otimes (\rho_{67} d\xi + \dots + \rho_{01} d\xi)$. There is an $h \in A^0$ such that $h(p_{01}) = 0$ and $dh = \rho_{67} d\xi + \dots + \rho_{01} d\xi$. It is easy to see that for this h holds

$$[N(\Omega)] = \Big(\omega^{(7)} \big| (\rho_{67} d\xi + \dots + \rho_{01} d\xi)\Big) = (-h\omega^{(7)}) = (7\rho\omega^{(7)}) \in H^0 \bar{B}_3(A^\bullet, a).$$

Similarly we see $[M(\Omega)] = (\rho\, \omega^{(7)}) \in H^0 \bar{B}_3(A^\bullet, a)$. From this we derive:

$$[L(\Omega)] = \frac{1}{156}[N(\Omega)] - \frac{1}{24}[M(\Omega)] = (-\frac{32}{156}\rho\, \omega^{(7)}) \neq 0$$

□

Remark 5.11. Since $\mathfrak{P}^2(f_\lambda, \theta\frac{\partial}{\partial t}) = \mathfrak{P}^2(f_\lambda, \frac{\partial}{\partial t})$ for all $\theta \in \mathbb{C}^*$, we may consider the map

$$\begin{array}{ccc} \mathbb{C}^* & \rightarrow & \mathrm{Ext}_{\mathrm{MHS}}\left(\bigoplus_{\vec{v},\vec{w}} H^1(A^\bullet);\;\; \mathfrak{P}^2(f_\lambda, \frac{\partial}{\partial t})\right) \\ \theta & \mapsto & \left\{0 \rightarrow \mathfrak{P}^2(f_\lambda, \frac{\partial}{\partial t}) \rightarrow \mathfrak{P}^3(f_\lambda, \theta\frac{\partial}{\partial t}) \rightarrow \bigoplus_{\vec{v},\vec{w}} H^1(A^\bullet)^{\otimes 3}\right\}. \end{array} \tag{5.6}$$

The fact that $[\omega^{(7)}] \otimes [\omega^{(7)}] \otimes [\bar{\omega}^{(7)}] \in F^2(H^1(A^\bullet)^{\otimes 3})$ and $L(\Omega) \notin F^2H^1(A^\bullet) = 0$ can be used to prove that even the map (5.6) is not constant.

REFERENCES

[A'C73] N. A'Campo. Sur la monodromie des singularités isolées d'hypersurfaces complexes. *Invent. math.*, 20:147–169, 1973.

[A'C75] N. A'Campo. La fonction zeta d'une monodromie. *Comm. Math. Helv.*, 50:233–248, 1975.

[BK86] E. Brieskorn and H. Knörrer. *Plane Algebraic Curves.* Birkhäuser, Basel, Boston, Stuttgart, 1986.

[BK91] D. Bättig and H. Knörrer. *Singularitäten.* Lectures in Mathematics, ETH Zürich. Birkhäuser, Basel, Boston, Stuttgart, 1991.

[BS74] J. Briançon and H. Skoda. Sur la clôture intégrale d'un idéal de germes de fonctions holomorphes en un point de $\mathbb{C}^n$. *C. R. Acad. Sc. Paris*, t. 278:949–951, 1974.

[Che73] K. T. Chen. Iterated integrals of differential forms and loop space homology. *Ann. of Math.*, 97(2):217–246, 1973.

[Che75a] K. T. Chen. Connections, holonomy and path space homology. In *Differential Geometry*, volume 27 of *Proc. Sympos. Pure Math.,*, pages 39–52, Providence, R. I., 1975. Amer. Math. Soc. Part I.

[Che75b] K. T. Chen. Iterated integrals, fundamental groups and covering spaces. *Trans. Amer. Math. Soc.*, 206:83–98, 1975.

[Che76] K. T. Chen. Reduced bar constructions on de Rham complexes. In A. Heller and (eds.) M. Tierny, editors, *Algebra, Topology and Category Theory*, pages 19–32, New York, 1976. Academic Press.

[Che77a] K. T. Chen. Extensions of C^∞ function algebra by integrals and Malcev completion of π_1. *Adv. in Math.*, 23:181–210, 1977.

[Che77b] K. T. Chen. Iterated path integrals. *Bull. Amer. Math. Soc.*, 83:831–879, 1977.

[CK82] E. Cattani and A. Kaplan. Polarized mixed Hodge structures and the local monodromy of a variation of Hodge structure. *Invent. math.*, 67:101–115, 1982.

[Cou34] R. Courant. *Differential and Integral Calculus*, volume 1. Blackie & Son Limited, London and Glasgow, 1934.

[Del71] P. Deligne. Théorie de Hodge, II. *Publ. Math. IHES*, 40:5–58, 1971.

[GS75] P. Griffiths and W. Schmid. Recent developments in Hodge theory: A discussion of techniques and results. In *Proc. Bombay Colloq. on Discrete Subgroups of Lie Groups*, pages 31–127, Bombay, 1975. Oxford Univ. Press.

[Hai87a] R. M. Hain. The de Rham homotopy theory of complex algebraic varieties I. *K-Theory*, 1:271–324, 1987.

[Hai87b] R. M. Hain. The de Rham homotopy theory of complex algebraic varieties II. *K-Theory*, 1:481–497, 1987.

[Hai87c] R. M. Hain. The geometry of the mixed Hodge structure on the fundamental group. *Algebraic Geometry, Bowdoin, PSPM*, 46:247–282, 1987.

[Ill94] L. Illusie. Logarithmic Spaces according to K. Kato. In V. Cristante and W. Messing, editor, *Barsotti Symposium in Algebraic Geometry*, pages 183–203. Perspectives in Math. 15, Academic Press, 1994.

[Jab86] E. R. Jablow. Quadratic vector classes on Riemann surfaces. *Duke Math. J.*, 53(1):221–232, 1986.

[Kae97] R. H. Kaenders. *On De Rham Homotopy Theory of Plane Algebraic Curves and their Singularities.* PhD thesis, defended at the Catholic University of Nijmegen, 1997. http://www.math.ruu.nl/people/kaenders/ or http://www.pica.nl/frames/index.html.

[Kat89] K. Kato. Logarithmic Structures of Fontaine-Illusie. In J. I. Igusa, editor, *Algebraic Analysis, Geometry an Number Theory*, pages 191–224, Baltimore, 1989. Johns Hopkins University Press.

[KN95] K. Kato and C. Nakayama. Log Betti Cohomology, Log Étale Cohomology and Log De Rham Cohomology of Log Schemes over $\mathbb{C}$. *Prepublications, Univ. Tokyo, UTMS 95-16*, 1995.

[Lam75] K. Lamotke. Die Homologie isolierter Singularitäten. *Math. Zeitschrift*, 143:27–44, 1975.

[Lê72] D. T. Lê. Sur les nœuds algebriques. *Compos. Math.*, 25(Fasc. 3):281–321, 1972.

[LT81] J. Lipman and B. Teissier. Pseudo-rational local rings and a theorem of Briançon-Skoda about integral closures of ideals. *Mich. Math. J.*, 28:97–116, 1981.

[Mor78] J. W. Morgan. The algebraic topology of smooth algebraic varieties. *Publ. Math. IHES*, 48:137–204, 1978.

[Sch73] W. Schmid. Variation of Hodge Structure: The Singularities of the Period Mapping. *Invent. math.*, 22:211–319, 1973.

[Spa66] E. H. Spanier. *Algebraic Topology.* McGraw-Hill, New York, 1966.

[Ste76] J. H. M. Steenbrink. Limits of Hodge Structures. *Invent. math.*, 31:229–257, 1976.

[Ste77] J. H. M. Steenbrink. Mixed Hodge structure on the vanishing cohomology. In P. Holm, editor, *Real and Complex Singularities, Oslo 1976*, pages 525–563, Alphen a/d Rhijn, 1977. Sijthoff & Noordhoff.

[Ste95] J. H. M. Steenbrink. Logarithmic Embeddings of Varieties with Normal Crossings. *Math. Ann.*, 301:105–301, 1995.

[SZ85] J. H. M. Steenbrink and Steven Zucker. Variation of Mixed Hodge Structure. *Invent. math.*, 80:489–542, 1985.

[tD91] T. tom Dieck. *Topologie.* de Gruyter, Berlin, 1991.

[Usu96] S. Usui. Recovery of Vanishing Cycles by Log Geometry. *Preprint*, 1996. Dept. of Math. , Osaka University.

Department of Mathematics, Utrecht University, Postbus 80.010, 3508 TA Utrecht, The Netherlands

E-mail address: `kaenders@math.ruu.nl`